to the same risks arising from unhealthy environment; and, again, if phthisis were directly contagious, it is remarkable that it should be chiefly contagious in badly ventilated houses, and on certain soils; and it is further difficult to see why drainage of the land and elevation above the sea level should in any way affect its contagiousness. Dr. Dewar, of Arbroath, makes the significant remark that in all his cases of supposed infection (No. 166), "the patients lived in small confined houses, and slept in the 'box beds' in use in Scotland." "During twenty-five years," he says, "I have not seen one case of contagion in the airy houses of the well-to-do."

2. As to the dissemination of phthisis by means of dust, it is well known that Koch regards this as the most common way in which it is produced—by the drying up and pulverisation of matter expectorated upon the ground, on the floors of dwellings, or on handkerchiefs.

Long before the discovery of the bacillus of tubercle, also, this view was taken by some of the most competent observers; thus Dr. Farr, in his Report to the Army Commission, remarked that "the prevalence of phthisis in the armies of Europe is probably due in part to the inhalation of expectorated tubercular matter, dried, broken up into dust, and floating in the air of close barracks."

The retention of the bacillus either in dust or in the vapour of the air would fully account for most of the cases of the supposed direct infection recorded by the Collective Investigation Committee, and especially for those in which the virus appeared to be introduced from without into houses previously entirely free from the disease (see Cases No. 188, 194, 196, and 255). It accounts fully for all the additional cases mentioned by Dr. Burney Yeo in his able defence of the doctrine of contagion in the *JOURNAL*, April 18th, 1885.

3. We have now reached, perhaps, the most difficult part of our inquiry, as to whether any conditions outside the mere presence of the bacillus are necessary, or at any rate auxiliary, to its infective power.

(1). With regard to the influence of free ventilation, it is not only possible, but highly probable, that in many of the places where its good effects have been observed, other conditions as well have come into operation, and especially greater cleanliness and more complete removal of dust and other refuse. Where people care for fresh air, they also frequently show a preference for cleanly surroundings, both of house, and clothes, and person.

(2). Somewhat similar remarks may apply to the influence of an elevated site upon even indirect infection through dust, for in these situations we probably have both more movement of air and less dust.

(3). But no such explanations as these will touch the remaining two points that we have to consider. If there is no condition outside the body that influences the viability or the virulence of the bacillus, it is very difficult to account for the comparative immunity of frosty or sub-arctic regions, and for the intensity of the tubercular action in hot or tropical climates. It is certain that chronic inflammations of the lungs are not more common in the latter, and thus one source of the disease is not so frequently present; whilst in the cold regions, where these affections are rife, phthisis is comparatively rare and much less virulent. We have no reason to suppose that bacillus-laden dust is absent in the crowded towns and close rooms which are to be found in Canada, for instance, and yet the disease does not spread as it does in the warmer and more genial air of Italy and the tropics.

(4). Again, it affords no explanation of the influence upon such infection of damp or undrained subsoils, nor yet of the improvement in the phthisis rate that has been shown to follow thorough drainage of a town.

To account for these several points, it is necessary to assume the existence of some condition necessary to infection by the bacillus, which shall either enable it to live longer outside the body, or which may even increase its power of attack.

I have ventured to suggest that the tubercle bacillus, like some other infective microzymes, may actually take up an increase of virulence by a sojourn for a time in some medium external to the body, either in polluted ground air or in an atmosphere saturated with aqueous vapour from the lungs.

I do not know whether the same observation has been made by other physicians to consumption hospitals, but it has certainly struck me that a fresh infection of the lung in persons already suffering from phthisis is much more likely to take place from without than from within the body. Many times have I seen in-patients, with sputum teeming with bacilli, improve in health, gain weight, and almost lose the physical signs of their disease; and then they

have returned home, and in a few weeks they have come back to the hospital, sometimes with fresh disease in the damaged lung, sometimes in the opposite lung that had been previously healthy, and sometimes with laryngeal phthisis.

Such facts as these certainly seem to show that there was something in the air of their own homes much more infective than the bacilli of which they were themselves the hosts. The one fact that might seem to bear against this theory is the high temperature required for the development of the organism; and the greater virulence of the disease in hot climates probably shows the influence of this condition, but it is by no means proved that in cold climates the bacilli contained in damp and foul dwellings may not in some way increase their power for evil.⁷

CASE OF CÆSAREAN SECTION FOR OSTEO-MALACIC PELVIS.

By WM. MCGOWAN, M.D.,

Medical Officer Maghera Dispensary, co. Derry

ON December 4th, 1889, I was called upon to attend in labour I. McC., wife of a farm labourer. I saw her at her residence about two miles distant, at 7.30 A.M. The patient, aged 37, a small, delicate, emaciated woman, was mother of five children, and all her previous labours had been natural and easy. I myself attended her at her last confinement on August 17th, 1881, when the pelvis was quite natural and roomy, and she had an easy labour, not even the forceps being required. After this I lost sight of her, but there was the history of her having, immediately following this confinement, to keep her bed for over six months from what was called "rheumatic pains." For some six months or so more she was only able to be carried to a chair at the fire. After this she began to go about a little on crutches, and gradually gained strength, but remained permanently "bent in the back," "sunk," and "stooped." During the latter months of gestation, she said her health had been very bad, and scarcely anything was retained on the stomach, so that she was very weak. When I saw her she had been about forty-two hours in labour, and yet, as she said, "it was doing no good; the lump had not gone down, as it used to do." Examination revealed an extreme degree of pelvic contraction and deformity. The tubera ischia were less than three-quarters of an inch apart, almost obliterating the pubic arch; so closely approximated were they that the index finger could only with some difficulty pass between them. The sacrum and coccyx projected forwards, leaving only a narrow chink between them and the approximated ischia. For want of space, caused by this narrowing of the outlet, it was impossible to make out the presentation. As delivery by the natural passage seemed to me an impossibility, I sent for my friend Dr. Thompson, of Bellaghy, who arrived shortly after 10 o'clock. On examining her, he agreed with me that the only chance of delivery must be the Cæsaean section. The nature and dangers of the operation were explained, and the sanction of the patient and her husband having been unhesitatingly obtained, we determined to lose no time in carrying out the operation, as the powers of Nature were already greatly exhausted by the prolonged labour. Giving directions to have the house thoroughly heated and other preparations made, I returned home for the necessary instruments, etc., and proceeded again to the house at 12 o'clock, accompanied by Dr. Thompson and Dr. Hueston, of Maghera, who kindly afforded me their valuable advice and assistance. The patient was lying in the kitchen of a farm labourer's cottage, so that the surroundings were not favourable for a serious operation. However, there was no alternative. The rectum was well cleared out by an enema, a catheter passed, and the patient placed on a suitable table, and put under the influence of chloroform. With the strictest antiseptic precautions possible under the circumstances, the usual incision was made, and without any difficulty I speedily removed a living child, healthy, and fairly well developed. In making the incision in the uterine wall, I came on the placenta attached to the anterior surface. This I cut through, and the child and placenta were removed without any trouble from hæmorrhage. Owing probably

⁷ The case against infection by sputum dust only without some increase of virulence from external conditions is well argued by Dr. Candler in his work on the *Prevention of Consumption* (Kegan Paul and Co.). Many of Dr. Andrew's observations in his *Lumleian Lectures* point in the same direction.

to exhaustion caused by the prolonged labour, the uterus did not contract well. After ascertaining that the passage to the vagina was patent, the uterus was sutured by chromicised catgut passed deeply into the tissue. The peritoneal cavity was sponged out, and the wound in the abdominal wall secured by six silk sutures passed deeply, so as to include the whole thickness of the wall and peritoneum, and superficial ones through the skin between these. The wound was dusted with iodoform, covered with wood-wool, and secured by four broad strips of adhesive plaster and a flannel binder over all. The external genitals were covered with carbolic tow and cotton wool. During the suturing of the external wound, vomiting set in from the chloroform, but it soon ceased, and the patient was put to bed, with a pulse, though weak, as good as before the operation. A hot water jar was put to her feet, her head covered with flannel, and a hypodermic injection of morphine given. In the evening I found her in a satisfactory condition, but retching and vomiting at intervals. Ordered nothing to be given by the mouth except small quantities of brandy and iced soda-water. Passed the catheter.

December 5th. Comfortable, but retching a little; passed catheter. In the evening she complained of rather severe pain in the right side of the abdomen with accelerated breathing; pulse 136, temperature 99.4°. Gave a quarter of a grain of morphine hypodermically, passed catheter, and syringed out the vagina with carbolic acid lotion. One grain of opium to be given every four hours until pain ceased.

December 6th. Found her much better, having slept after the morphine; pain almost gone; expresses herself as feeling comfortable; had passed urine voluntarily.

For the next two days her condition was fairly satisfactory; scarcely any pain; temperature never higher than 99.4°, but pulse continued to increase in rapidity and weakness. During the entire treatment, nourishment was given by means of enemata consisting of beef-tea, brandy, egg, Bengel's food, etc., which were well retained. The vagina was syringed out regularly, and the discharge remained healthy.

On the evening of the 8th, I found her worse. Delirium and restlessness had set in and continued until her death, which occurred at 6 o'clock P.M. on December 9th, the sixth day after operation.

On removing the dressings after death, I found the wound in the abdomen healed, and not a drop of pus was observed. There was gaping between two of the sutures, but this was evidently caused by her restlessness before death, when she could only with difficulty be restrained from tearing off the dressings. I got permission only to reopen the abdominal wound, so that the examination was very limited, and it was also, under the circumstances, hurried. I found the stomach and intestines much distended with flatus. There was marked peritonitis with adhesions, which probably arose on the day after the operation. The uterus was forced down by the distended intestines, was well contracted, and there appeared complete union of the uterine wound. There was no blood or effusion in the abdominal cavity.

REMARKS.—As she got over the period of collapse, and escaped the second great danger (namely, septicæmia), hopes were entertained of recovery. The cause of death was peritonitis, and I may call attention to the obscurity or "latency" of the symptoms. Except for two or three hours she scarcely suffered any pain; her temperature was never above 99.4°. The most marked symptom was the accelerated pulse, gradually increasing in rapidity and weakness, which might have been an indication of exhaustion or prolonged collapse, independent of peritonitis, taking into consideration her enfeebled condition at the time of operation. Notwithstanding the unfavourable result to the mother, the case seems to me of interest not on account of anything novel in the performance of the operation, or, unfortunately, in the result, but from the nature and rarity of the deformity necessitating the operation.

There can be no doubt of this being a case of *mollities ossium*. The pelvis was characteristic of this deformity and contraction. The nature of the outlet has been already described, and from the slight *post-mortem* examination made I found irregularity and deformity of the brim corresponding to the outlet, although not to such an extreme extent. The iliac and pubic bones were pressed inwards, narrowing the cavity of the pelvis transversely, and the vertebral column had sunk into the pelvis, and overhung the brim.

I regret very much not having had the advantage of a complete *post-mortem* examination so as to have taken measurements.

The case is of interest in several respects. 1. The rarity of the disease in this country. In the north of Ireland, at least, I believe it to be exceedingly rare. 2. The extreme degree of the deformity; and, 3, the complete ossification of the previously softened bones, and, as a consequence, the absolute necessity of the operation performed. Spiegelberg says, "It is extremely rare for complete recovery to take place, or for the pelvis again to become ossified." Barnes, in describing his new method of embryotomy designed "to effect delivery in extreme cases of pelvic contraction," where Cæsarean section was previously considered the only resource, and defining the cases suitable for it, says: "Where two fingers can barely pass between the tuberosities of the ischia, there will be insufficient room for manipulation, and it will be scarcely possible to guide the *écraseur* through the pelvis." "But in these cases," he adds, "the bones will often open up under pressure applied from within." Again, Spiegelberg says: "That most of such pelves are dilatable, and labours *per vias naturales* have become more common. Casati asserts that he has almost always found the osteo-malacic pelvis (which is so common in Milan) to be dilatable, and has extremely rarely been obliged to resort to Cæsarean section, only in two out of forty-one cases." These quotations show how rare it is for complete ossification to take place. Even if the head had entered the pelvis in this case, which it could not do, delivery would have been equally impossible, as the bones at the outlet were absolutely rigid and unyielding. This was easily tested on account of the close approximation of the ischia, where the slightest yielding could have been detected. No doubt the complete consolidation of the bones was promoted by the long freedom from child bearing. The morbid process which deprived the bones of their calcareous matter had ceased to progress, and they, in the course of time, had become re-ossified.

As regards the etiology of the disease, about which so little is known, I may mention that my patient attributed her illness to a damp, cold, and badly lighted house. So convinced was she of this fact that, as soon as she was able, she removed to another and drier house, after which she stated she improved much more rapidly. No doubt this may have been the exciting cause, under the lowering influences of the puerperal state, in a constitution delicate at the best.

ON THE HISTOLOGY OF A CASE OF SO-CALLED "DUCT CANCER."

By ARTHUR E. BARKER, F.R.C.S.,
Surgeon to University College Hospital, etc.

THE patient from whose breast I have made these microscopic sections and drawings was a lady, aged 52, of healthy appearance but spare figure. Her attention was first drawn to her right breast some three years before she first came under observation, by a slight serous discharge from the nipple; this had continued ever since. For about a week before I first saw her, on February 4th, 1889, this discharge had been tinged with blood. On examination of the breast I found it very much atrophied and scantily furnished with fat. The nipple was puckered at one side and hard at the base, but little if at all retracted. The skin over the breast, was in no way affected except by a trace of puckering at one spot on being pinched. There was a distinct hard nodule the size of a filbert, just above and to the inside of the base of the nipple, and another in the substance of the breast close to it. Manipulation of these knots caused an escape of bloody serum from the nipple. This fluid contained many large round corpuscles undergoing fatty change and also red corpuscles (Plate I, c & d). The former (c) were many times larger than the latter (d). There had been no pain in the breast until recently, and at the moment of examination it was said to be quite absent. The glands in the axilla were markedly knotty, but not enlarged, and were freely movable.

There was no history of abscess in the breast, nor of any injury except a severe bite of the nipple twenty-two years ago during nursing. The part is described as having been nearly bitten off, and the irregularity at its base has always been attributed, and probably correctly, to this accident.

In view of all the facts of the case, I concluded that this was an instance of so-called "duct cancer," and thus described it to the patient's son who was the medical man who had brought the patient for consultation.