was specially impaired; and in all a total absence of paralysis of the limbs.

Though I have occupied so much of your time, I have only been able to treat of—and that in many respects very imperfectly—the functions of the cortical centres so far as concerns sensation and motion. There is another question which I have not considered at all, namely, the relations of the cerebral hemispheres to the functions of organic life. This, however, is a subject which is still involved in so much obscurity, and in reference to which there are at present so few facts which are not susceptible of different modes of interpretation, that I think it well to wait for further light before hazarding any definite views of my own. And I feel it all the less necessary to do so, seeing that the subject in one of its principal aspects, namely, the relation of the hemispheres to the thermic functions of the body, has been recently so ably placed before you by my predecessor, Dr. MacAlister.

On the psychical aspects of cerebral localisation I have touched only incidentally. This of itself would require a volume, and that mainly of speculation. As to the questions which I have treated more fully, and on which so many differences of opinion at present exist, and will probably still continue, I shall be content if the facts and considerations which I have brought before you contribute to their solution, if only by stimulating work on the part of others, with a view to arriving at conclusions which shall be acceptable alike to physiologists and physicians. For the true conception of the functions and relations of the cerebral hemispheres and their constituent centres is not only of the highest scientific and philosophical interest, but of direct and important practical bearing on the diagnosis and treatment of cerebral disease.

THREE LECTURES ON SOME POINTS RELATING TO INJURIES TO THE HEAD.

Delivered at the Royal College of Surgeons of England.

By WILLIAM HENRY BATTLE, F.R.C.S.,

Hunterian Professor of Surgery and Pathology, Royal College of Surgeons
Assistant-Surgeon to the Royal Free Hospital; and to the East London
Hospital for Children; formerly Resident Assistant-Surgeon at, and
Surgical Registrar to, St. Thomas's Hospital.

LECTURE II.

MR. PRESIDENT AND GENTLEMEN,—I would now direct your attention to other evidences of fracture of the base of the skull, the escape of blood from the seat of fracture, and its appearance under the conjunctiva or over the mastoid process. An effusion of blood into the cellular tissue of the orbit is one sign of the presence of a fracture of the anterior fossa of the base of the skull. This effusion of blood may be limited to the eyelids at first, and then spread to the ocular conjunctiva; it may take place into the back of the orbit, and then spread forwards, only showing itself in the conjunctiva after the lapse of some hours or days, and then spread to the lids; it may be small in quantity, or sufficient to produce exophthalmos. The value of orbital extravasation as a sign of fracture is largely dependent on the question of its position (subconjunctival or not), and the quantity in which it presents. Other signs, such as epistaxis from the nostril of the one side on which the fracture is, and the position of the injured part must be considered. Most are familiar with the appearance of the eye when there is this hæmorrhage under the ocular conjunctiva; in extreme cases it forms an elevated brownish-red circle, which encloses the cornea, and contrasts strongly with the colour of the iris of the patient. Usually disappearing in the course of a fortnight or so, it may last a long time, the blood not being absorbed for weeks; I have watched its gradual disappearance over a period of from six to seven weeks. In four cases, all of which recovered, there was much protrusion of the eye on admission, and that to an extent sufficient to interfere with the action of the ocular muscles. In one case, that of a man of 25, there was slight proptosis on admission; next day there was more marked but still moderate protrusion of the eyeball, which was displaced downwards and towards the middle

line, the movements being limited in every direction, chiefly outwards. There was no ophthalmoscopic change. On the third day there was considerable subconjunctival ecchymosis. In sixteen patients there was orbital effusion in the lids when they came in, which subsequently spread to the conjunctiva. In four subconjunctival ecchymosis appeared on the second day, in three on the third, and in one on the sixth. Of the fatal cases, ecchymosis of the conjunctiva was not an invariable symptom, for in two out of fourteen there was none, the fracture being correctly diagnosed by the severe epistaxis, and in one, where the lesser wing of the sphenoid was splintered, the hæmorrhage did not show in the conjunctiva for at least ten hours after the injury. Again, in the case in which the anterior clinoid process was the only part of the base broken, contusion of the lids and extravasation in the conjunctiva, were evident from the first.

In fractures of the malar and superior maxillary bones blood may be extravasated into the lids and under the conjunctiva; this is a rare occurrence which might lead to error in diagnosis. Mr. Holmes¹ has recorded a case. Another sign of fracture of the base of the skull is the appearance of extravasated blood in the mastoid region after an injury to the head. I must ask you to consider this sign somewhat fully, for I consider it under certain circumstances to be a most important indication that the posterior fossa of the skull is the seat of the fracture. Sir Prescott Hewett writes: "Extravasation of blood, and consequent discoloration of the skin, appearing in the mastoid region some hours after a severe injury to the head, may lead to the suspicion of a fracture involving the posterior part of the base; and all the more valuable will this sign become if the injury did not bear directly upon this region, and especially if it bore upon the opposite side of the head."² Observations on this subject lead me to the following conclusions with regard to "mastoid ecchymosis." That it appears in the first place in front of the apex of the mastoid process. That it often spreads upwards over the mastoid in a line, slightly curved, and with the convexity backwards, its direction being approximately that of the outline of the external ear, from which it is distant half to three-quarters of an inch. At the end of three or four days after its appearance it diffuses itself forwards and backwards, chiefly in the latter direction, is most marked in the original line, and then gradually disappears. That it usually shows from the third to the fourth day after the injury, but its appearance may be delayed until the twelfth or fourteenth day. If the injury which is in the occipital region is to the right of the middle line, the ecchymosis will appear on that side over the anterior part of the apex of the mastoid process. That it may be accompanied with œdema and tenderness over the process. That the duration of the time which elapses between the receipt of the injury and the appearance of the extravasation is to some extent an indication of the distance of the fracture from the process, the blood being compelled to travel by a certain anatomical route in order to reach the surface. That unless search be made for the extravasation it is very apt to be overlooked, as the ear conceals it, especially if the ear is large and the head of patient has not been shaved. Should the fracture have taken place in the middle line of the occipital bone, or should there be two or more lines of fracture diverging towards the foramen magnum from a point in the middle line the extravasation may appear on both sides and at a varying interval.

The following examples illustrate these statements. A man, aged 41, fell and received a scalp wound over the posterior part of the left parietal bone. He was admitted into hospital unconscious, bleeding freely from the ears and nose. On the fourth day there was ecchymosis behind the left ear, from which flowed clear watery fluid. A woman, aged 50, knocked down by a hansom cab, had considerable hæmorrhage from the right ear, which ceased during the first night. On the fourth day there was well-marked ecchymosis over the right mastoid process. A man, aged 24, thrown from the driving seat of a van, was admitted into the wards unconscious, with very free hæmorrhage from the right ear. On the third day there was an extravasation of blood over the mastoid continuous in a less degree down the side of the neck. Over this area there was much tenderness. This extravasation disappeared in sixteen days, but the tenderness remained for some time longer. A man, aged 40, was knocked down, and struck the back of his head on the pavement. On the sixth day there was considerable discoloration over the occipital bone and behind the ear, and he threw his arms about when touched in that situation. A man, aged 32, was admitted on

¹ JOURNAL, 1885, p. 967.

Holmes, *A System of Surgery*, edited by Hulke, vol. i, p. 591.

the fifth day after a fall from a loft, with marked ecchymosis in the mastoid region, and a small scalp wound in the posterior parietal region on the same side. A youth, aged 17, fell fifteen feet, and was found to be unconscious, and with a jagged wound of the scalp on the right side. On the fifth day confusion was seen behind the right ear. A man, aged 47, was picked up insensible after having been run over in the street. On the seventh day there was premastoid ecchymosis on the left side, but it was not until the fourteenth day that it appeared in front of the right mastoid process. A man, aged 37, received a blow from a man's fist, and fell on the back of his head on the pavement, and was brought to the hospital on the third day. There was then ecchymosis over the right mastoid process, which, with other symptoms, was considered to indicate a fracture in the posterior fossa on the right side. A year later he was admitted under the care of Dr. Bristowe with epilepsy, and died. At the *post-mortem* examination the diagnosis was proved to be correct. A man, aged 47, fell seven feet from a wall, producing a superficial scalp wound just behind the right parietal. On the third day some ecchymosis appeared over the right mastoid process. A boy aged 7 fell down a stairway, receiving a small wound in the left occipital region. On the seventh day some ecchymosis appeared over the left mastoid process. A boy aged 12 fell backwards off a brick wall about five feet in height on concrete pavement. A hæmatoma was found in the left occipital region. On the twelfth day there appeared some bruising over the left mastoid, especially in the region of the apex of that process. A man aged 21 fell off a cart, his head striking the ground, and was admitted with hæmorrhage from the ear. There was no wound of the scalp; but on the fourth day, when his head was shaved, bruising over the back of the head was found, and on the fifth day ecchymosis over the right mastoid process, which appeared at the very tip in the first instance. A boy aged 7, who had fallen down some stone steps, presented ecchymosis behind the left ear on the third morning. A man aged 31 fell seven feet, and complained of tenderness over the mastoid process the same evening; this continued, but it was only on the third morning that ecchymosis appeared over the right mastoid process, just behind the ear, with a little puffiness over the process. Eight days later a note was made to the effect that the bruising over the right mastoid process which followed the outline of the external ear was beginning to fade, though the staining over its anterior border was still well marked, and there was slight œdema on deep pressure, but no longer definite tenderness. A woman aged 55 was knocked down in the street by a cab. On the fifth day there was a well-marked swelling about the position of the posterior superior angle of the left parietal, and ecchymosis was noticed over the left mastoid process. This discolouration was visible for a period of four weeks, and when it had disappeared there was still some tenderness. A man aged 37 fell backwards from a cart on a macadamised road. There was hæmorrhage from the nose, mouth, and left ear. On the fourth day there was ecchymosis over the mastoid, commencing to extend backwards. A policeman aged 44 was thrown violently from a light cart on a stone pavement. There was bleeding from the nose and both ears, which continued for some hours. On the third day there was bruising visible over the left mastoid process.

A summary of these cases gives the following result:—that when, in patients with supposed fracture of the posterior fossa, ecchymosis was first perceived, it was on the third day in six cases, on the fourth day in three, on the fifth day in four, on the sixth day in one, on the seventh day in two, and on the twelfth day in one, whilst in one case it appeared also on the opposite side on the fourteenth day. I am careful to say perceived, for in more than one instance it had probably been overlooked for a time. I have already drawn attention to one case in which a necropsy proved that there had been a fracture of the posterior fossa, of which this extravasation had been the only external sign. What is the evidence afforded by the examination of the fatal cases in which mastoid ecchymosis was observed during life?

An omnibus driver, aged 62, who had fallen on his head on the kerb when mounting to the driver's seat, died nine days later from delirium tremens. On the fifth day ecchymosis was noticed over the right mastoid process. At the *post-mortem* examination Dr. Sharkey found two fractures running from the occipital protuberance obliquely downwards on the right and left. That on the right side was far the most severe, and ran across the groove of the lateral sinus, and across the right petrous bone and middle ear.

The fracture on the left only amounted to a crack, which did not allow of movement of the bones on each side of it, and stopped short of the temporal bone.

A boy, aged one year and nine months, who presented, in addition to other symptoms, bruising over the tip of the left mastoid process on the third day, died fourteen days after the injury, and Dr. Sharkey reported as a result of the examination:

"The left occipital bone was divided by a fracture which ran from the margin of the foramen magnum along the left side to its junction with the parietal; it did not quite go through the margin of the foramen magnum," etc.

The fracture on the left side in the omnibus driver whose case is alluded to above did not give rise to extravasation over the mastoid process, although he lived nine days; it only amounted to a mere crack. That it is necessary for the production of this sign that there should be a complete fracture, one penetrating both tables, was also shown in a patient aged 40, who died from meningitis after an injury to the posterior part of the skull. He was supposed from other symptoms to have received a fracture of the occipital bone, but no mastoid extravasation developed.

At the *post-mortem* examination there was a fissured fracture extending across the left posterior fossa to the middle fossa of the same side, but it only involved the inner table of the bone. The reason why ecchymosis should appear in the situation of the mastoid apex is capable of an anatomical explanation. When a fracture extends through the occipital bone towards the foramen magnum there is an extravasation of blood along the line of fracture usually between the dura mater and the bone, as well as into the muscles and intermuscular planes across which it runs. This extravasation, which varies in quantity, gradually makes its way along the lines of least resistance; in this instance the planes of cellular tissue between the muscles towards the mastoid process, aided in its course by the tonic contraction of the muscles of the part, by movements of the head, and, to some extent, by the pressure of the neck on the pillow. The difficulties presented by a direct route towards the surface in this region are very great, the structures are dense, the muscles overlap to a marked extent, and are sometimes tendinous in origin. The valvular overlapping of the muscles in this part of the body is illustrated in Fig. 4, and by preparations in the Museum

Fig. 4.—Illustrating the attachment of the muscles in the occipital region. From a preparation in the Anatomical Department of St. Thomas's Hospital. 1. Trapezius muscle. 2. Complexus. 3. Rectus capitis posterior major and minor. 4. Deep cervical fascia. 5. Sterno-mastoid. 6. Splenius capitis. 7. Trachelo-mastoid. 8. Digastric. 9. Obliquus superior. 10. Styloid process. 11. Glenoid fossa. 12. Occipital condyle.

of the College. Any pressure from within would only make the route to the surface less practicable, excepting the way round the mastoid process. The muscles are covered in by the firm investment of the deep cervical fascia. This, as you will remember, is attached along the superior curved line of the occipital bone from the external occipital protuberance behind to the posterior margin of the sterno-mastoid in front, where it divides, and after enclos-

ing that muscle, the two investments unite again in front of it. The superficial layer passing forwards forms the parotid and masticatory fascia, and then is attached to the zygoma. A prolongation of the deep layer passes behind and to the inner side of the parotid gland to the styloid process. Ascending under cover of the parotid gland to the groove between the cartilage of the ear and the mastoid process is the posterior auricular artery, and it appears to me that it is along this vessel that the extravasation makes its way, probably aided by its pulsation. This would account for the appearance of the ecchymosis in front of the apex of the mastoid process and the course which it takes afterwards over the process.

Mr. E. C. Stabb, the senior resident medical officer at the Royal Free Hospital, has, with me and in my absence, kindly made a series of experiments on the dead subject to test the truth of these statements. The object was to find out the direction in which a fluid would spread when injected amongst the muscles in the occipital region. The fluid used was a solution of Berlin blue with glycerine and water, and the injection was made by means of a syringe and cannula. In order to make the condition more closely approximate that which obtains after a recent linear fracture of the posterior fossa, the cannula was passed towards the foramen magnum and the injection forced in as the instrument was withdrawn; in all cases the puncture was closed, and in some pressure was made over the occipital region by means of an elastic bandage, which was removed several hours afterwards when the region was examined. In six cases (aged 11 months, 18, 36, 39, 44, and 66 years) the solution was injected deeply about an inch and a half midway between the external occipital protuberance and the mastoid process. In all these experiments the tendency of the fluid was to pass forwards and downwards in the inter-muscular planes, reaching beneath the sterno-mastoid muscle as far as its anterior border, and in several cases further forwards along the digastric and stylo-hyoid muscles. In one instance, that of the child aged 11 months, it passed nearly down to the clavicle beneath the sterno-mastoid. In no case did it pass through the deep fascia. In four cases (aged 19, 33, 59, and 69) the solution was injected about half to three-quarters of an inch behind the mastoid process. In a very emaciated subject, aged 69, the fluid appeared superficial to the deep fascia just below and in front of the apex of the mastoid process, but not in any quantity, and, so far as could be seen, not in connection with any vessel. The deep fascia was very thin. A better marked result was obtained in the man aged 19, where the fluid was found distinctly passing through the deep fascia and running upwards with the posterior auricular vessels. It had also passed forwards beneath the parotid gland. These experiments, then, go far to confirm my theory as to the route taken by extravasations in the suboccipital region in their progress to the surface, especially when we consider the altered conditions of the tissues which exist after death, the different character of the fluid, the greater resistance to diffusion of fluids, and the shorter time necessarily allowed for the diffusion. The probability is, then, that as the blood extravasated about the fracture is dispersed, it is arrested in its progress across the middle line by the dense fibrous structures met with there, but is gradually forced outwards to the submastoid region, and continuing its course meets with the posterior auricular artery which guides it to the surface. The reason why the occipital artery which is encountered earlier by some of the extravasated blood making its way towards the mastoid process does not guide the extravasation through the fascia and muscles to the surface is probably explained by its oblique course with regard to the muscles, any increase of tension under the fascia of the region making the muscles act in a valve-like manner and pressing them against the vessel.

Extravasation of blood into the neck is said to be an occasional cause of a discoloration of the integuments over that region, and an aid to diagnosis in cases of fractured base; but, as in the case of the "black eye," is more frequently the result of direct injury and bruising of the part where it manifests itself; and it must be borne in mind that ecchymosis about the mastoid process will occasionally be met with under similar circumstances. I was recently asked to see a case of supposed fracture of the posterior fossa of the base of the skull in which this was present; there was, however, no difficulty in discriminating between the two conditions, for the discoloration had appeared soon after the infliction of the injury, was not limited to the mastoid region or more intense in that part, and was apparent also over the ear and temporal region.

ON THE OCCURRENCE OF OPTIC NEURITIS IN HEAD INJURIES.

For many years it has been known that injuries to the head are occasionally followed by optic neuritis, and isolated evidences of this fact are to be met with in our literature; but no systematic or continued investigation was on record in this country before the communications of Mr. Edmunds and Mr. Lawford to the Ophthalmological Society on that subject. These gentlemen have investigated this question from a clinical as well as from a pathological aspect, and in their last paper, which was read in July, 1887, the result of their investigations in twenty-four cases is given, with the conclusions to which they have come.³ I do not think it necessary for our present purpose to consider the cases brought forward by them in which optic neuritis was not found, either on ophthalmoscopic examination during life or on microscopical examination of the optic nerves after their removal from the body after death. Frequently during the progress of a patient towards recovery after an injury to the head—often, too, when the tendency was to a fatal termination—repeated examinations with the ophthalmoscope have revealed no change in the condition of the discs, and this statement applies to all kinds of head injury from slight concussion to compound comminuted fractures of the skull with wound of the brain substance. Then, again, it is well to exclude those in which injury to the frontal region was followed by signs of inflammation in the eye of the corresponding side—inflammation which not infrequently passed on to atrophy of the disc and permanent blindness, as cases which I will mention later prove. Two of the cases given by Messrs. Edmunds and Lawford⁴ come under this heading, and twelve under the former. We have, then, ten remaining from their series for consideration, and to these I add seventeen which have been observed by me since they made their communication to the Ophthalmological Society. It would perhaps be as well to mention here that their investigations were made in cases which I was enabled to examine, and that the majority of those, the notes of which I propose to read presently, were also examined by them, by Mr. Nettleship, or by the ophthalmic assistant at St. Thomas's Hospital, with the exception of Case VIII, which Mr. Mackinlay kindly saw. These seventeen cases are as follows: Nine recoveries, in which the base was supposed to have been fractured; three fatal cases, in two of which a fracture of the base was proved at the *post-mortem* examination, and in one of which a fracture of the middle fossa was diagnosed, but a *post-mortem* examination not obtained; four recoveries from concussion of the brain, in which optic neuritis was found; and one fatal case of bullet wound of the brain, in which there was a second attack of optic neuritis.

I will now read short notes of the cases in which optic neuritis was present, and in which recovery from the head injury as well as from the neuritis ensued. In the majority of them examination of the urine proved the absence of albumen, and no evidence of syphilis could be obtained.

CASE I.—A butcher, aged 32, was under the care of Mr. Clutton in St. Thomas's Hospital from August 19th to September 10th, 1887. He had fallen from a ladder 10 feet high four days before. He suffered from very severe headache, and there was a mark of injury over the back of the head, with slight wound. On the fifth

Fig. 5.—Temperature Chart in Case I. Optic Neuritis.

day after admission the edges of both optic discs were noted as hazy and obscured; veins full; tortuosity of some smaller vessels at the edges of the optic discs. On the nineteenth day Mr. Nettleship reported that both optic discs were hazy; veins rather full;

³ *Trans. Ophth. Soc.*, vol. ii, p. 208.

⁴ *Ibid.*, vol. v.

changes more marked in the left, where they were conclusive of papillitis; in the right, taken alone, they were of doubtful meaning.

CASE II.—A man, aged 24, was under the care of Sir Wm. Mac Cormac from April 8th to May 8th, 1887. He had been pitched on his head from a cart. There was profuse and prolonged bleeding from the ears, and he was unconscious for some hours. Next day he was restless, and fighting to get out of bed, and at night delirious, with a temperature of 104° ; he was very irritable at times. On the eighth day Mr. Robert Nairne found both discs slightly hazy, but without enlargement of vessels. On the eleventh day the left disc was more clearly defined, but examination was difficult, as the patient was very irritable, and resisted.

CASE III.—A greengrocer, aged 47, was under the care of Mr. Mackellar from September 29th to October 20th, 1887. He had been run over, and continued unconscious for some time. On recovery he was drowsy, and complained of headache. Mr. Wingfield-Roll, who examined the eyes, reported on the sixth day that the appearances were highly suggestive of optic neuritis; both

Fig. 6.—Temperature Chart in Case III. Optic Neuritis.

optic discs were rather hazy, with slight swelling, especially in the right; the veins were full; there were no hæmorrhages; and on the eleventh day the condition was more marked.

CASE IV.—A man, aged 37, under the care of Mr. Mackellar from July 29th to September 10th, 1888. He was admitted in a drowsy condition, with headache on the third day, after having been knocked down and injured his head. On the eighth day after the accident both optic discs were swollen, and their margins could not be made out; the vessels over the discs were very indistinct, but swollen over the adjoining part of the retina; and on the twenty-second day Mr. Lawford, who examined the patient, reported there was no swelling of the discs, but the vessels were all bordered by white lines, and the tissue over the surface was somewhat opaque. The appearances would quite support the diagnosis of an attack of optic neuritis some time previously.

CASE V.—A man, aged 47, under the care of Mr. Croft from July 18th to August 5th, 1888. He had fallen from a wall, and been unconscious for half an hour. There was a superficial scalp wound behind the right parietal eminence. He complained of some headache, and his mind was obscured. On the fifth day there was impairment of vision, and he could not see so clearly with the right as with the left eye. In the right eye the vessels were swollen and tortuous, and the disc very indistinct; the left disc was not quite clearly defined, but otherwise it appeared normal. On the ninth day there was still well-marked optic neuritis.

CASE VI.—A boy, aged 12, under the care of Mr. Mackellar from December 20th, 1887, to February 9th, 1888. He had fallen from a brick wall about five feet in height on concrete pavement; was supposed to have been unconscious for an hour; symptoms of cerebral irritation. On the fourteenth day he was still very drowsy with headache, and examination of the eyes by Mr. Lawford showed the presence of optic neuritis, moderate swelling of the discs, with much streaky haze, veins full and bending, but no hæmorrhages.

CASE VII.—A woman, aged 23, under the care of Mr. Mackellar from July 8th to the 25th, 1888, was admitted unconscious with hæmorrhage from the right ear and mouth, and subconjunctival ecchymosis of the right eye. Afterwards she was noisy and restless. On the ninth day the eyes examined with the ophthalmoscope revealed haziness of the discs, and the margins were ill-defined.

Fig. 7.—Temperature Chart in Case VII. Optic Neuritis.

CASE VIII.—A woman, aged 55, under my care at the Royal Free Hospital from December 24th, 1888, to February 13th, 1889. She had been knocked down in the street; she afterwards suffered from headache and drowsiness. The eyes were examined on more than one occasion by Mr. Mackinlay, who reported on the twenty-sixth day that there was undoubted optic neuritis in the right eye. The fundus of the left eye could not be examined on account of the presence of cataract. Nine days later the inflammation was diminishing.

CASE IX.—In one other case, that of a carpenter, aged 22, who was supposed to have received a fracture involving the anterior and middle fossæ of the base; slight haze of the disc was noticed on the fifth day, but Mr. R. Nairne, who also examined the eyes, did not think it sufficient to indicate the presence of optic neuritis.

OF THE THREE CASES IN WHICH THIS SYMPTOM WAS PRESENT IN FRACTURES OF THE BASE, WHICH TERMINATED FATALLY.

CASE X.—An omnibus driver, aged 62, under the care of Sir William Mac Cormac, who had fallen and received a scalp wound at the back of his head, presented symptoms of fracture in the posterior and middle fossæ of the base of the skull. On the sixth day there was commencing optic neuritis. He died on the tenth day with symptoms of delirium tremens. At the *post-mortem* examination there was great bruising of and hæmorrhage into the anterior parts of the brain. No meningitis.

CASE XI.—A leather-dresser, aged 40, under the care of Mr. Sydney Jones from February 2nd to the 12th, 1886, was admitted with very severe headache three days after a fall on the back of his head in the street. He became restless, and meningitis developed. On the tenth day after the accident Mr. Johnstone noted some haze about the optic discs, especially the right, but the man was very irritable, and they were not well seen. At the *post-mortem* examination there was well-marked basal meningitis, and a fissure in the posterior fossa of the skull extending forwards to the petrous bone.

CASE XII.—A boy, aged 7, under the care of Mr. W. Anderson from August 26th to September 5th, 1886. He had been knocked down and run over in the street, and presented signs of fracture in the middle fossa of the base. Meningitis with hemiplegia developed, and on the seventh day Mr. Lawford reported optic neuritis in both eyes. The patient died three days later, but an examination of the body could not be obtained.

CASES OF CONCUSSION, FOUR IN NUMBER, IN WHICH OPTIC NEURITIS DEVELOPED.

CASE XIII.—A man, aged 45, under the care of Mr. Croft from September 14th to October 12th, 1887. He was standing on the edge of a crate, which turned over and fell on him. He became unconscious, and was admitted with a contusion on the right side of the head and swelling of the right shoulder, stertor, with blowing respiration, a small rapid pulse of 72, and the pupils equally dilated and not acting to light. Two days later he was more conscious, and the respiration was quieter. On the seventeenth day, on ophthalmoscopic examination, the left optic disc was slightly swollen and hazy at the margin, with bending of the vessels; one vessel was obscured by new tissue. The right optic disc presented appearances of a similar character, but they were less marked. The highest temperature was 99.6° .

CASE XIV.—A tramcar conductor, aged 26, was under the care of Mr. Mackellar from July 8th to the 29th, 1888. He had fallen off the car steps on his head. He never became unconscious, but

was sick soon afterwards. When admitted, after a long walk, he appeared dazed. There was a hæmatoma over the vertex and right temple; pupils normal. He vomited from time to time during the remainder of the day and suffered from headache. On the following day there was a slight rise of temperature to 100.2° ; the sickness had passed off, but there was still headache. On the seventh day there was much pain in the right temporal region, and the right pupil was the larger. On the tenth day both optic discs were found to be much swollen and hazy, and the retinal vessels were enlarged. In this case the temperature was generally over 99° and under 100° in the evening until the ninth day.

CASE XV.—A carman, aged 40, under the care of Sir W. MacCormac from October 25th to December 15th, 1886. He had fallen from a cart and received a superficial scalp wound in the occipital region. Was suffering from shock on admission, then passed into a deep sleep, which gave rise to a drowsy condition, from which he was roused with difficulty. The temperature was frequently elevated, 100° to 102° , occasionally rising to 102.6° ; and on the sixth day he had a severe convulsive seizure. Afterwards there were drowsiness and frontal headache. The fits continued at intervals until the tenth day, when they ceased. After this there was delirium, with sleeplessness and occasional fits. On the fourteenth day he was sensible again, complaining of headache, but was still violent at night. The evacuations were passed involuntarily. Was quite sensible on the twenty-fourth day, and then improvement continued. Mr. Nairne reported haziness of the discs, with dilatation and tortuosity of the vessels, on the sixth day. Mr. Lawford examined the eyes eighteen days later, and found slight but definite papillitis present.

CASE XVI.—A gentleman, aged 30, who came under my care a week after falling on his head. There had been loss of consciousness for an hour or two, with vomiting. Since that time drowsiness and severe headache with intolerance of life, but no blindness. Examination of the discs showed well-marked swelling and haze about the tenth day. He was kept quiet. Under treatment the symptoms all rapidly subsided.

The following is a most interesting example of bullet wound of the brain:—

CASE XVII.—A boy, aged 11, was under the care of Mr. Mackellar from April 13th to October 12th, 1887. There was a wound caused by a revolver bullet of small size, two inches above and an inch and a half in front of the left external auditory meatus. Two days after admission optic neuritis developed. Mr. R. Nairne reported: "Both discs slightly hazy and congested. The large vessels, both arteries, and veins dilated; all the small vessels running over the edge of the disc appear more evident than normal." On the third day he was trephined. On the sixth day Mr. Nairne noted more haze, with definite swelling of the discs on each side, especially at the lower edge. On the eighth day the swelling was more marked in the right eye; there were no hæmorrhages. On the eighteenth day, still marked neuritis in both eyes; swelling not great, but more in the right than in the left. Much shot-silk appearance of retina and a few bright spots in the right yellow spot region. On the thirty-seventh day Mr. Wingfield-Roll reported that there was still slight haze, with moderate swelling; veins on both sides large and tortuous; and on August 8th, nearly four months after the injury, some streakiness about the optic discs, with a little haze of the retina surrounding them. One or two dots of black pigment near the optic discs, possibly the remains of hæmorrhages. More streakiness and haze in the right. Hernia cerebri developed after the operation to a considerable extent, but gradually subsided, and the patient went to a convalescent home. He was again in the hospital, under Mr. Mackellar's care, from October 29th to December 4th of the same year with traumatic epilepsy. Ophthalmoscopic examination showed the veins very tortuous near the discs, which were not pale. They had a good colour, but some streakiness at the edges; there was nothing further to indicate that there had been optic neuritis on any previous occasion. When he left he was very dull, and avoided the sunlight; had no fits or headache, but staggered occasionally in walking. He was again admitted about eight months after receipt of the injury, and died early in 1888, forty-one days later. He had suffered from from fits, pain in the left side of the head, loss of power in the right arm, right facial paralysis, vomiting, and bulging of the scar. After admission Mr. Wingfield-Roll found moderate papillitis, with tortuosity and abrupt bending of the veins at the edges of the disc, obscured by streaky haze; changes less in the left eye. And two days later Mr. Nettleship reported "well-marked papillitis of each, with moderate steep swelling, without hæmorrhages."

On the twenty-third day, aspiration through the scar drew off a drachm of clear fluid; on the twenty-seventh, Mr. Mackellar explored the brain, removing more bone. Hernia cerebri again developed, and an abscess cavity was incised. At the *post-mortem* examination there was a thick-walled abscess cavity in the posterior part of the left frontal convolution, extending into the white matter of the parietal and temporo-sphenoidal lobes and basal meningitis. The bullet was found in the white matter of the right frontal lobe low down, surrounded by apparently healthy white matter. The track made by the bullet to reach that situation could not be traced. I have given this case more in detail than the others, as it is an example of neuritis affecting the optic nerves for a second time, when the first attack had been severe, but completely recovered from. It will be noticed that, although the inflammation was very marked in many of these cases, no hæmorrhages were at any time found by those who examined them, and in only one was there any complaint of blindness. Taking, then, the four cases of recovery from optic neuritis (that is, excluding the two cases in which one eye only was involved) and the two other cases which proved fatal, and in which evidence of inflammation of the nerves was present during life, we have a total of six from the series by Messrs. Edmunds and Lawford which we can compare with those which I have just given. I would not, however, have it thought that I am detracting from the value of their observations on the subject. This is not my intention. I am simply desirous of investigating the usefulness of this sign as an evidence of fracture of the base of the skull or injury to the base of the brain, and put on one side those cases which do not appear to assist in the elucidation of the question as to the importance of optic neuritis as a clinical sign.

I should like, however, before proceeding further, to refer to the results of the microscopical examinations made by Messrs. Edmunds and Lawford in four cases which proved fatal from the severity of the injury, and in which the base was involved. In two of them which were examined the patients had only lived twenty-four hours after the injury. The result is thus summarised: "In those four cases in which there was neuritis the changes observed with the microscope were, in one case, present throughout the whole length of the nerves, but confined to their surface (perineuritis), and in the remaining three cases the changes did not extend throughout the whole length of the nerve, but were only found at its proximal part, and in one of these three the hyperneuculation was more marked at the surface of the nerve than in its interior."⁵ The microscopical changes are well illustrated in their paper, and through the courtesy of Mr. Edmunds I have been enabled to examine the microscopical sections of the nerves from which the illustrations are taken. The change from the normal which is found consists in a considerable increase in the number of staining corpuscles seen in the sections, especially in the trabeculae between the nerve bundles. Sections from one case show also the existence of fluid between the outer and inner nerve sheath, and others from another case the presence of inflammation almost entirely limited to this space, with but slight changes in the nerves themselves. It is evident, then, that the inflammatory process may commence very soon after the injury, and microscopical evidence may be afforded within a comparatively short time. What is the period at which we are most likely to meet with it at the bedside? This is a question that I am not able to answer definitely; I can only mention the times at which it was found. Often the patients were too irritable to permit of examination during the first day or so after admission, and others were not examined until some special symptom, such as headache, made it appear possible that there might be a neuritis. Of the cases in which the base was fractured, in three it was found on the fifth day, in three on the sixth, in one on the seventh, in two on the eighth, in one on the ninth, and in one on the tenth day, and not until the fourteenth in one and the twenty-sixth in another, though repeated examinations were made. In the cases where concussion was present without evidence of injury to the bones it was found on the sixth day in one, the tenth in two, and the seventeenth day in one; whilst in the case of bullet wound of brain it was found when the boy was examined two days after the injury. It is not possible to give the approximate date of onset of the second attack.

Messrs. Edmunds and Lawford conclude that "optic neuritis occurs with greater frequency in those cases in which the base

⁵ *Ibid.*, vol. vii, 1887, p. 208. See also *St. Thomas's Hospital Reports*, vol. xi, p. 71.

of the brain or skull is affected than in those in which it is not.⁶ The cases which I have given support this conclusion. I would even go further, and add that it is much more likely to follow injuries inflicted on the posterior part of the skull than on the lateral or anterior aspects. Thus it was found in two patients in whom fracture of the middle fossa was diagnosed, in one with probable fracture in the middle and anterior fossæ, in three with fracture of the middle and posterior fossæ, and in six in which fracture of the posterior fossa was alone present—that is, there was a fracture involving the posterior fossa in nine out of twelve fractures of the base in which optic neuritis was found—a very large proportion.

Dr. C. S. Jeaffreson, in a recent lecture on Optic Neuritis,⁷ says that injuries to the head are not a very common cause of optic neuritis unless they are severe and attended with fracture in the neighbourhood of the optic foramen. I agree with the earlier part of his statement, but have not met with any cases that appear to confirm the latter as to the frequent existence of fracture in the vicinity of the optic foramina. It may be, however, that his remark is intended to apply more to the affection of one nerve, and is founded on the statement of Van Hölder as to the frequency of fracture in that situation.

That optic neuritis may be the result of extension of inflammation of traumatic origin from the membranes of the brain is shown by examples only too numerous. Here the meningitis has produced symptoms leading to diagnosis of the disease before the ophthalmoscope has given evidence of the inflammation of the nerves; it is simply confirmatory of the opinion to which the surgeon has already come. Messrs. Edmunds and Lawford are inclined to think the neuritis to be due to a meningitis of the base even in the cases that recover, and this seems the most likely explanation; if it is so, however, we must suppose a very localised and subacute inflammation not sufficient to produce any effect on the temperature curve, for, as shown by the charts of the temperature in cases apparently uncomplicated by wound of the soft parts, severe cerebral contusion or fracture communicating with the external air, the temperature may be below normal when the optic neuritis is at its height, judging from ophthalmoscopic examination. It is not improbable that the inflammatory process is started most usually by a contusion of the brain substance in the neighbourhood, and then passes along the cerebral membranes to the optic nerves. We know how frequently contusion of the under part of the frontal lobes is found after death when the fracture is situated in the posterior fossa of the skull; and I have shown that it is in these cases that we most frequently meet with optic neuritis after a fracture of the base.⁸

INJURIES OF THE CRANIAL NERVES.

Loss of function of one or more of the cranial nerves is not unusual after an injury to the head of some severity, but cannot be described as a symptom of fracture in every case, even when the evidence of the injury to the nerve is at once manifest. The olfactory nerves have been torn across in a fracture of the base of the skull, and it is stated that paralysis of these nerves not infrequently follows severe injuries to the head without evidence of fracture. Loss of smell was complained of by only one of the patients who were under my observation. Examples of injury to these nerves were afforded by the *post-mortem* examination of two cases; in one there was contusion of the frontal lobes of the brain, with both fracture of the posterior and anterior fossa, and the olfactory nerves were much involved; in the other there was a similar injury to the brain in the frontal region with a fracture in the posterior fossa, and the left olfactory bulb was crushed like the brain substance. These cases are confirmatory of the opinion expressed by Sir Prescott Hewett⁹ that loss of smell is most likely to follow injuries in which the anterior lobes of the brain are driven against the bone and bruised. The optic nerves may be injured in a thrust wound of the orbit, the instrument which produces the fracture of the bone dividing or lacerating the nerve. They may also be divided by a bullet, sometimes without injury to the brain, in its progress across the head from temple to temple. Such cases are recorded in the *Surgical History of the American War of the Rebellion*, and are rare in civil practice. I saw one such, however, under the care of Mr. Sydney Jones¹⁰ last year, in which the injury had been self-inflicted with a revolver.

A curious instance of damage to one optic nerve from a bullet wound was under my close observation in 1876. A boy, aged 15, was admitted under the care of Sir W. Mac Cormac with a wound of the upper and inner angle of the right orbit, caused by a bullet from a revolver a short time previously. This injury was immediately followed by loss of sight in the eye of the opposite side. A probe passed along the track of the bullet went backwards and to the left for three inches and a half. He was suffering from shock but no symptoms developed. I saw him some years later, and the eye remained disabled, with an atrophied disc. Pressure upon the nerves by a fragment of the broken sphenoid bone has produced complete blindness.

Dr. Berlin¹¹ read a paper at the International Medical Congress of 1881¹² on Injuries to the Optic Nerves and Ophthalmic Artery from Fracture of the Optic Canal, and detailed three cases, one of gunshot injury, with a fissured fracture of the roof of the canal, a second of fissured fracture through both optic foramina, and a third of laceration of the internal carotid, with fracture of both optic canals, in each of which there was a laceration of the nerves. The injuries which caused these fractures were necessarily severe and fatal. Van Hölder considered fracture of the optic canal as very frequent in fractures of the base, and Dr. Berlin thought that sudden single amaurosis, following immediately after injuries to the head was usually due to this cause.

Mr. Nettleship¹³ in a clinical lecture gives an account of some cases of injury to the optic nerve which he had observed. Three of them were under treatment in St. Thomas's in 1881, and I have notes of two cases¹⁴ under care since that time in the surgical wards, in both of which there was evidence of severe injury to the head, with proptosis, followed by atrophy of the disc and blindness. In one, atrophy was visible on the seventh day, and Mr. Nettleship noted on the twentieth day that there was still interference with the action of the muscles, and a thickening of the orbital roof. A coal porter,¹⁵ aged thirty, was admitted under the care of Mr. Sydney Jones on March 27th, 1884, having fallen twenty feet half an hour previously. He had been unconscious for a short time. He was a stout, heavy man, suffering from shock, cold and shivering, but conscious. Pulse 112. There was a wound of the upper lip, a wound of the nose, at the bottom of which could be felt a fissure of the bone. Much ecchymosis of the cellular tissue of the eyelids of both eyes was found, and on the right side it was already present under the ocular conjunctiva. There was slight hæmorrhage from the nose, mouth, and left ear. The bones of the face were fractured and separated, and fracture of the radius on both sides. The patient was noticed to be blind of the left eye on the fifth day, the left pupil was larger than the right, and did not respond well to the action of light. As the swelling in the eyelids subsided, it became possible to examine the eyes more fully, and on the eighteenth day a note was made to the effect that the external rectus of the left eye was almost completely paralysed, the pupil did not act to light, and the disc was pale with slight haze, and it was thought that there had been direct injury to the optic nerve, possibly by fracture through the optic foramen. The appearance of the left optic disc on the twentieth day resembled that found after embolism of the central artery of the retina. The various fractures gradually consolidated. The hæmorrhage from the ear was slight and the discharge which followed it of small quantity. There was no vomiting. The temperature for the first week was higher than normal, reaching from 100.8° to 102.° in the evening, and being 99° or 100° in the morning; after this it was normal. The strabismus was very marked when he left the hospital, after a residence of forty-five days. The left optic disc was extremely pale, and the arteries considerably diminished in size. There were some white lines along the vessels. The right eye continued apparently normal. The man never recovered from the effects of these grave injuries, and was almost constantly under medical care for various ailments, dying at last on February 4th, 1887, from general tuberculosis. There had been no improvement in the condition of the left eye; internal strabismus and complete atrophy of the disc remained. At the *post-mortem* examination made by Dr. Hadden there was "a united fracture, evidently quite old, of the middle fossa of the skull, which extended from near the apex of the petrous bone obliquely forwards and outwards to the anterior

⁶ *Trans. Ophth. Soc.*, vol. vii, p. 212.

⁷ *Lancet*, vol. i, 1890, p. 892.

⁸ *Op. cit.*

⁹ *Loc. cit.*, vol. iii, p. 614.

¹⁰ *Lancet*, vol. i, 1889, p. 424.

¹¹ *Med.-Chir. Trans.*, vol. xiv, p. 348.

¹² *Transactions*, vol. iii, p. 115.

¹³ *St. Thomas's Hospital Reports*, vol. ii, p. 116.

¹⁴ See series by Messrs. Edmunds and Lawford.

¹⁵ *Vide the Lancet*, vol. i, p. 105, 1886; *ib.*, vol. i, p. 877, 1887.

margin of the fossa. The margins of the longitudinal fissure of the brain were pretty firmly adherent in the anterior half, and the tips of the frontal lobes were adherent to the dura mater. There were evident signs of old meningitis at the base. The pia arachnoid was thickened and opaque, especially over the optic nerves, chiasma and tracts, the crura and pons. The optic nerves were dull grey, and the chiasma and tracts distinctly changed in colour. There was no marked diminution in size. The left third nerve was greyer than the right, and more flattened. The corpora quadrigemina were normal in size and appearance." Scattered tubercles were found in the brain. The fracture does not appear to have involved the optic foramen.

A man aged about 30 was brought to me at St. Thomas's Hospital in 1888, complaining of loss of sight in the right eye. A few months previously he had been an inmate of the hospital suffering from concussion and an injury to the right frontal region (? fracture), but had not complained of his eyesight. Mr. Lawford reported atrophy of the disc, and the eye was quite blind.

Mr. Nettleship¹⁶ says on the subject: "But in the more common cases the accident is an injury to the head, probably causing fracture of the roof of the optic canal, with crushing of and hæmorrhage into the nerve, or perhaps occasionally tearing of the nerve on the brain side of the canal. Both conditions have been shown to occur in fatal cases of fracture of the base of the skull."

Another possible cause for blindness after these injuries is the occurrence of hæmorrhage into the sheaths of the optic nerves. In none of the *post-mortem* records of cases in the series here brought forward was fracture in the neighbourhood of the optic foramen found, but in eight, where the damage to the base was severe, blood was found extravasated in the sheaths of the nerves. And in one at the end of eight days there was also slight neuritis. It is a remarkable fact, and one of considerable importance, that in these cases of injury to the head which I have related, and in which optic neuritis was found on ophthalmoscopic examination, if the neuritis was found in both eyes complete recovery ensued, but if after an injury to the frontal region a condition of neuritis developed in one eye only, the disc presented at a later date complete atrophy, and the patient remained permanently blind on that side. Paralysis of one or more branches of the third nerve may ensue as a consequence of injury to the head, but it was very rarely seen, the difficulty in movement presented by various ocular muscles appearing to be the result of hæmorrhage, and the interference which it caused to their action, rather than to any injury to the nerves supplying them. When paralysis of the third nerve was present it was due to compression by an extravasation of blood at the base. In one fatal case in which it was not possible to get a *post-mortem* examination paralysis of the sixth nerve was noticed on admission, soon after the accident. In five patients who recovered this paralysis was present when they were first examined; of these the injury had been very severe in all, and the squint continued without sign of improvement when they passed from observation. In one of these it was associated with optic atrophy, and in one with simultaneous loss of power in the facial nerve of the same side. In all a fracture of the middle fossa was diagnosed. Of the four others in whom it appeared at a later date—on the second, third, fifth, and thirteenth day—the manifestations of injury were severe, and the squint continued without improvement when last seen, with the exception of the patient in whom it appeared on the third day, and was associated with proptosis, and later with optic atrophy; in his case there was gradual improvement in the squint. It is probable that when the nerve is paralysed from the first the paralysis is due to rupture of the nerve, though a case recorded¹⁷ does not appear to support this. In that case it is stated that the squint did not develop until three weeks after the injury, yet at the *post-mortem* examination the sixth nerve was found snapped across. One can hardly think there was no paralysis before three weeks after the injury. Although in these examples the state of the nerves did not, with one exception, improve under observation, it is the experience of others that in several instances power has been gradually regained.

In a child, aged $3\frac{1}{2}$ years, suffering from concussion, temporary paralysis of this nerve was observed for a period of six days, but with reference to squint in children, the statements of the mothers as to its absence before the head injury do not always agree with what is said by other relatives. There is another thing to be borne in mind, and that is the possibility of the squint being a

manifestation of syphilis. An old patient of mine, who had suffered severely from tertiary manifestations some years ago, recently fell forwards accidentally and struck his forehead against the mantelshelf in his room. Three days afterwards double vision gradually developed, which made it difficult for him to move about anywhere. Under iodide of potassium he recovered at the end of two months. That such a symptom as strabismus is important from the interference with vision that may follow it is well known, and as the prognosis depends much on the date of its appearance, it should be decided by the medical man when he first sees the patient whether strabismus is present or not, for the question of damages in a law court may depend much on this point, as in a case recently related to me.

Paralysis of the facial nerve is an accompaniment of many fractures of the base; indeed, like hæmorrhage from the ear, it may be regarded as a most valuable sign of fracture in the middle fossa. Mr. Pick,¹⁸ in his remarks on some cases of fracture of the base, states that in thirteen well marked instances this symptom was present in ten, and that it developed at once in two and in a period of from two to six days in eight. It was not met with in anything like this proportion in the cases under my observation; in only fifteen was paralysis of the facial nerve sufficiently marked to attract attention, if we exclude two in which the patients were the subjects of old hemiplegia. In six the paralysis was present on admission, and persisted when the patients (four in number) who recovered were able to leave the hospital. In the two fatal cases it was found that the fracture which extended through the internal auditory meatus had produced considerable laceration of the nerves. In the others it appeared as follows: On the second day in a man aged 31, after severe hæmorrhage from the ear, in whom the fracture was proved to have passed across the internal auditory meatus without lacerating the nerves. On the fourth day in a man aged 48, who recovered. On the fifth day in a man aged 29, who died from erysipelas and pyæmia. On the eighth day in a man aged 62, who died from delirium tremens. The fracture involved the internal auditory meatus, but no coarse lesion of the nerves was seen. On the tenth day in a boy aged 10. There was much œdema of the scalp and suppuration in a scalp wound at the time. The paralysis had nearly gone in a month. On the tenth day in a man aged 23, preceded by deafness the day before. The paralysis had nearly disappeared in a fortnight. On the sixteenth day in a man aged 37, with an attack of erysipelas. The hæmorrhage from the ear had been very free at first. The paralysis was transient. On the eighteenth day in a man aged 40. Deafness before the accident increased during the period of hæmorrhage, four days. The paralysis lasted six days, and during the time deafness was more manifest.

OBSERVATIONS

ON

THE TREATMENT OF PERNICIOUS ANÆMIA

BASED ON A STUDY OF ITS CAUSATION

WITH NOTES OF A CASE.

By WILLIAM HUNTER, M.D., M.R.C.P.,

John Lucas Walker Research Student, University of Cambridge; Arris and Gale Lecturer, Royal College of Surgeons.

(Continued from page 1.)

CAUSATION.—EXCRETION OF AROMATIC SULPHATES.

THE further series of observations I made on the urine in the above case related to the cause of the extensive blood destruction prevailing in this condition. I concluded as the result of my first investigation, for reasons I have already indicated, that to produce a condition of blood and of the liver such as was found in pernicious anæmia, the action of a special poison was necessary. With regard to the nature of the poison or poisons thus generated, I stated that my observations up to that time had not supplied me with any definite information. From analogy, however, afforded by the action of the toluyene-diamine, more especially from the limitation of its action to the portal blood, I concluded that "the poison was probably of a cadaveric nature produced within the gastro-intestinal tract—in excessively small

¹⁶ *Loc. sup. cit.*

¹⁷ *Journal de l'Expérience*, 1843.

¹⁸ *JOURNAL*, 1865, p. 530.