clots. In this case an abscess forms, and the poison of pyæmia is thus circulated. The necropsy showed that the left profunda femoris artery had been the seat of a small aneurysm about the size of a hazel nut, and that this had burst at last into the muscles of the thigh. In this way a column of blood was driven with peculiar force with each pulsation and injected into the muscles, compressing suddenly the anterior crural and the great sciatic nerves. This fully accounted for the agonising pain experienced. In this case there was no reason to believe that the valvular disease was of infective character; or, in other words, that it was an instance of so-called ulcerative or malignant endocarditis. The clinical, no less than the *post-mortem*, evidence was against this view of the case. In a mechanical sense, the profunda femoris artery presents a favourable position for an obstructing embolus, and I think it probable that a secondary thrombus may have extended from this vessel along the external circumflex artery.

And now in respect of treatment of this and such like cases. I regret to say there is little to be done. The indication is to secure rest and to maintain good nutrition of the cardiac walls. Without an exact diagnosis we acted rightly in this case. And here I would observe, by way of parenthesis, that you will often have to treat cases in which you are unable to make a complete or accurate diagnosis. You are not to withhold measures of relief from any patient because you do not know exactly with what you have to deal. Remember that our business is always to do our best according to the light we have, and hence we have often to prescribe for symptoms whose full significance we have imperfect knowledge of. We must feel our way with caution and hope in time to gain more insight, and we can often do good.

The modern therapist is too much disposed to hold his hand and, in deference to what he calls scientific practice, to wait, do nothing, and let Nature take charge. But unfortunately Nature is not always kind, and in her workings causes much suffering and misery, which, as practitioners of medical art, we can surely sometimes relieve. I would, therefore, urge a little less science and a good deal more art in many points of practical medicine at the present time. Our forefathers had more of the art than we have, and we have lost it in proportion as we have striven to be always scientific. Never be ashamed to say you do not know. We know a good deal and we shall know more, but we must help our patients as far as we can, and often while in ignorance of much that is going on.

In this case, then, in respect of treatment I find no room for regret. I do not see that more could have been done to afford relief or to ward off the rupture of the small aneurysm. I pass round the specimen that you may study it for yourselves, and Mr. Mark has made a drawing which is reproduced on the preceding page. I could wish to tell you something more encouraging in respect of the treatment of such cases as these, but I know not where to find it. Happily they are not common, and it is, at any rate, encouraging to know that many cases of aortic disease follow a happier course than this untoward one. Why in one case there should be this proliferative (vegetative) form of endocarditis, and in others no such overgrowth should occur we do not know, but it is probably a matter of this is proclivity and personal peculiarity.

To trace the processes back somewhat remotely, we may say that, with care to avoid undue strain and exposure, rheumatic disease may be averted or held in check, and that with strict temperance in all things, many of the worst effects of cardiac and vascular disease may also be kept in abeyance. In some of these cases benefit may be had from the employment of iodide of potassium and iron, and the undue action of the heart may be moderated, provided always the patient is sufficiently fed and nourished, by aconite given carefully and its effects frequently noted.

Five or ten minims of the tincture may be added to any other medicine which is employed, and in aneurysms which are accompanied by a frequent pulse I have found on several occasions much relief from this drug. It cannot, however, be employed while treatment after Tufnell's method is being carried out, since the patient is insufficiently nourished to bear it with safety. A case which well illustrates this you may see in No. 5 bed in John Ward at the present time.

PSILOISIS (LINGUÆ ET MUCOSÆ INTESTINI).

(With Chromolithograph.)

By GEORGE THIN, M.D.

In temperate and cold climates diseases of the respiratory organs are common, and diseases of the intestinal canal in adults in comparison rare. In tropical and subtropical climates, on the contrary, the relation is reversed. I once took occasion to express this fact by remarking that in the tropics disease strikes below the belt. But although cough is a symptom common to various affections of the respiratory organs, we do not use such terms as "the pulmonary flux of cold climates," or "arctic cough," whilst in regard to the intestinal derangements of tropical countries we do not seem to have got beyond the primitive stage designated by such expressions as "tropical diarrhoea," or "the intestinal flux" of hot countries.

The intestinal canal of an adult man is about eight mètres long. Nearly every disturbance of the structure or function of this long tube is attended by fluid discharge from some part of the mucous membrane, that is, by diarrhoea; but it does not require much consideration to understand that the flux may be produced by diverse causes acting in different parts of the canal. In the *Practitioner* for September, 1883, I published a short and very incomplete paper on an intestinal affection known in the East as "Indian sprue," the term given to it by the Dutch physicians of Batavia, where the disease is common, and who appear to have been the first to consider that there is a specific affection with symptoms which separate it from other forms of intestinal disease attended with diarrhoea. In the *Practitioner* for 1887 I published a series of papers giving a fuller account of the literature of the disease so far as it was known to me, and more particularly reporting in detail six cases of the affection which had been treated by myself.¹

Briefly summarised, psilosis may be described as a chronic affection of the intestinal tract unaccompanied by fever (or any abnormality in the urine), with a remarkable and unusual disposition to an affection of the tongue and buccal mucous membrane, and with tenderness in the gullet. There is no blood or mucus in the stools, and no straining. As a rule the diarrhoea is not acute, or in any sense of the word severe, although it may be so paroxysmally. The motions may be considered characteristic. They are either of a pale straw, yellowish colour, frothy, quite unfurrowed (in the first stages being chiefly passed in the early morning), or (and this is especially so in chronic cases which have lasted some time) they consist of a greyish-white pultaceous mass, often in a state of apparent fermentation, the motion being found covered with froth.

In some cases the symptoms in the tongue and mouth are particularly troublesome, the patient being unable to swallow anything that is hot either by temperature or seasoning, and even the weakest alcoholic drinks are not tolerated. A remarkable peculiarity of the affections of the tongue and mouth is their fugitive character. The tongue or mucous membrane may become the seat of an eruption which is almost herpetic at the outset, soon leaving the surface bare; or the whole tongue, or patches of it, may simply become gradually red and bare. In early cases this appearance may entirely disappear in a few days, but eruptions on the tongue and mouth frequently precede the relapses which are so common. The disease begins insidiously, is usually very chronic, and even after it is apparently cured the patient is very liable to a relapse. While it continues the patient becomes more and more emaciated, and often extremely anæmic. There is no jaundice and no bile in the urine, which is usually light coloured. The liver has a tendency to become smaller, and often becomes very much shrunken.

It has been long recognised from the clinical symptoms that certainly in the mouth, and probably in other parts of the canal, there was bareness or rawness of the mucous membrane, and it was to express this condition that I ventured to propose the term "psilosis" in order to invite the attention of observers to one of the

¹ These papers were republished by Messrs. Churchill in a small pamphlet, entitled, *Psilosis or Sprue; its Nature and Treatment*, and I wish to refer such of my readers as are interested in the subject to the *Practitioner* or to this pamphlet. I am obliged to lay particular stress on this reference, as the space now at my disposal does not permit me to discuss at present either the symptoms or the treatment.

SUSPENSION.—Tiberghien (*Ann de Méd. et de Chir.*, Bruxelles, 1390, p. 21) reports excellent results from suspension in twenty-six cases of locomotor ataxy with various complications.

most characteristic symptoms of the malady. I have had recently further opportunities of studying this disease, and, incomplete as these studies are, I believe it will be advantageous to publish some of them in order to suggest criticism and stimulate observation. Through the medium of the JOURNAL I hope that they will reach observers who are now treating cases both in England and abroad.

Psilosis, although not a very common disease, certainly cannot be considered rare. It constitutes a danger which specially threatens middle-aged and elderly residents in the Straits, in Batavia, in certain parts of China, in Ceylon, and probably some other parts of India. In patients who do not do well it is an exceedingly wearing and distressing malady, and in elderly or exhausted constitutions, the chances of rapid recovery are by no means so great as they are in younger people. The marvellous manner in which many of the patients recover when put on milk diet is one of the most striking things connected with the disease. The great majority of young patients, and many even of the elderly ones, respond at once to a diet of this kind properly carried out. Unfortunately the patients frequently do not consent to the necessary restrictions until the disease is comparatively far advanced: but even then in many of the younger patients, and in not a few of the older ones, a satisfactory cure may be obtained. A good result, however, cannot be reckoned on with certainty.

In the present paper it is my object to direct attention to two conditions which are probably peculiar to psilosis. One of these—the state of the tongue—I have thought it well to illustrate in the coloured drawing which accompanies this paper. The other point to which I now direct attention is the pathology of the disease. In 1888 I felt compelled to state that I was able to say almost nothing about its pathology and absolutely nothing of its pathological histology.

The patient, to whose kindness I am indebted for permission to have the drawing made which is reproduced in the accompanying plate, is a middle-aged man, who had spent the best part of twenty years in the south of China. After the first ten years' residence he visited England, and whilst at home suffered from unmistakable symptoms of psilosis, from which, however, he quite recovered, and then returned to China. After his return he was from time to time the subject of intestinal irritation, which disappeared on milk diet. He finally returned to England three years ago. When the cold weather began, the first winter after his return, the symptoms from which he suffered ten years previously returned; his digestion and general condition became bad, and he passed about three somewhat loose motions daily. When I first saw him, six months afterwards, he had lost two stones in weight, was free from fever, had an exhausted, worn appearance, and had not passed healthy motions during the whole winter. He was passing two to three motions daily, sometimes large in quantity, described as being yellowish in colour and of the consistence of jam. I purposely omit at present important details in connection with the case in order to devote the available space to the description of the tongue. The third time I saw him I described it as follows:—

"The whole surface of the tongue is smoother than natural, not furred, with a few cracks; but, taken as a whole, it is neither red nor glazed. On the surface of the right side, near the middle, there is a deep sinuous cleft extending longitudinally for over an inch. This cleft is very red, and has the appearance as if it bled slightly. Under the tongue in front there are several projecting pustular appearances and several rose-coloured or raw-looking patches. These make the tongue so tender that he cannot protrude it. On the inner surface of the under lip there are several small red patches and two pustules. The mucous membrane of the cheeks and roof of the mouth is very pale and anæmic. On the right side, in the centre of the cheek and extending backwards, there are several patches of a dark rose colour, the borders of which are not well defined."

The patient steadily improved under appropriate diet, and four weeks after the above description was written I noted regarding the tongue that "the fissure which bled on the surface is hardly to be detected. The surface of the tongue is pale; certain parts have a pinkish colour and are without fur, other parts having a deep red and spongy appearance. The point is a deep pink and raw looking. On the right side, extending from where the fissure was to the edge of the tongue and dipping under the edge, is a deep red surface. The pustules that were under the tongue have disappeared, and he can now protrude it." The claret-coloured and irregular patches on the cheeks were smaller, and there were

fewer of them than on the previous occasion. After this time the patient improved rapidly, and the tongue got so well that, although I had made arrangements to have it drawn, the delay of one week was sufficient to render the appearance too indistinct to be valuable.

The general condition improved even more rapidly than that of the tongue. Within eight weeks of his being put on milk diet he had gained a stone in weight and passed healthy, formed motions. He passed successfully through the various gradations of diet described in the pamphlet to which I have referred, and became comparatively well. Unfortunately he had a relapse during the following winter, and I saw him in the end of December in a condition very similar to that for which he first consulted me—the condition being that of typical psilosis. The state of the tongue continued to vary for some time, improvement not supervening so rapidly on milk diet as it did on the former occasion. During a period of exacerbation in the condition of the tongue a drawing was made by Mr. Burgess. About this time my notes described the tongue as follows:—

"The upper surface is of a pale, pinkish-rose colour, the reddest parts running indefinitely into the whiter parts; not glazed. On the under side of the tongue on each side of the raphe there are circular, macerated-looking spots, deeply red, with small white patches. On the left side of the tongue, on the under surface, there is an irregular patch, nearly as large as a sixpence, distinctly limited on its borders with little islands of white epithelium in it. The borders are slightly whitish, the other parts are of a deeper red than usual. There is a similar patch under the right surface of the tongue. On the soft palate there are several erythematous patches." The patient was still free from fever, passing, on an average, two pulpy, whitish grey, offensive motions daily, sometimes partially formed, and the condition generally was that recognised as characteristic of the disease in its full development. There has been in both attacks a history of the usual sensitiveness to acrid or hot substances, with more or less tenderness of the tongue.

It is characteristic of the state of the tongue in this disease that it varies rapidly. Within a week of the time that the drawing was made, the upper surface was much less cracked and altogether smoother, and the white and red patches on the under surface of the tongue had almost entirely disappeared, both the upper and under surfaces presenting simply a uniform rosy tint, deeper than that of the healthy tongue, with a smoother surface and with the cracks fewer and shallower than is shown in the drawing. This rapid change in the condition of the tongue is remarkable, and, so far as I am aware, is peculiar to this affection. In patients who do well under treatment, of whom fortunately there are many, well-defined appearances as characteristic as those shown in this drawing often disappear within ten to twelve days. It is of importance that attention should be directed to this point, as one case has come under my observation in which the disease was mistaken for syphilis, and the patient subjected to antisyphilitic treatment.

A case that unfortunately ended fatally under my care in the beginning of this year enables me to throw some light on the symptoms of this disease, particularly those that characterise the advanced stages. I saw the patient, a man about 57 years of age, in this country some five or six years ago. Although previous to that time he had suffered from severe attacks of renal calculus and, apparently, gallstones, he looked healthy and well, and much younger than his age. When he came this time under my care, he was in a hopelessly exhausted condition, and I am indebted to Dr. Henderson, of Shanghai, and Mr. Low, who took charge of him on the passage to England, for valuable notes connected with the case. Briefly stated, the main facts are as follows:—

In the winter of 1887 he was very severely chilled by exposure amongst ice and subsequent travelling in the extreme north of China. After that time, he ceased to notice perspiration, and had occasional attacks of diarrhoea. In March, 1889, still in the extreme north of China, he had a severe feverish illness, after which diarrhoea was very marked. At this time he lost 65 lbs. in weight in three months. Change of air was tried without benefit. Dr. Henderson, who happened to be at Chefoo in August, 1889, saw him there, and, from the notes with which he has kindly furnished me, I find that the diarrhoea continued severe, the motions being liquid, pale yellow in colour, scanty, sometimes very frothy and offensive, but containing no blood. The patient was thin, anæmic, and sallow. The tongue was pale, smooth, and polished, with lines on the surface. No aphthae.

Dr. Henderson again saw him in October, at Shanghai, when the motions are described as being either liquid and yellow, or thicker and offensive, sometimes pale, sometimes watery. The pulse averaged 86 to 88, and the temperature always remained normal, or below normal. Mr. Low, who took charge of the patient from Shanghai to England, states that there were at first from two to five motions daily, the discharge being loose, yellow, sometimes pasty, sometimes frothy and bad smelling; the tongue being described as glazed, and red without papillæ. His general condition steadily deteriorated during the voyage, and in December, some time after leaving Bombay, as he appeared to be sinking. Mr. Low put him on a pure milk diet with a little arrowroot, and whilst on this diet the motions became thicker and the tongue not so red. He began to sleep better and feel better. After a week of the diet the tongue is described as being pale with some papillæ, the temperature remaining normal; pulse about 88.

On January 23rd, the weather became very rough, and the patient was in consequence much worse, shallow ulcers appearing on the tongue and the side of the mouth. With a temporary abatement of the storm the patient at once improved, but the weather in the Mediterranean having become again stormy, he got much weaker, the diarrhoea became greatly aggravated, and he arrived in England in a state of extreme exhaustion and prostration, passing frequent watery, odourless motions. The tongue was pale red, smooth, and free from fur. On the inner surface of the lower lip there was an eruption, herpetic in character, about the size of a fourpenny piece. With care and rest in bed, and milk diet, the diarrhoea diminished in intensity, and the watery motions gave place to yellow feculent discharges. The patient's strength steadily deteriorated, and he became the subject of extreme sleeplessness and feelings of intense *malaise*, and he died exhausted a fortnight after he landed. The progress downwards continued without intermission from the time he came under my observation, the symptoms from sleeplessness and nervous exhaustion being of an exceedingly painful kind. During life a cardiac murmur was recognised. One of the chief features of the case was intense thirst. The mouth, but more particularly the throat, was very sensitive, and occasional small quantities of champagne, from which he got considerable relief, could only be swallowed when considerably diluted with water. Dr. Wethered was so kind as to undertake the necropsy and the histological examination of the tissues. How fortunate it was that this task fell to his hands will be best understood after his report has been read, and by those readers who are most conversant with work of this kind. Dr. Wethered's report is as follows:—

Pathological Report.—The necropsy was made on January 25th, 1890, thirty-six hours after death; rigor mortis was well marked, the body was greatly emaciated; the rib cartilages were ossified. On opening the abdominal cavity nothing unusual was noticed in the position of its contents; the intestines were of a peculiar brownish-yellow colour. On exposing the thoracic cavity, the heart was observed to be almost covered by the lungs, which were emphysematous and rather cedematous. Both pleuræ contained about half a pint of clear fluid, but there were no adhesions or other signs of pleurisy, except at the base of the left lung, which was firmly adherent to the diaphragm, but this condition was evidently of old standing. There was an adherent pericardium, these adhesions also being not of recent origin; they extended all round the heart, and there was about half a drachm of clear fluid between the adhesions. The heart was small in size and firmly contracted; there was still a fair amount of fat round it; the substance appeared normal to the naked eye. The right auricle contained a small quantity of soft red clot, but the other cavities were empty. The mitral orifice admitted the tips of two fingers with difficulty, but no vegetations were seen on the cusps, although the edges were thickened. All the other valves were healthy. The aorta was slightly fatty. The liver was soft and friable, but exhibited no change of structure. There was slight perihepatitis, and at the margin of the right lobe were evidences of some old fibroid change, with dilated bile ducts. There was a fair amount of fat in the omentum. The spleen was soft. Both kidneys were healthy; the capsules peeled easily. The pancreas was rather shrunk. A small calculus was found in the bladder. Digestive tract.—The tongue was clean, papillæ rather enlarged, and the epithelium in some places seemed to be absent. The walls of the œsophagus were a little thickened, but showed no other change to the naked eye. The stomach showed some *post-mortem* digestion, and its surface was covered with an excess

of mucoid-like material. The intestines gave a peculiar velvety feel to the hand, as though there was some thickening of their coats, but this was afterwards shown not to be the case. The walls of the duodenum were thin, and the whole of the interior lined with a thick layer of mucoid material, which could be easily scraped off, leaving the mucous coat of the intestine exposed. The

Fig. 1.—Vertical section through part of the tongue. A, normal epithelium; B, epithelium partially destroyed; C, a point where the epithelium has disappeared; D, partially broken down epithelium (low power).

Fig. 2.—The part marked as B in Fig. 1 as seen by a high power. A, connective tissue of the tongue; B, leucocyte; C, partially detached epithelium. Hartnack VIII. Eyepiece 3, tube in.

walls were so thin as to be perfectly translucent, and the finger could be distinctly seen through them. No other change was apparent to the naked eye. The jejunum and ileum showed similar appearances, the characteristic features being the layer of mucoid-like material and the thinning of the walls. It was this

Fig. 3.—Vertical section through the œsophagus, showing the absence of the lining epithelium. A, mucosa thickly infiltrated with leucocytes; B, muscularis mucosæ; C, submucosa with leucocytes and connective tissue; D, inner layer of muscle; E, outer layer of muscle cut transversely; F, inner surface of œsophagus; G, point in the muscularis mucosæ. OO Verick. Eyepiece 4, tube out.

layer that gave the peculiar feel to the intestines described above. Congestion was absent, there being a few patches of injection

only. The Peyer's patches could not be made out. Passing to the large intestine, similar changes presented themselves. In the rectum the mucoid-like material was very thick, and on gently scraping it off a few abrasions of the coats were seen, exposing

Fig. 4.—Portions of Fig. 3 more highly magnified. The point marked *x* in Fig. 3, showing that the esophagus is lined by a granulation-looking tissue. A, leucocytes; B, connective tissue. Hartnack VIII. Eye-piece 3.

Fig. 5.—The point marked *c* in Fig. 3 more highly magnified, showing unaltered bundles of the muscularis mucosae close to the surface. A, bundles of involuntary muscular fibre (muscularis mucosae); B, leucocyte; C, connective tissue. Hartnack VIII. Eye-piece 3.

the muscular coat. Nowhere was there any trace of ulceration or deposit of fibrinous membrane.

Fig. 6.—Vertical section through the jejunum. A, free surface composed of mucoid-looking unformed material; B, Lieberkühn's crypts unchanged; C, connective tissue of submucosa; D, circular layer of muscular fibre cut transversely; E, external muscular layer; F, crypt showing first stage of destruction; G, inflammatory focus in the mucosa. Hartnack III. Eye-piece 3, tube out.

Microscopic Examination.—The portions of intestine were thoroughly hardened in Müller's fluid, embedded and cut in celloidin and stained with picro-lithium-carmin. *The tongue* (Fig. 1).—The epithelium was seen to be completely destroyed in some places, the destruction gradually passing into healthy tissue. The throwing off of the epithelial cells is well seen in Fig. 2. The muscular layers seemed to be quite healthy, so that the morbid process could not have penetrated deeply. *The œsophagus* (Figs. 3, 4, and 5).—Very marked and extensive changes were noticed

Fig. 7.—Vertical section through ileum. A, unformed mucoid-like substance lining the gut; B, remains of Lieberkühn's crypts; C, strongly developed fibrous tissue taking the place of the submucosa; D, circular layer of muscular fibres cut transversely; E, external layers of muscular fibres. Hartnack III. Eye-piece 3, tube out.

Fig. 8.—Remains of a Lieberkühn's crypt in mucoid-like substance lining ileum. A, membrana propria; B, detached epithelial cell; C, leucocyte. (The cells did not show evidences of breaking up, but there are evidences of loosening from the surrounding tissues.) Verick immersion No. XII. Eye-piece 1, tube out.

here. The epithelium entirely destroyed, and the mucosa densely infiltrated with small round cells. The inflammatory products were seen to have penetrated to the muscular layer; the sub-mucous coat was thickly packed with exudation cells, and the glandular substance to a great extent destroyed. *The stomach and duodenum.*—No abnormal changes were noticed in these. *The jejunum* (Fig. 6).—Effects of the disease were again apparent here. Slight inflammatory action was noticed, as shown by the exudation cells and some destruction of the Lieberkühn's follicles. The epithelium of these seemed to be very little affected. *The ileum* (Figs. 7 and 8).—It is here that the most marked changes were

found. The mucosa was almost entirely destroyed, being replaced by a structureless substance enclosing leucocytes, and here and there the remains of a follicle. These remains are shown under a high power in Fig. 8, in which it will be seen that the epithelium lining the follicle is beginning to be detached. The submucosa appeared much thickened, and the fibrous tissue abundant and more than usually solid in consistence. The walls of the vessels were also thickened, altogether denoting a more chronic process than in the other parts of the intestinal tract. The muscular layer was thinned. The cæcum and rectum showed no changes. Other sections were stained with methyl blue in order to search for micro-organisms. As is usual with sections of intestine, some were found, but they were not more numerous than usual, and no distinctive grouping could be made out.

The pathological changes described by Dr. Wethered are not only new but explain, in an unexpected way, some of the most important symptoms of the disease. Their importance will be best appreciated by those who have had the care of patients suffering from it abroad and in this country. An œsophagus entirely stripped of its lining epithelium and its glandular structures, and an ileum in which the absorbing and secreting surface is practically useless, constitute pathological changes of the greatest gravity and suggestiveness. We now understand how it is that patients complain not only of tenderness of the tongue and mouth, but find great difficulty in swallowing hot or acrid substances. Some of my patients have suffered great distress from their inability to swallow certain kinds of food, and one unfortunate sufferer often spent hours in endeavouring to hawk up morsels of bread which neither would go down nor up, and which, in the light of what we now know, probably stuck in pockets of the gullet unlubricated by the mucus which is needed to smooth the passage of the bolus. Another patient, in whom it was impossible to find a tender spot anywhere else, always complained of a certain uneasiness when the fingers were pressed firmly upwards under the sternum in the direction of the gullet, and occasionally complained of a choking sensation when attempts were made to swallow solid food.

So little seems to be known of the exact changes which take place in the ileum in the process of digestion that it is very difficult *a priori* to understand what the effect would be of the practical obliteration of that part of the gut, but it is quite certain that such a large extent of surface must perform functions of a very important kind. Here again, as in the case of the œsophagus, the pathological changes bear out the results of clinical observation. As the stomach, liver, and pancreas were unaffected, digestion must have taken place in the first instance, but as soon as the bolus descended to the ileum further digestion, at all events in so far as it depended directly on the condition of that gut, would be arrested. As regards absorption, not only is the absorbing vascular surface practically destroyed, but the sub-mucous connective tissue is converted into a thick sclerotic layer which must have interfered with the absorbing functions of any of the deeper vessels that may have remained. Not only so, but the thick coating of mucoid-like substance that covered the free surface of the bowel must have prevented the contact of the contents with the mucosa, such as it was. The result of such a condition must interfere directly with assimilation and nutrition, and these are functions which are profoundly affected in this disease.

The bolus, by the time that it has reached the ileum, has acquired, at all events in the centre, an acid reaction which, to some extent, must be neutralised by the alkaline succus entericus. The destruction of the crypts of the ileum must therefore lead to the failure of the supply of this alkaline juice, and the acidity must by so much be increased, and possibly also the chemical constituents of the bowel contents. With this condition we may with probable justice associate the frothy fermenting appearance of the stools. The large masses of feculent discharge passed by some patients it is also fair to attribute to the want of absorption in the ileum.

Nothing has been discovered in this or in any other case to account for the peculiar pathological changes described, but their wide extent suggests some morbid agent acting from the free surface, leading to a chronic and low form of congestion of the superficial blood vessels with consequent gradual destruction of the epithelial structures. It is not to be forgotten that in the stomach, with its acid secretion, no destruction had taken place. The fact that the stomach, duodenum, and, to some extent, the jejunum, with the liver and pancreas, are practically uninjured accounts

for the long duration, with alternating improvement and relapses, that characterises the history in many of these cases. The acute relapses of diarrhoea to which these patients are all liable after recovery is also explained by the condition of the ileum, because it is probable that even in cases in which apparently good recovery has taken place the membrane, with its vessels, is to some extent permanently impaired, and therefore is easily affected by the ordinary causes of intestinal catarrh.

Space will not at present allow me to pursue the subject further. Before long I hope to resume its consideration, more particularly in its practical aspect. I would only now remark that the fresh knowledge which has been gathered shows how necessary it is that patients who acquire this disease should recognise its gravity in its early stages, and sacrifice everything towards seeking recovery in good climates and by appropriate treatment. On the other hand, I think we ought to recognise that in cases which appear to be hopelessly affected attempts should be made to render what remains of life less painful, and to humour idiosyncrasies more perhaps than has hitherto been done.

NOTE ON MESENTERIC AND OMENTAL CYSTS.

By SIR SPENCER WELLS, BART., F.R.C.S.,
Surgeon to the Queen's Household.

A VERY able paper by Dr. Rasch, read last November at the Obstetrical Society, on a Case of Large Chylous Cyst of the Mesentery, and published in the 31st volume of the *Transactions* of that Society, induces me to bring before the profession notes of two cases which have occurred in my practice since the publication, in 1885, of my last work on *Abdominal Tumours*. In that work there is a short chapter on Mesenteric, Omental, and Pancreatic Cysts, and on Undescended Testicle, with some account of a case where I opened and drained a mesenteric cyst, and of another where I removed, in 1882, a solid mesenteric tumour as large as an adult head. Although this tumour proved to be a sarcoma, the lady remains, eight years after the operation, in good health.

Dermoid Mesenteric Cyst.—The first of the two cases I now record bears some resemblance to Dr. Rasch's case of chylous cyst, but it contained fatty or stearoid matter, instead of chyle, and I have scarcely a doubt that it was congenital. The patient was an unmarried girl, aged 19, whom I saw in October, 1888, with Mrs. Garrett Anderson, who was disposed to regard a central fluctuating abdominal tumour as a large hydatid tumour of the liver. Dulness extended from the lower edge of the liver downwards for about four inches below the umbilicus. Thence to the pubes there was resonance, and vaginal examination excluded the idea of any pelvic tumour. Both flanks were clear on percussion. The cyst or tumour could be pushed rather freely from side to side, not at all upwards or downwards. This and the rather doubtful fluctuation, and absence of hydatid fremitus, or any sign of liver disease, were against the supposition of hydatid disease of the liver, and the history pointed to some congenital tumour of the mesentery or omentum. It was, therefore, arranged that I should make an exploratory incision, and I did so on October 21st, 1888, commencing in the middle line just below the umbilicus, and carrying it downwards for about three inches. On opening the peritoneum and pushing aside a coil of intestine, the mesentery appeared in cyst-like form, and yellowish-white in colour. On dividing the mesentery and holding the edges of the opening carefully forward so that none of the contents could escape into the peritoneal cavity, yellowish semi-solid fatlike matter escaped. More was pressed out, and some bundles of fine hair were also removed by the fingers. Then the cavity, which extended backwards quite down to the spine or root of the mesentery, was thoroughly cleansed by fine sponges. On considering the question of drainage, the difficulty of keeping such a large deep cavity thoroughly aseptic, and the doubt whether reaccumulation would or would not take place, it was decided not to drain, but to close the wound, including the opening in the mesenteric sac with the peritoneal edges of the opening in the abdominal wall. The fatty matter removed weighed six pounds. Some of it was kindly examined for me, at St. Thomas's Hospital, by Mr. Shattock, who reported it to be "a