AN ADDRESS ON ANÆSTHESIA.

Delivered before the International Medical Congress, Berlin, August, 1890.

By H. C. WOOD, M.D., LL.D.,
Philadelphia.

THE most brilliant modern achievements of the science and art of medicine in the direct saving of life are connected with surgery. These great achievements have been rendered possible by two epoch-making discoveries—antiseptics and anæsthesia. The long array of fatal cases of poisoning by carbolic acid, by iodoform, by corrosive sublimate, and by other antiseptic agents; the hundreds of deaths from chloroform, ether, and other anæsthetics, all bear witness to the verity of that strange law, in obedience to which the progress of the human race is so often at the sacrifice of the individual. Antiseptics has outgrown the dangers of its youth, and to-day the measures that are meant to save very rarely kill. On the other hand, the death roll of anæsthesia is daily added to—added to, according to my belief, at a rate that has not changed in forty years. Though this be true, from far-off Australia comes the news that jury and judge have condemned to heavy penalty a chloroformist who had lost his patient; and in England itself a well-known medical journal lends support to such a verdict by affirming that “deaths from chloroform are preventable, that with due care they may be avoided,” and that, therefore, when they occur they are the result of ignorance and carelessness. If this be true, five hundred deaths and more—the result of ignorance or carelessness! Five hundred surgeons, including such names as Billroth, Jaeger, Simpson, McLeod, Agnew, Hunter McGuire, and others of equal rank, guilty of manslaughter! And still the carnage goes on. Surely, under such circumstances, the subject of anæsthesia is worthy of the attention even of this, the most learned medical gathering of the nations that the world can furnish. Antiseptics, the gift of the Old World to humanity; anæsthesia, the gift of the New World, which made the fruits of antiseptics possible—surely it is fitting that I, standing here to-day before you all as the representative of the newer civilisation, should be the chosen mouthpiece for the renewed discussion of this old but pressing theme.

In attempting a fresh study of a well threshed-out subject, I propose to take advantage of the modern physiological methods, and to endeavour to discover by experiments upon the lower animals how anæsthetics kill, and what drugs or measures are most powerful in putting aside their lethal effects. This brings us face to face with the question, How far is it possible to adapt experiments to the needs of practical medicine, and to reason from the dog to the man? A full discussion of this subject would not be opportune, but it does seem necessary for our purpose to devote a few minutes to the pointing out of certain general guiding principles.

It ought to be acknowledged as a fundamental axiom, that no amount of experiments can overthrow a clinical fact, although when a contradiction between experimental and bedside observation seems to arise, such contradiction challenges the correctness of the alleged clinical and experimental facts alike, and should lead to careful re-examination. No amount of failure to purge a dog by elaterium proves that elaterium does not purge man; whilst, on the other hand, the discovery that digitalis increased the blood pressure in the lower animal very properly led to doubt as to the correctness of the, at that time general, belief that digitalis acts upon man as a cardiac sedative, and finally to the recognition of the falsity of the clinical observation upon which such belief rested.

Whatever difficulties may beset the path of the experimental therapist, it is certain that law is throughout the universe supreme; that man, at least in his physical nature, is only an especially developed animal; and if drugs act differently upon different animals, such action must be in obedience to certain laws, to us known or unknown.

Any attempt to fairly discuss these laws would lead us too far afield for the present. One law, however, treads so closely upon the matter at hand this morning, that it requires statement. This law is, that when an apparatus or system is of similar function, and of similar functional activity in different animals, the difference in the action of remedies is very rarely, if ever, in kind,

though it may be in degree. Throughout mammalia the heart has one general structure and one general function; the heart of the dog responds to the touch of digitalis precisely as does the heart of the man. The human brain is so much more highly developed than the brain of the lower mammal that it is, in fact, a new organ or apparatus, and its relation to drugs changes with the change of structure and of function. The scope of this law in regard to anæsthesia is not far to seek. The functions especially compromised in lethal anæsthesia are respiration and circulation. Surely these functions are similar throughout mammalia, and surely we ought to be able to safely reason concerning them from the dog to the man.

Recently, however, alleged clinical facts have been challenged by high authority, upon the strength of experimental results. Under these circumstances, nothing must be at once abandoned, everything must be re-examined. These re-examinations I have made, and I may be pardoned, perhaps, if I affirm that a complete study of the clinical and experimental evidence brings out, not a discord, but a most beautiful concord—that concord between experimental and practical medicine which so often fails to appear simply because we cannot fit together the fragments of truth in our possession.

Although numerous substances have been tried, there are to-day in use practically only three anæsthetics—nitrous oxide, ether, and chloroform. Of these, nitrous oxide stands apart, because it produces loss of consciousness, not by virtue of any inherent properties, but simply by shutting off from the nerve centres the supply of oxygen.

It has been asserted that the changes of circulation produced by the inhalation of nitrous oxide are essentially different from those of mechanical asphyxia, and that therefore nitrous oxide does not act as an asphyxiant. It must, however, be borne in mind that the phenomena of mechanical asphyxia are largely due to the presence of an excess of carbonic acid in the blood, whilst in the asphyxia produced by nitrous oxide there is no excess of carbonic acid, so that the phenomena present are simply the outcome of a lack of oxygen. It is, therefore, *a priori*, to be expected that the phenomena of mechanical and of nitrous oxide asphyxia should differ to a certain extent. To determine the way in which nitrous oxide inhalation affects the circulation, I have, during the past winter, in connection with my assistant and friend, Dr. David Cerna, made a long series of experiments. The result has been to show that usually the inhalation is followed by a rise of the arterial pressure, accompanied by a great disturbance of the pulse; the pulse at first becoming irregular and tumultuous, but by-and-by settling, so that when anæsthesia is complete the pulse wave is remarkably large and full, and the rate very slow. The rise and fall of the arterial pressure in nitrous oxide anæsthesia was found to vary remarkably, not only in different inhalations, but in different periods of the same inhalation. Sometimes the rise was sudden, sometimes it was slow and gradual; sometimes it was maintained until near death, sometimes it was interrupted very early; sometimes it was not very well marked, sometimes it was enormous.

In all our experiments respiration ceased while the heart was still in full activity. Indeed, instead of the gas acting as a cardiac depressant, it appeared to act as a cardiac stimulant, although it paralysed the vasomotor apparatus. Thus, during complete anæsthesia, faradisation of the sciatic nerve always failed to register itself in an increase of the blood pressure, although the heart was beating very powerfully, and although the pneumogastria had been previously severed; whilst late in the poisoning—at a time when the respiration had absolutely ceased, and the animal was in this respect dead, and without the power of self recovery, and when the arterial pressure also had fallen almost to zero—the pulse waves were frequently still nearly three times the normal.

We made but few experiments as to the action of artificial respiration upon the animal dying from nitrous oxide, but these experiments proved that even after complete paralysis of the respiratory function, artificial respiration is capable of rapidly bringing the animal back to life. The heart lives on through nitrous oxide anæsthesia long after the respiratory function has been abolished, and even when the strong, full pulse fails, and the heart has almost ceased to quiver, recovery is still hopeful, because the loss of function has been caused, not by the presence of a poison, but by the absence of oxygen; and although the paralysis may be complete, the life power sleeps before it dies, and is ready to awake at the touch of fresh oxygen.

These experimental results are in strict accord with clinical observations. The S. S. White Dental Manufacturing Company supply a very large, if not the largest, portion of the apparatus and material used for the administration of nitrous oxide in the United States; and, in answer to my inquiry, Dr. J. W. White, their President, writes me that a computation based upon their own sales, and a knowledge of those of their rivals, has reached "the somewhat appalling result, that anæsthesia by nitrous oxide gas is probably effected in three-quarters of a million of cases annually in the United States." Most of these inhalations have been given, not by trained physicians, but by comparatively untrained and often very ignorant dentists; have been given to patients in a sitting or semi-sitting posture; have been given apparently without thought or care to the general community, as the units presented themselves; to the healthy and to the diseased alike; and the result is, out of many millions of inhalations, only three deaths recorded as directly due to nitrous oxide. Could anything be safer?

A suggestive and very practical fact which came out in our experiments is that sometimes during an inhalation of nitrous oxide the rise of the arterial pressure is extraordinary and abrupt. Not long since, in the city of Philadelphia, a gentleman arose from the dentist's chair after an inhalation of nitrous oxide, staggered, and fell in an apoplexy. Is it not easy to perceive that when the arterial system is diseased the great strain of a sudden rise of blood pressure may produce rupture?

Some years since Dr. Kenderdine, a Philadelphia surgeon of local note, died of diabetes, which he insisted was produced in him by the inhalation of nitrous oxide. This is in accord with the researches of the French physician, Dr. Lafont, who reported a case in which sugar appeared in the urine twice in a patient after the inhalation of the gas, and who also caused in himself, and in dogs, temporary glycosuria by such inhalations. Further, Dr. Lafont noticed in a case of mitral insufficiency temporary albuminuria.

I am not aware that these very suggestive statements of the French physician have given rise to any research, except five experiments made recently upon healthy men, with negative results, by two medical students of the University of Pennsylvania, Messrs. George S. Woodward and Alfred Hand, jun. I do not believe that ordinarily the inhalation of nitrous oxide is followed by sufficient disturbance of the circulation to register itself in the urine, but the negative evidence of Messrs. Woodward and Hand is not sufficient to render it improbable that in exceptional cases the inhalation of nitrous oxide may produce albuminuria or glycosuria. Such phenomena, if they occur, are in all probability not directly produced by the nitrous oxide, but are due to the disturbances of capillary circulation caused by it.

However these facts may be, it seems to me that great caution should be used in the administration of nitrous oxide to persons the coating of whose arteries is diseased, and it is probable that when widespread atheroma exists, ether is a safer anæsthetic than nitrous oxide.

When respiration has been suspended in nitrous oxide anæsthesia, the overwhelming indication is certainly for the employment of artificial respiration.

Notwithstanding the great safety and the many advantages which attend the anæsthetic employment of nitrous oxide, the gas can never be used for the general purposes of the surgeon, on account of the excessive fugaciousness of its influence.

The perfect anæsthetic will be a substance which has the power of paralysing the sensory nerve trunks without affecting other functions of the body. If such drug exists it yet awaits the coming of its discoverer. Probably until such a sensory-nerve paralyrant is found chloroform and ether will maintain the complete supremacy which they now have; and in the further discussion of my subject I shall confine my remarks to them. Lack of time limits this discussion to:

1. The method in which these two drugs kill, both in man and in the lower animal, that is, whether they destroy life through the circulation or the respiration.

2. The comparative fatality attending the use of these two agents, and the reasons for the difference.

3. The comparative disadvantages between the two agents, and the best method of securing the desired results.

4. The treatment of accidents occurring during ether or chloroform anæsthesia.

In regard to the method in which anæsthetics kill, my own teaching hitherto has been: First, that although ether in mode-

rate doses acts as a stimulant to the circulation, yet, in overwhelming amount, it is capable of depressing the heart, but that such depression of the heart is always less than the depression of the respiration, and therefore ether kills always through the respiration; secondly, that chloroform may produce death by paralysis of the respiratory centre, or by a simultaneous arrest of respiration and circulation, but that primary paralysis of the heart may occur, and is especially prone to do so when the chloroform vapour has been given in concentrated form.

I think that these views are in accord with general professional belief, but it has recently been alleged that they are at variance with experimental evidences, so that a re-examination is necessary. What, then, are the clinical facts?

If any credence is to be attached to the statements of competent witnesses, who have recorded human deaths during anæsthesia, it is certain that in some cases, under the influence of chloroform, the pulse and respiration have ceased simultaneously; whilst in other instances the respiration has failed before the pulse; and in still other cases the pulse has ceased its beat before the respiratory movements were arrested.

Usually ether arrests respiration in man before it paralyzes the heart, but the collection of records made by Dr. J. C. Reeves certainly show that the fatal result may be produced by syncope. Thus Dr. Ernest H. Jacobs, in a report of a fatal case, asserts positively "the pulse ceased, the breathing continued." It would seem that we must allow that ether in the human subject may cause death in the same methods as does chloroform.

Such then are the clinical facts; or in other words, such are the results of observations made upon the human subject. What are the results of observations made upon animals?

The general teaching in regard to chloroform has been recently challenged by Dr. Lauder Brunton, who, as the result of 450 experiments made by himself upon the pariah dogs of India, has reached the conclusion, as published in the London *Lancet*, that however concentrated the chloroform may be it never causes death from sudden stoppage of the heart. In the physiological laboratories of the University of Pennsylvania, for some years, several hundred dogs have been used annually, and a very large proportion of these dogs have been, at the end of an experiment, killed by chloroform. The observations of Dr. Reichert, Professor of Physiology in the University, Dr. Hobart Hare, Demonstrator of Therapeutics, and myself, have been concordant in showing that chloroform is a cardiac paralyrant, and often does kill dogs by a direct action upon the heart or its contained ganglia. The statements made concerning the Hyderabad Commission, however, led Dr. Hare and myself to a careful and thorough restudy of the subject. Some of our experiments were made by injecting chloroform into the jugular vein; others by administering it by inhalation in the usual way.

The action of the chloroform seems to be not seriously modified by the method of administration. We definitely proved that in the dog chloroform has a distinct, direct, paralyzing influence on both respiration and circulation; that the respiration may cease before the heart beat, or the two functions be simultaneously abolished; but that in some cases the heart is arrested before respiration. We have several times seen the respiration continue as long as one, and even two minutes after the blood pressure has fallen to zero, and the pulse has completely disappeared from the carotid artery.

The correctness of our experiments, we claim, must be acknowledged. The experiments have not only been witnessed by a number of persons, but I have with me to-day tracings which I will gladly show anyone especially interested in the subject. I do not desire to express any doubt whatever as to the correctness of the experimental data of Dr. Brunton; I simply claim that both sets of experiments, although they have yielded different results, have been correctly and properly performed. It may be that the high heat or other climatic conditions surrounding the pariah dog make his heart less sensitive to the action of chloroform than is the heart of the dog bred in northern climates. That the thought of the different constitutions of animals in different climates is not absurd, is shown by the fact that some years ago—after I had affirmed before the Physiological Section of the International Medical Congress at London, that if certain asserted results were obtained upon European dogs, said dogs must differ from those of America, and had been met with a smile of incredulity—Dr. Brown Séquard rose and stated that he had experimented upon hundreds of dogs on both continents, and that there was a distinct difference between the animals, the vascular system of

the European dogs being much more developed, and operations upon them being, therefore, much more bloody than was the case with the American dog.

A very curious parallel might be traced at this point between the experimental and clinical evidence in regard to the effect of climate upon the action of chloroform. In the Southern United States chloroform is used with great freedom, and with great alleged safety; and as long ago as 1878, Dr. Langdon B. Edwards, editor of the *Virginia Medical Monthly*, wrote: "It is one of the most peculiar effects I have ever known in medical practice—the difference of experience in Europe and the North with chloroform and ether as compared with that of the South—the high rate of mortality in the North, and the low rate in the South." Further, in a recent letter to me, Sir Joseph Fayrer affirmed the extraordinary safety of chloroform in India, and stated that he knew of no death from it as having occurred in that country, although its use is universal.

In a series of experiments which I have recently made myself to determine the changes in the circulation produced when ether anaesthesia is carried on to death, I have found that in the first periods of anaesthesia the blood pressure is usually elevated, and that it is usually quite high at a time when the respirations are very shallow and imperfect, and the dark colour of the blood shows that it is heavily charged with carbonic acid. It is not, however, very rare for the blood pressure to remain near the normal, and I have seen the blood pressure begin to fall in the very first stages of ether anaesthesia; moreover, in at least two experiments, death occurred from syncope, the respiration continuing for one or two minutes after the complete cessation of the circulation. In an experiment in which the fall of blood pressure was most pronounced, and the arrest of the heart most complete, the dog was sick from the mangle, and it is possible that the weakened heart was more susceptible than is the normal heart to the depressing influence of ether.

So far, then, as concerns the method in which ether and chloroform kill, I claim most urgently that there is no contradiction between the results as obtained by the bedside and in the physiological laboratories, and that a complete broad study of the clinical and experimental evidence leads to one conclusion, namely, that chloroform and ether are capable of paralysing the respiration and the circulation; that in some cases one function, in other cases the other function, is primarily arrested; but that ether is less prone to produce a primary arrest of the heart than is chloroform.

In the discussion of the second point which I have raised, namely, as to the comparative fatality attending the use of ether and chloroform, I shall not occupy time with any elaborate setting forth of the clinical evidence. In regard to the number of recorded deaths, I shall content myself with accepting the latest statistics at hand, namely, those collected by Dr. Lawrence Turnbull, who has found 375 deaths reported from chloroform, and 52 from ether. I do not believe that these figures nearly represent the total mortality; I doubt very much whether one-third of the deaths from anaesthesia are reported; certainly not one-third of the cases I have had personal knowledge of having been publicly recorded. Moreover, the pressure to conceal deaths from chloroform is greater than when the lethal result is due to ether. The surgeon who uses ether feels that he has employed the safest anaesthetic, and that he will receive no blame if a death occurs from it, and feels also that he has a rare case to put on record, which will give his own name a permanent place in anaesthetic literature; whereas the surgeon who uses chloroform knows that if death occur from the anaesthetic, a very large proportion of the profession, at least in the United States, will condemn him either in public or in secret for the use of this drug, and that he will be fortunate if he escape being publicly condemned by a coroner's jury. Moreover, deaths from chloroform are only too common, so that the surgeon has nothing to gain and much to lose by publication of a chloroform death, and, if possessed of the average human nature, holds his peace.

It seems to me impossible to get at the exact number of anaesthetic deaths, or the proportionate fatality of ether and chloroform. Lyman considers that in regard to chloroform, the ratio of deaths to inhalations is 1 in 5,860; Richardson affirms that it is 1 in 2,500 to 3,000; Andrews puts it for ether at 1 in 23,204; and Lyman at 1 in 16,542.

Without claiming strict accuracy for any of these figures, I think that it can be asserted that the probable ratio of deaths from chloroform is three to five times that of deaths from ether.

When we come to study the effects of chloroform upon the lower animals, we find that it varies very distinctly in its action on the different species. The cat seems to withstand the fatal influences of chloroform with a power worthy of its reputed "nine lives." Many years ago, Professor Schiff called attention to the fact that the use of chloroform as an anaesthetic in the dog is usually attended with the loss of many animals. Professor Martin, of the Johns Hopkins University, writes me that the margin between complete chloroform anaesthesia in the dog, and chloroform death, is a very narrow one. This certainly is our experience in the University of Pennsylvania; we have never been able to use chloroform as an anaesthetic without losing a very large proportion of our dogs.

Clinical and experimental results, namely, the results of experiments made in the physiological laboratory upon the lower animal, and the results of experiments made in the amphitheatre upon the higher animal, man—are again concordant. Chloroform is much more inimical than ether to animal life. The cause of this singular fatality is not, however, chiefly the cardiac action of chloroform. Chloroform is more apt to cause cardiac arrest than is ether, but it is also much more prone than is ether to cause death by failure of the respiration. Almost invariably, when ether is withdrawn before the dog is absolutely in the grasp of death, recovery occurs; but over and over again I have noticed that although the chloroform was taken away whilst the respirations were still being maintained with regularity, the arterial pressure much above zero, and the pulse very apparent, yet the symptoms of cardiac and respiratory failure continued to increase until the fatal issue was reached.

It seems to me that certain general facts or principles in regard to anaesthesia must be considered as established:

1. That the use of any anaesthetic is attended with an appreciable risk, and that no care will prevent an occasional loss of life.
2. That chloroform acts much more promptly and much more powerfully than ether, both upon the respiratory centres and the heart.
3. That the action of chloroform is much more persistent and permanent than that of ether.
4. That chloroform is capable of causing death either by primarily arresting the respiration, or by primarily stopping the heart, but that commonly both respiration and cardiac functions are abolished at or about the same time.
5. That ether usually acts very much more powerfully upon the respiration than upon the circulation, but that occasionally, and especially when the heart is feeble, ether is capable of acting as a cardiac paralyser, and may produce death by cardiac arrest at a time when the respirations are fully maintained.
6. Chloroform kills, as near as can be made out, proportionately three to five times as frequently as does ether, partly, no doubt, because it is more powerful in depressing the heart, but largely because it lets go its hold much less rapidly than does ether when inhalation ceases. Is it not possible that this "holding on" is because it is less volatile than ether, and can we not here get a hint why chloroform is less deadly in the South than in the North? The diffusibility of vapours or gases is in inverse proportion to the square of their densities, and the vapour of chloroform would certainly diffuse itself with far greater rapidity at 90° F. than at 70° F.

The comparative advantages and disadvantages of the two anaesthetics in practical medicine are so well known that only one or two points seem to force themselves upon our present attention. I cannot see that the surgeon is justified in putting the life of the patient to unnecessary risk of chloroformisation, except under special circumstances. I believe, moreover, that much of the unpopularity of ether is due to its improper administration. It is so easy to embarrass the respiration seriously by the folded towel, as commonly used, that not only are the struggles of mechanical asphyxia almost invariably produced, but probably death itself sometimes caused. Especially is there danger of death being thus caused mechanically in the advanced stage of etherisation, when the patient is too thoroughly etherised to struggle, and when the attention of the etheriser is, it may be, attracted by some novel and difficult operation. I confess myself to once having nearly killed a patient in this way.

A proper apparatus is certainly preferable to the folded towel. Various have been invented, but, as the time is short, I shall only mention one, which seems to me a practically perfect mechanism, although it is probably little known this side of the Atlantic.

The inhaler invented by Dr. O. H. Allis is based upon the theory that the patient to be etherised should be supplied with a full abundance of air, saturated with the vapour of ether. It consists essentially of a series of foldings of muslin on a wire framework, arranged almost like the gills of a fish, so as to allow the air to pass freely through, but everywhere come in contact with ether. It should be placed upon the face of the patient dry, and the ether gradually poured on from a bottle with a tight cork, pierced by two small glass tubes, one short for the entrance of the air, the other long, and reaching nearly to the bottom of the bottle for exit of ether. When properly used the Allis inhaler practically does away with the sense of suffocation, and the consequent struggles which have made etherisation alike so repulsive to patient and surgeon.

In order to determine the rapidity with which etherisation can be produced by this inhaler, Dr. M. H. Wilson kept for me notes of thirteen consecutive cases in the clinic of the Jefferson Medical College Hospital in Philadelphia. The average time required for the production of complete unconsciousness was 8 minutes. The average time during which anaesthesia was fully maintained was 32 minutes; and the average amount of ether used during this time was $7\frac{1}{2}$ ounces. In twenty-one surgical cases occurring this spring in the clinical service of Professor J. William White, of the University of Pennsylvania, the average time for the production of complete anaesthesia with ether, used through Allis's inhaler, was $7\frac{1}{2}$ minutes. The results arrived at in these two clinics are so close that 8 minutes must be considered as the average time required for full etherisation by this apparatus.

In discussing the treatment of the accidents of anaesthesia, the results obtained at the bedside naturally press forward for careful consideration; but in going over the subject from this point of view, I have found so little that was novel, and so little that was satisfactory to myself, that I shall not occupy the time of this Congress with any conclusions drawn from reported cases, or personal experience in chloroform accidents. I do not think myself that the problem can be solved by any such study of cases. Death is so near and so terrible, time is so absolute, moments so important, that no surgeon would be willing or justified in waiting for the effect of any one remedy; and when a man is dosed with alcohol, nitrite of amyl, hypodermic injections of ether, digitalis, atropine, and other powerful agents; faradised, slapped, douched, stood on his head, subjected to chest movements for artificial respiration, and various other measures too numerous to mention; who can tell, if by chance he recovers, why he has done so? or who can point out, if by chance he die, what is the remedy whose omission or commission has led to the fatal result?

The problem is a very complex one, not to be worked out amidst the excitement and responsibilities of the amphitheatre. Only in the physiological laboratory can its various elements be separated and studied each by itself, without regard to the individual life which is at stake.

In the physiological laboratory two distinct paths open, each promising to lead to some positive knowledge. We may, on one hand, enter upon the study of the minimum fatal dose of the anaesthetic, and of the results by the consentaneous or subsequent administration of its supposed physiological antagonist; or we may investigate the effect of remedies upon functions that are failing under the influences of the anaesthetic.

The objections to the first of these methods have been, in the present instance, overwhelming. The accidents seem to be independent of the amount of anaesthetic inhaled; and such a method of investigation would have required far more time than was available after I had had the honour of being asked to address this body. Death is produced by chloroform and ether through paralysis of the respiration and the heart, and the method of experimental study which I have employed consisted in a study of the action of powerful agents upon these functions when oppressed by chloroform. I have selected chloroform chiefly because it is the more powerful agent of the two anaesthetics, and the more certain in its lethal results.

The experiments have all been made upon dogs, by one plan. The carotid artery and also the trachea, having been connected with a recording drum, so that the movements of the circulation and the respiration could be consecutively recorded, the animal was anaesthetised, and when the blood pressure had fallen almost to zero, and the respiration had ceased, or nearly ceased, as the case might be, the remedy to be tried was injected into the jugular vein, through a cannula which had been previously inserted.

The more important remedies which have been used by clinicians for the averting of threatened death during anaesthesia are ether, alcohol, ammonia, nitrite of amyl, digitalis, atropine, and caffeine, alterations of position, and artificial respiration.

Although, at least in America, hypodermic injections of ether have been frequently employed even in ether accidents, such use is so absolutely absurd that it does not seem to me to require any experimental evidence of its futility. Ether in the blood acts as ether, whether it finds entrance through the lungs, through the rectum, or through the cellular tissue; and the man who would inject ether hypodermically into a patient who is dying from ether, should, to be logical, also saturate a sponge with the ether and crowd it upon his unfortunate victim.

Instead of simply stating the results obtained in my experiments, I have thought it would be more interesting to show reproductions from some of my tracings. The first drug that I shall report upon is caffeine. I have injected it during the cardiac failure produced by chloroform, in doses varying from 3 to $7\frac{1}{2}$ grains, and have never been able to perceive any distinct alteration in the arterial pressure, and no consistent distinct change of the pulse either in number or force. So far as the experiments go, they certainly indicate that the drug has no influence upon the heart that is being overpowered by chloroform. I may also state here, that it is not possible in any of my tracings to make out any influence exerted by caffeine upon the respiration.

With atropine I have made a few experiments, the results being almost as negative as with caffeine. Ten cubic centimetres of a 2 per cent. solution of the atropine injected into the jugular vein of a chloroformed animal, altered the rate of the pulse beat, but had no apparent effect or influence upon the arterial pressure, or upon the respiration, and in no wise prevented final cardiac arrest.

Of all drugs, that which I think is usually most relied upon by clinicians as a cardiac stimulant in anaesthesia, as in other cases of heart failure, is alcohol. The chemical and physiological relations of alcohol to ether and chloroform are, however, so close, that many years ago I became very doubtful of the value of this drug as a stimulant to a heart depressed by an anaesthesia.

These doubts continually grew stronger from what I saw and read as to the effects of the administration of alcohol during anaesthesia, and were finally changed into conviction by the experiments of R. Dubois,¹ who found that in the animal to which alcohol had been freely given, much less chloroform is required than in the normal animal, to anaesthetise or to kill; or, in other words, that alcohol intensifies the influence of chloroform and lessens the fatal dose.

In my own experiments with alcohol an 80 per cent. fluid was used, diluted with water. The amount injected into the jugular vein varied in the different experiments from 5 to 20 cubic centimetres, and in no case have I been able to detect any increase in the size of the pulse, or in the arterial pressure, produced by alcohol, when the heart was failing during advanced chloroform anaesthesia. On the other hand, on several occasions the larger amounts of alcohol apparently greatly increased the rapidity of the fall of the arterial pressure, and aided materially in extinguishing the pulse rate.

The effects of ammonia upon the failing heart of chloroform anaesthesia have been in my experiments uncertain; sometimes distinct, although very fugacious, and sometimes imperceptible. Twenty cubic centimetres of a 10 per cent. solution of aqua ammonia fortior (*United States Pharmacopoeia*), in some cases produced an immediate rise in the arterial pressure, and even fugaciously registered itself in the respiratory rate, but perhaps more frequently it failed in its influence.

The influence of injections of digitalis has been in a number of experiments very pronounced in producing a persistent gradual rise of the arterial pressure with an increase in the size of the individual pulse rate. In several instances death was apparently averted by its injection, and I saw in one or two cases, where large amounts of the digitalis had been employed, sudden systolic cardiac arrest, indicating that digitalis, if in sufficient amount, is able to victoriously assert itself in opposition to chloroform. Moreover, when I have given chloroform to dogs whose hearts were already under the influence of digitalis, there has seemed to be a peculiar steadying or sustaining power combating the circulatory depression naturally produced by the anaesthetic, and I believe that in all cases of weak heart in man a full dose of digitalis before the administration of chloroform would greatly lessen the danger of cardiac collapse.

¹ *Progrès Méd.*, 1883, xi, 951.

With the nitrite of amyl four experiments were made; in some of these from four to ten drops of the nitrite of amyl were injected in the jugular vein; in others the nitrite was used by inhalation. No distinct effect upon the arterial pressure was in any instance produced, and usually no alteration in the size of the pulse waves, although sometimes the pulse did appear to be a little fuller.

Of all my experimental results, those which have been reached with strychnine have been the most surprising. The injection of strychnine into the jugular vein usually produced a gradual rise of the arterial pressure, and always caused an extraordinary and rapid increase in the rate and extent of the respiration. Thus I have seen the respiration, which had practically ceased for ten seconds, suddenly, under the influence of an injection of two-tenths of a grain of strychnine, become at once very large and full, and reach a rate of 130 a minute.

A series of elaborate experiments made upon the effect of the position of the animal on the blood pressure in the carotid and other arteries, has very clearly proved that the body of the animal whose circulation has been paralysed by chloroform acts in a measure like a tube filled with fluid. Thus, if the feet of the dog were raised vertically above the head, whilst the latter remained upon the table, an immediate rise of pressure occurred, even though the heart had ceased beating entirely; provided that the head of the animal was kept upon a level with the table. If, however, the head of the animal was depressed below the level of the table for a distance equal to, or greater than the length of the body of the animal, a decrease of the arterial pressure occurred at once, although the animal was in a vertical position. The phenomena observed were precisely such as would have been produced if the cannula had been inserted into a tube filled with fluid, instead of the carotid artery, and the elevation and depression of this tube had registered itself on the recording drum, in obedience to the ordinary laws of hydrostatics. The phenomena were entirely independent of any beat of the heart, and were readily produced when the animal was dead, provided the death had not occurred too long previously. Sometimes, even a very few minutes after the cessation of the heart beat, it was impossible to produce the changes of pressure upon the drum. This I believe to have been due to coagulation of the blood occurring very early after death to a sufficient extent to interfere with the liquid properties of the fluid. In no case was any effect upon the respiration produced by change of position in the animal. In a number of cases, however, when the feet were elevated, the heart, which had entirely ceased beating, recommenced its work, and I have several times seen a pulse entirely disappear when the animal was taken from the vertical to the horizontal position. On the other hand, very frequently it was impossible to affect the cardiac action by changing the position of the animal. Nevertheless, the phenomena spoken of occurred too frequently to be a mere outcome of chance, though I several times noted that the heart was usually more affected by alternately elevating and depressing the feet of the animal than by keeping it in a steadily elevated or horizontal position.

When the circulation has practically ceased, under the depressing influence of an anæsthetic, inverting the body must cause the blood which has naturally collected on the enormously relaxed vessels of the abdomen, to flow into the right side of the heart and distend it, and this distension—this increase of pressure—appears at times to have a sufficient momentary influence to stimulate the failing organ.

The theory which has been advocated by some therapeutics—that inversion of the body is of value in the accidents of anæsthesia, because it causes the vital centres of the brain to be supplied with blood—is probably incorrect. The respiration in anæsthesia fails, not through want of blood in the respiratory centres, but because the blood contains a poison which paralyses these centres.

The most remarkable results which I have reached in bringing about recovery of animals to all ordinary intents and purposes dead, were obtained through the use of artificial respiration. Thus, I have seen an animal, in which no respiratory movements whatever had taken place for two minutes, and in which, during that time, no movements of blood had occurred in the carotid artery, and in which, therefore, the heart had ceased its beat, rapidly and permanently restored by artificial respiration.

At one time in these researches, it appeared as though after any dose of chloroform by inhalation, the animal could be resuscitated by artificial respiration, even though heart and lungs were completely paralysed by the drug; but finally I did find a case in which artificial respiration failed.

The results of my experiments with the lower animals may be summed up: that nitrite of amyl, caffeine, and atropine are of little or no use in chloroform poisoning; that alcohol, when given in small amounts, has no influence, but that when given largely, materially assists in paralysing the heart and producing fatal results; that ammonia has some little influence upon the heart, but that of all substances tried, digitalis was by far the most powerful in stimulating the failing circulation; indeed, my experimental results indicate that it is the only known drug which is of any real practical value in such cases. Next, or perhaps even before digitalis, strychnine seems to be of value in the accidents of anæsthesia, because, whilst having some influence on the circulation, it affects powerfully the respiration. For many years chloroform has been used in practical medicine as the physiological and practical antagonist to strychnine, and it seems rather odd that strychnine should never have been employed as the practical antagonist to chloroform.

The one measure which in practical value far surpassed all others for the restoration of the dying animal was artificial respiration, and I have no doubt that a great majority of the deaths which have occurred in man from anæsthesia might have been avoided by the use of an active artificial respiration. The difficulty with artificial respiration as it has been hitherto practiced upon man, after the Sylvester or other methods, is its inefficiency; whereas the artificial respiration which I used on animals was very active, indeed, much more efficient than natural breathing in causing circulation of air through the lungs, and therefore in removing excess of the anæsthetic from the residual air of the lungs and from the blood.

The use of what may be called "forced" artificial respiration by the physiologist, so naturally suggested a similar practice in man, that the celebrated John Hunter invented for the purpose an apparatus which consisted of a bellows so constructed that when it was extended one compartment drew in air from the lungs, whilst the other drew air from the atmosphere; and when it was closed the process was reversed, the fresh air being thrown into the lungs, the foul air into the atmosphere. In 1867, Richardson, of London, invented an apparatus more elegant and portable, although identical in principle with that of John Hunter's; but I have not found that either Hunter or Richardson treated by forced artificial respiration an actual case of disease or poisoning. In 1875,² Dr. John Ellis Blake reported a successful case of aconite poisoning, in which life was apparently saved, although there was no pulse for over three hours, by artificial respiration, with the use of oxygen. In this case Marshall Hall's method was at first used, but later, a small rubber tube was connected directly with a copper reservoir of condensed oxygen, the other end of the tube terminating in a small nozzle, which was inserted in one nostril. Four hundred gallons of oxygen were thus used, but how far the force of the compressed gas was employed to dilate the lungs is not very clear; and it is doubtful whether this case should be considered as one of forced respiration. The first physician to use forced respiration in actual human poisoning, with a clear idea of its value and power, so far as my reading goes, was Dr. George E. Fell.³

It is plain that the bellows constructed by John Hunter and by Richardson are unnecessarily complex and faulty in principle. There is no need whatever of drawing the air out of the fully-filled lungs. Every physiologist knows that when the muscular system is completely paralysed by curare or even by death, the chest walls have sufficient elasticity to force air out of the lungs, and all ordinary laboratory apparatus for artificial respiration is based upon this fact. For forced artificial respiration in man, an ordinary bellows of proper size is all that is required for the motive power.

The real difficulty—the point to be especially investigated and studied—is as to the connection between the bellows and the lungs. Hunter and Richardson simply placed a tube in one nostril, closing firmly the other nostril and the mouth of the subject.

Dr. Fell at first used a tracheal tube, the insertion of which, of course, necessitated the performance of tracheotomy. In one case, however, a simple mask covering the mouth and nostrils was a perfect success. I have had no opportunity of trying the apparatus on the living, but have made a series of experiments upon dead bodies, which have demonstrated that usually a face mask is all that is necessary for the performance of artificial re-

² *Boston Med. and Surg. Journal*, vol. xxxi.

³ International Medical Congress, Washington, 1887.

spiration. Before using the mask, the tongue should be well drawn forward, and, if necessary, fixed in this position by an ordinary piece of silk suture run through it, which can be held in the hand of the operator. If in any individual case the mask fails, an intubation tube may be introduced into the larynx. It do not believe it will ever be found necessary to perform a tracheotomy.

Dr. Fell's apparatus consists of a pair of foot-bellows by which air is forced into a receiving chamber, which is connected with an apparatus for warming the air, and a valve which can be opened and shut by a movement of the finger. This valve in turn leads to the tracheal tube. When the valve is opened the air rushes through the chamber into the lungs and expands them; the finger is lifted, the valve shuts, the lungs contract; and so the respiration goes on. I have no doubt that this apparatus is efficient in practice, but it is open to the serious objection of being unnecessarily complex and costly.

A much simpler, cheaper, and probably equally efficient apparatus may consist simply of a pair of bellows of proper size, a few feet of indiarubber tubing, a face mask, and two sizes of intubation tubes. There should also be set in the tubing a double tube, with opening similar to that commonly found in the tracheal cannula of the physiological laboratory, so that it is in the power of the operator to allow for the escape of any excess of air thrown by the bellows. This whole apparatus can be prepared at a very trifling expense, and it seems hardly necessary to point out its probable value in various narcotic poisonings, and other accidents, in which death is produced by a paralysis of the respiratory centres of temporary nature. The proper use of it—at least with the face mask—could be taught to persons without special medical skill, so that it not only ought to form a part of the surgeon's outfit, but might be of great service in life-saving stations, about gas works, etc.

In conclusion, I may be allowed to state that if the results and deductions arrived at in this address are, as I believe, correct, the rules for the proper treatment of accidents during anaesthesia can be summed up in a very few words:

Avoid the use of all drugs, except strychnine, digitalis, and ammonia.

Give the tincture of digitalis hypodermically.

Draw out the tongue, and raise up the angle of the jaw, and see that the respiration is not mechanically impeded.

Invert the patient briefly and temporarily.

Use forced artificial respiration promptly, and in protracted cases employ external warmth and stimulation of the surface by the dry electric brush, etc., and, above all, remember that some at least, and probably many, of the deaths which have been set down as due to chloroform and ether have been produced by the alcohol which has been given for the relief of the patient.

ON THE TREATMENT OF CHRONIC BRIGHT'S DISEASE.

Read before the International Medical Congress, Berlin, August, 1890.

By T. GRAINGER STEWART, M.D.,

President of the Royal College of Physicians; Professor of Practice of Medicine in the University of Edinburgh; Physician in Ordinary to the Queen for Scotland.

It is no easy task to condense into a short speech so wide a subject as that with which I have to deal, nor is it easy to render interesting a subject so familiar. I should certainly not have dreamed of offering to undertake it, and yet when requested by the Committee I could not refuse: for it is clear that at a Congress such as this, not only is there an opportunity for promulgating new views and original discoveries, but for members of different nationalities comparing the results of their experience in regard to the treatment of important maladies.

I agree with the Committee in thinking the treatment of chronic Bright's disease so important as to warrant its discussion even by those who do not propose to present new views or promulgate important discoveries. The masterly speeches of Senator and von Ziemssen at the recent Congress in Vienna are examples of what may be done in this direction, but rather increase the difficulty by showing how impossible it is to tread in their footprints. I am almost tempted to take their speeches as my text, and go

over the ground which they have traversed, indicating in how far my experience leads me to their conclusions, and in how far I differ from them; but this would presuppose a knowledge on the part of all my hearers of what they have said, which might, perhaps, not be justified. I shall, therefore, deal with the subject in the way which commends itself to me.

Under the term chronic Bright's disease I include all those forms of organic kidney disease which belong to the group first defined by Richard Bright, including therefore diseases of the tubules, of the Malpighian bodies and stroma, as well as of the blood vessels; the parenchymatous and the interstitial nephritis or cirrhosis, and the waxy degeneration as well, for all of these, individually and associated with one another, are referable to Bright's disease. I consider a case chronic when it has lasted more than three months. The indications for treatment may be recognised as consisting in:

1. The removal of the cause or causes, or of some individual cause which still remains in operation.

2. The curing, if it be possible, of the morbid action which has occurred, or is still going on in the kidney, or at least the limitation of the action to the part which has been primarily involved.

3. The removal or modification of the various results, whether of the nature of symptoms or complications, which follow upon the renal lesion, as, for example, the drain of albumen, the dropsy, the uræmic symptoms, the gastro-intestinal symptoms, the serous inflammations, and the vascular and cardiac changes consequent upon the process.

In considering the treatment of any disease, one likes to keep in view the climate and the clothing, the diet and the medicinal treatment; and in discussing Bright's disease it might be most satisfactory if one could speak of each of these in relation to each of the indications; to the causes and to the morbid processes, as well as to the various symptoms and complications bringing out the importance of the different elements of management and treatment in relation to each; but this would be too tedious and perhaps too formal for our present purpose. It will therefore be best if I consider first the general hygienic and dietetic treatment, and then say a few words as to the treatment of the morbid processes, and the symptoms and complications arising from them.

Exposure to cold and damp or to sudden changes of temperature is a fruitful cause, not only of the initiation, but of the prolongation, of kidney disease. Much advantage flows from residence in warm climates and scrupulous avoidance of exposure to cold. Residence in Southern Europe, in Algiers, in Egypt; a voyage to the Cape and a sojourn there, or a similar trip to Australia or New Zealand, or to some of the West Indian Islands, often promotes recovery or hinders the increase of disease. I have no doubt that many patients were formerly sent from home whose cases were really referable to the category of what we now term functional albuminuria, but such a mistake is not likely to happen now with our better knowledge of these conditions, and I have known some of these cases return absolutely uninfluenced by the change of climate.

All will agree that patients suffering from Bright's disease should avoid cold, and especially variable climates, and winter in the South if possible. The remark applies both to the tubular affection and the cirrhotic process, and, seeing how liable the amyloid disease is to be complicated by these, it, of course, applies to it also.

Practitioners who have experience of treatment of phthisis in high altitudes tell us that it is useless to send there patients who suffer from albuminuria as well as phthisis; and while it may be argued that if the albuminuria were simply due to waxy disease, which in its turn is caused by the pulmonary malady, it would be a wise course to treat the kidney by influencing the lung affection. Still experience shows that the practitioners are right; and while the waxy process itself might be benefited, the inflammatory changes which so often attend upon it must be aggravated.

I have seen a good many cases in which malarious influences had to do with the renal disease, and in which, accordingly, it was essential to insist upon removal from the insalubrious districts in which the patients habitually resided, and to keep in view the use of quinine, arsenic, or other antimalarious remedies.

The clothing of patients ought always to be such as to protect from cold and to favour the action of the skin. Woollen under-clothing should be regularly used, and should be changed whenever it is damp from perspiration, and light warm wraps should be carried about, so as to be thrown on if occasion arises. It is