SOME RECENT RESULTS FROM THE STUDY OF LABOUR BY MEANS OF FROZEN SECTIONS AND CASTS.

[WITH LITHOGRAPHIC PLATES.]

Read in the Section of Obstetric Medicine at the Annual Meeting of the British Medical Association, held in Birmingham, July, 1890.

By A. H. F. BARBOUR, M.D., F.R.C.P.E.,

Lecturer on Midwifery and Diseases of Women Edinburgh Medical School, etc.

DURING the past twenty years important light has been thrown on the phenomena of labour, through the study of it by means of frozen sections. During the past year there has been placed at my disposal some very valuable material for this line of inquiry, and I propose now to bring it before you. One preparation is unique; it has never been got before, and, as far as one can judge from probabilities, it is not likely to be stumbled on again. I have to thank my assistant, Dr. Webster, for his invaluable aid, without which the investigation could not have been carried out. The detailed results have been published by us jointly in the Laboratory Reports of the Royal College of Physicians, Edinburgh, for 1889.

Before taking up the preparations we have made, let me say a word on the nature and value of this method, and what it has already done. By freezing we fix parts of the body in the position in which they were at death. We thus get the nearest approach to the condition during life, although there are certain changes that take place at death,¹ for which allowance must be made. By making a section we lay open to view the relations along the plane of the section; such a section shows things exactly as they are—nothing is left to the imagination. It is remarkable how much of what we are accustomed to regard as “knowledge” is imaginary; we get one fact, and fill in all the rest from the imagination. We have only to compare any of the numerous diagrams in obstetrical works with an actual section to see how many erroneous ideas creep in with the single fact which the diagram was intended to express. Take one illustration: the peritoneal cavity is always represented as an open space, which it never is, the peritoneal surfaces being always in apposition unless they are separated by exudation.

By making a series of sections through a pelvis at short distances, and putting the results of the whole together, we render that pelvis practically transparent—we see it through and through; while by selecting any section (say a vertical mesial one), and repeating it in different cases at short successive intervals of time while labour is going on, we can follow the process from beginning to end. Labour is a process implying the movements of a body along a tract. To follow these movements was almost as difficult as to study the movements of a horse running along the road. Photography has solved this last problem by catching the horse's limbs in the different positions, and fixing them for comparison. What photography has done for the horse sectional anatomy will do for labour. Each section is an instantaneous photograph of what was going on immediately before death; thus, the English translation accompanying Chiara's plate of the frozen section of a foetus undergoing spontaneous evolution makes a happy hit by its very literalness when it describes the child as “*caught in the act through corpsy congelation.*”

In working up our material, we have used not only sections but casts. A section gives us relations only along its plane. The range of our inquiry is much extended when we take casts of cavities and masses. In labour we get the whole surface of the genital canal along with its capacity, and also the form and relations of the foetal mass. In the case of two of the preparations, we have for the first time used this method, and it promises to be of great value. It will be most convenient to look separately at the results from sections and the results from casts.

PART I.—RESULTS FROM THE STUDY OF SECTIONS.

1. *Vertical Mesial Sections.*—Plates I and II give an extremely interesting and instructive series of identical sections at different periods in the progress of labour; and in passing from one of these sections to another we can read the changes as if we were

turning over the pages of a history. The sections are, however, at distant intervals of time. The first is during pregnancy at the beginning of the eighth month; in the second, the first stage is commencing; the third shows us the second stage well advanced, the child's head about to be born; in the last, the third stage has finished, the placenta and membranes being newly expelled.

In every case, death occurred from circumstances independent of labour—pulmonary and laryngeal phthisis, heart disease, and cedema of the lungs with hydro-thorax. The second stage case is a most remarkable one in this respect, for the patient's heart failed immediately before the child was born, and when there was no one at hand to interfere and save at least the child. The other two sections of normal labour already published from the second stage (by Braune and Chiara) are with the membranes unruptured, and therefore cannot compare in value with this one. Fortunately for the practice of obstetrics, the chances of such a case occurring again are almost *nil*. It is a very important fact, and one that enhances the value of these sections that the cause of death in each case was independent of and not such as to affect labour.

The drawings which I now hand round are faithful reproductions of the frozen sections both in form and colour and the illustrations which accompany this paper are reduced sketches from them. The full-sized sections will shortly be published in a third edition of my *Atlas of the Anatomy of Labour as Exhibited in Frozen Sections*.² They will also be published separately as a supplement to the former editions of the *Atlas*. It will be best to select one structure at a time and follow it through the four sections, naming these for shortness of reference A, B, C, D, and the following will be found to be of special interest (see plates I and II).

The Bony Pelvis with its Soft Parts.—The antero-posterior diameter for the four pelves, as seen in the vertical-mesial sections, are as follows:—

	Sections	A	B	C	D
		Inches.	Inches.	Inches.	Inches.
Brim	...	4 $\frac{1}{2}$	3 $\frac{3}{4}$	4 $\frac{1}{8}$	4 $\frac{3}{8}$
Cavity	...	4 $\frac{1}{2}$	4 $\frac{1}{4}$	4 $\frac{1}{2}$	4 $\frac{1}{2}$
Outlet (to end of sacrum) ...	...	4	4 $\frac{1}{4}$	4 $\frac{1}{2}$	4 $\frac{1}{2}$

A and C may be regarded as normal pelves, while B shows slightly rickety characters, and D has a longer conjugate than normal. But the chief interest in the frozen section is that it shows to what extent the diameters of the bony pelvis are shortened by soft parts. Information in regard to this can be obtained from frozen sections alone; and as during the second stage the soft parts are altered in position and also compressed, measurements made from sections from this stage will give us the truest idea of the extent to which the available space as measured in the bony pelvis is reduced by the soft parts lining it. Section C is, therefore, of great value as showing this for the antero-posterior diameters, and we find that there is a shortening by soft parts at the conjugate of the brim, posteriorly of about $\frac{1}{2}$ inch, and anteriorly about $\frac{3}{8}$ inch; in the cavity, anteriorly $\frac{1}{4}$ inch and posteriorly $\frac{1}{2}$ inch, and at the outlet (to end of sacrum), anteriorly $\frac{3}{8}$ inch and posteriorly $\frac{1}{2}$ in. Thus there is a shortening at the brim of $\frac{1}{2}$ inch, in the cavity of $\frac{1}{2}$ inch, and at the outlet of $\frac{1}{2}$ inch.

Bladder.—The bladder is distended in A and almost empty in B, but whether distended or empty, it is in both entirely a pelvic organ. In passing I should mention that the vertical extent of the dilated bladder in A does not give an accurate idea of the amount of urine in the bladder, for while the vertical expansion is $1\frac{1}{2}$ inch, the transverse is 5 inches. This shows us how the bladder by transverse expansion accommodates itself to the pressure of the pregnant uterus. While the bladder during the first stage thus remains still a pelvic organ (other sections from more advanced periods in this stage bear this out), in the second stage a remarkable change in its relations takes place, for C shows us that it is now abdominal as well as pelvic. Note that the whole bladder has not become abdominal; the neck of it is still low in the pelvis. The portion within the pelvis is compressed by the child's head, and cannot distend with urine, while the part in the abdomen being free is the part which fills. Hence a small quantity of urine in the bladder produces a remarkably high position of that organ in the lower part of the abdomen, and a long catheter is necessary to reach this part and empty it. It is not the moderate distension of the bladder with urine, however, that accounts for that organ being partly abdominal in C. There is an

¹ For the discussion as to the nature of these changes see the Laboratory Report referred to.

To illustrate Dr Barbour's Paper on Some Recent Results from the Study of Labour, etc.

FIG. 1.—VERTICAL MESIAL SECTION OF A MULTIPARA AT THE BEGINNING OF EIGHTH MONTH PREGNANCY ($\frac{1}{2}$).

Reduced from Pls. XIII. and XVI. of the Author's Atlas, On the Anatomy of Labour as exhibited in Frozen Sections, 3rd Edition.

FIG. 2.—VERTICAL MESIAL SECTION OF A MULTIPARA. FIRST STAGE. FULL TIME (4).

To illustrate Dr Barbour's Paper on Some Recent Results from the Study of Labour, etc.

FIG. 1.—VERTICAL MESIAL SECTION OF A PRIMIPARA. SECOND STAGE. FULL TIME (4).

Fig. 1 reduced from Pl. XVIII. of the Author's Atlas, 3rd Edition.

To illustrate Dr Barbour's Paper on Some Recent Results from the Study of Labour, etc.

PLATE II.

FIG. 2.—VERTICAL MESIAL SECTION OF A MULTIPARA. END OF THIRD STAGE (4).

To illustrate Dr Barbour's Paper on Some Recent Results from the Study of Labour, etc.

PLATE III.

FIG. 1.

FIG. 2.

VERTICAL MESIAL SECTIONS AT END OF THIRD STAGE.

FIG. 1. RICKETY PELVIS (Stratz) (4).

FIG. 2. KYPHOTIC „ (Barbour) (4).

To illustrate Dr Barbour's Paper on Some Recent Results from the Study of Labour, etc.

FIG. 1.

FIG. 2.

FIG. 3.

FIG. 1. TRANSVERSE SECTION ABOVE BRIM. FIRST STAGE (1).
FIG. 2. " " " " SECOND " (1).
FIG. 3. " " " " OF UTERUS AND PLACENTA (PORTO CASE) (1).

Reduced from Pls. IV., VII., XVII., XIX., and XX. of the Author's Atlas, 3rd Edition
(except Figs. 4, 6 and 9).

FIG. 4.

FIG. 5.

FIG. 6.

FIG. 4. TRANSVERSE SECTION BELOW BRIM. FIRST STAGE (1).
FIG. 5. " " " " SECOND STAGE (1).
FIG. 6. " " " " END OF THIRD STAGE (1).

To illustrate Dr Barbour's Paper on Some Recent Results from the Study of Labour, etc.

PLATE IV.

FIG. 7.

FIG. 8.

FIG. 9.

FIG. 7. VERTICAL OBLIQUE SECTION OF EIGHT-MONTH PREGNANCY (1).
FIG. 8. " " " " FIRST STAGE (1).
FIG. 9. " " " " END OF THIRD STAGE (1).

FIG. 1. CAST OF INTERIOR OF UTERUS. EIGHT-MONTH PREGNANCY. SIDE VIEW (1).

FIG. 3. CAST OF FÆTUS. EIGHT-MONTH PREGNANCY. FRONT VIEW (1).

2. CAST OF INTERIOR OF UTERUS. EIGHT-MONTH PREGNANCY. FRONT VIEW (1).

FIG. 4. CAST OF FÆTUS. EIGHT-MONTH PREGNANCY. SIDE VIEW (1)

To illustrate Dr Barbour's Paper on Some Recent Results from the Study of Labour, etc.

FIG. 1.—LEFT HALF OF GENITAL TRACT. SECOND STAGE (1).

FIG. 2.—PLACENTA AND UTERUS FROM SECOND STAGE, SHOWING VENOUS SINUSES (1)

FIG. 3.—CAST OF GENITAL TRACT. SECOND STAGE. SIDE VIEW (1).

FIG. 4 CAST OF GENITAL TRACT. SECOND STAGE. ANTERIOR SURFACE (4).

FIG. 5 --CAST OF FŒTUS. SECOND STAGE. FRONT VIEW (4).

FIG. 6 --CASTS OF FULL-TIME FŒTUSES. (SPIRIT HARDENED) (4).

actual upward displacement of a portion of the bladder in the second stage of labour, so that even if empty it is partly abdominal.³

Passing now to D, at the end of the third stage, we find the bladder again an entirely pelvic organ, its position and relations resembling very much those found in B. The bladder, therefore, when labour is finished, comes to have the same position and relations it had before the second stage began.

Peritoneum.—Along with the bladder the peritoneum between it and the uterus undergoes upward displacement. The utero-vesical fold, which in A and B runs away down into the pelvis, rises in C to the level of the brim. This upward displacement of the utero-vesical peritoneum must be taken along with the upward displacement of a portion of the bladder which we have described, and with the elongation of the cervix uteri and lower uterine segment to be referred to immediately. The peritoneum of the pouch of Douglas also stands in C at a somewhat higher level than it did before the second stage began, being opposite the third instead of the last sacral vertebra, although it has not moved upwards in the same striking way as the peritoneum in front of the uterus has done. In D, after labour, the folds of the peritoneum have returned to the same position as they occupied before the second stage began.

Uterine Wall.—Passing now to the muscular wall of the uterus, we find that before labour (in A) it is of pretty uniform thickness; at the placental site it is rather thinner than elsewhere, and there are also slight inequalities which are due to the pressure of the foetus on the softer *post-mortem* tissues. In every case the cadaver was frozen in the dorsal posture, and it is only in structures which lie posteriorly that we shall find these minor pressure changes—for example, the thinning of the wall in A opposite the sacral promontory.

Comparing now the uterine wall as we see it in A, before labour has begun, with it in D after the uterus has emptied its contents, we find that it has become both thicker and shorter by the process of retraction, the nature of which we shall have to refer to immediately. In D both walls, as far down as the retraction ring, have increased remarkably in thickness⁴—to four times that before labour began—the fundus being not quite as thick as the anterior and posterior walls. With this increase in thickness there is a diminution in circumference, although not in the same ratio. We cannot compare exactly the length of the uterine walls before and after labour, because the circumference before labour does not include the cervix, and the thinner portion of the wall becomes folded up after delivery, so that it cannot be accurately measured; but we may say roughly that the loss in length is only about one-third, as against an increase in thickness of four times. A glance at the frozen sections will bear out this point that the increase of thickness is very much greater than the diminution in length.

These sections give us an exact idea of the change in the uterine wall produced by retraction in labour. We have now to inquire whether this process affects the whole wall throughout, and to do this we must look at A and D more carefully, and also at B and C. Although we spoke of the wall in A as being of uniform thickness. There is one exception to this general uniformity of thickness, namely, the lower portion of the anterior wall, which becomes distinctly thinner before it passes into the cervix. In B we find the relative thinness of this part even more marked; while in C it is so pronounced that the anterior wall might be sharply divided into two portions, an upper thick portion and a lower thin one, the line of demarcation being marked in C "retraction ring." In C the posterior wall has undergone a similar change, although there was no hint of its occurrence in A and B; in fact the posterior wall is in these sections as thick, if not thicker, in its lower portion than it is in the part above.

³ As first pointed out by Dr. Berry Hart from his study of Braune's section.

⁴ The actual measurements are:

Section A.		Section B.	
In.		In.	
Anterior wall...	$\frac{1}{4}$ (falling at placental site to $\frac{1}{16}$)	$\frac{1}{16}$	(p. s. $\frac{3}{16}$)
Anterior lower segment...	$\frac{1}{4}$	$\frac{1}{16}$	
Fundus ...	$\frac{1}{4}$	$\frac{1}{16}$	
Posterior wall ...	$\frac{1}{4}$ to $\frac{1}{2}$	$\frac{1}{4}$ to $\frac{1}{16}$	
Posterior lower segment...	$\frac{1}{16}$	$\frac{1}{4}$	
Section C.		Section D.	
In.		In.	
Anterior wall ...	(placental site $\frac{1}{4}$ in.)	1 to $1\frac{1}{2}$	
Anterior lower segment ...	$\frac{1}{4}$	Folded up	
Fundus ...	$\frac{1}{4}$	$\frac{1}{4}$	
Posterior wall ...	$\frac{1}{4}$	$\frac{1}{4}$ to $1\frac{1}{2}$	
Posterior lower segment ...	$\frac{1}{4}$	Folded up	

Before labour, then, we have the anterior wall thinner in its lower portion to begin with, while during labour this thinness increases, and a corresponding thinning of the posterior wall takes place. The uterine muscle, therefore, in the process of labour becomes differentiated into two portions—an upper, which becomes thick, and a lower, which becomes thin. This thinning portion is passive, and therefore more allied to the cervix and vagina below (which tend to elongate and thin in labour), than to the active contracting muscle above. Functionally, therefore, we should divide the genital tract in labour into two parts—an active, shortening, thickening portion, and a passive, lengthening, thinning one—the line of separation forming a ring in the tract. Now, what are we to call this ring? The patient from whom preparation C was taken did not probably die during a pain, that is, during a contraction (for the axis of the uterus is not thrown forward into the axis of the brim of the pelvis as it is in Braune's section), and yet the uterine muscle is relatively thick. Passing to Section D, we find the muscle here also thickened in proportion to the diminution of the uterine cavity, and this is the case with all *post-partum* uteri. This shows us that there occurs in the uterine muscle in labour a change by which it remains permanently shorter and thicker after a transient contraction or pain has passed off; and this permanent change is "retraction," which you will see is a different thing from contraction. In pushing a book intermittently along the table by means of the arm muscles, each muscle returns to its former length and thinness in the intervals of rest; but on driving the foetus along the genital tract the uterine muscle does not return to its former length and thinness between the pains, but to a great extent retains the dimensions it has acquired, so that it becomes progressively shorter and thicker. Now it is this change, and the extent to which it has occurred, that we see registered in such a section as C, and therefore I think the term "retraction ring" more suggestive as well as more accurate than the term "contraction ring," which has also been applied to it. This differentiation of the uterus into two parts is even more distinctly seen after the work of labour is done, as in Section D, where the thickening of the uterine wall is evident. The thinning of the part below is not so well marked in D, as the uterus has sunk down into the pelvis.

That the extent to which we see this thin portion in the uterus after labour is affected by the position of the uterus will be evident if we compare with the section in D those given in Plate III, also of uteri immediately after delivery and representing the two extreme types of pelvic deformity. In Plate III, Fig. 1, where the conjugate is shortened by rickets to 3.2 inches, the uterus is kept up in the abdomen, and the lower segment is thus drawn out; while in Fig. 2, where the conjugate is elongated by kyphosis to 6.6 inches, the sliding down of the heavy uterus has been favoured by the long, sloping sacrum, and the flabby lower segment is folded up below. The lower segment is thus more allied to the cervix and vagina; and physiologically we might divide the genital cavity into two parts, an upper active and a lower passive one, the line of demarcation being the retraction ring. The recognition of the fact that the genital tract consists of these two parts is of the greatest importance both for normal and pathological labour. For normal labour, we see that as soon as the contents of the uterus pass below the retraction ring they cannot be propelled forward by the muscular action of the walls of the tract, but only by the abdominal muscles. Looking at C, with regard to the force which will cause the child to be expelled, we note that the contents of the uterus acted upon directly by the muscular wall comprises only the part of the foetus above its arms; and further, that the breech is not in immediate relation to the uterine wall, but separated from it by liquor amnii. The part of the foetus below the retraction ring is acted upon indirectly through the part above it; but once the foetus has passed wholly below the retraction ring, the muscular action of the abdominal wall alone will operate on it.⁵ So also with regard to the expulsion of the placenta.⁶ Looking at Section D, we can see that if the placenta be above the retraction ring, the uterine muscle will act upon it, but if it be below that, only the abdominal muscles can act. We might compare the emptying of these two portions of the tract to the emptying of the bladder, or by the straining of the abdominal muscles where the bladder walls have lost their tone.

⁵ Schröder has drawn attention to this in his *Der schwangere und kreisende Uterus*, etc. Bonn, 1888.

⁶ See my former paper on the Anatomy of the Third Stage of Labour, *BRITISH MEDICAL JOURNAL*, February 11th, 1888.

As regards the pathology of labour, the passive character of the lower segment has a bearing both upon rupture of the uterus⁷ and according as that is done by the action of the bladder wall alone placenta prævia.⁸

Uterine Cavity and Contents.—The placenta, membranes, and foetus we shall refer to under "transverse sections" and "casts" respectively. As regards the question of cavity, a glance at the four Sections A, B, C, and D, shows that there is nowhere at any period of labour any empty space in the genital canal. The walls are always in apposition throughout unless separated by contents. The appearance of dilatation in the lowest third of the vagina in A, B, and D, was produced by a cotton plug placed in the vulva to mark its position. The appearance of a cavity in the *post-partum* uterus, as in D, is due to blood clot. Compare also the two sections in Plate III, in one of which the walls are in apposition, while in the other they are separated by blood clot. The fact that the walls of the genital tract are in apposition before labour and come into contact again as the contents are being expelled, is significant in its bearing on the exclusion of septic air.

2. *Results from the Study of Cross Sections.*—In Plate IV, Fig. 2, we have a section above the brim, taken from Case C, the patient who died during the second stage. For the sake of comparison with it, I have put alongside corresponding sections that I have made in other cases; Fig. 1 is from a first-stage case (not Case B), and though it does not show the abdominal wall nor intestines, it runs through the uterus and spinal column in much the same position as the second stage one; Fig. 3, is from a Porro uterus, in which uterus and placenta are cut through at a corresponding level; and Fig. 6 shows a section of the left half of the pelvis from a case dying immediately after delivery, this section again dividing the uterus antero-posteriorly, though, of course, to do this it has to pass through the cadaver at a much lower level, being below the brim instead of above it. Now I can scarcely conceive a more instructive picture of the changes in the uterus, as seen in cross section, which result from the process of labour. In Fig. 1 you have a muscular bag with thin flaccid walls lying moulded on the spine, containing liquor amnii and foetus, the liquor separating the parts of the foetus from each other and from the uterine wall, and having the placenta covering rather less than a third of its circumference. We see beautifully how much nearer the back of the foetus is to the uterine wall than its chest is, and how the foetal parts in section differ from the smooth back, giving us an anatomical basis for the auscultation and palpation of the pregnant uterus. In Fig. 2 we see the change in the size and consistence of the bag which the removal of the greater part of the liquor amnii has brought about; for the uterus has not been wholly emptied of the liquor amnii by the rupture of the membranes. The uterine wall is thicker, and though slightly indented by the spine, is not moulded on it. It and the placenta are in contact with a considerable portion of the foetus, the trunk of which forms the main part of the section, the limbs not appearing to the same extent as in Fig. 1, for a reason to be immediately explained. The placental site has also shrunk somewhat, and the placenta become thicker, though neither it nor the membranes are separated (determined by examination of the thawed specimen, though, of course, not demonstrable in a frozen section). Fig. 3 shows us uterine walls still more increased in thickness, and uterine cavity filled only with the placental mass, which has the walls everywhere in contact with it. Fig. 6 shows the uterine walls still more thickened, the cavity very much shrunk (for we have about one-half of it in this section), and the walls of the cavity in apposition, with the exception of a small amount of blood separating them.

The study of these sections brings out three points: The remarkable change in the thickness and consistence of the uterine wall which results from retraction; that the walls are always in apposition except for the contents; and that the placenta and membranes (we are not speaking at present of the membranes over the lower segment of uterus, we are speaking only of those over the retracting portion) remain unseparated until they form the only structures left in the uterine cavity, while after their removal, as already said, the uterine walls come into apposition.

The cross sections, given in Plate IV, Figs. 4 and 5, are from cases B and C (the first and second stages), and show us the relations of the foetal head to the pelvis; in Fig. 4 it lies almost trans-

verse, while in Fig. 5 it is partially rotated, being out of the right oblique though not yet in the conjugate diameter. In both cases it pretty well fills the pelvis. In Fig. 4 the cutting out of a block to preserve a vertical mesial section prevents us from following the sutures; but in Fig. 5 we see the margins of the occipital bone dipping below both parietal bones, while the relation of the frontals to the parietals and to each other is not affected. The head in this latter section blocks the pelvis, reducing the vaginal wall to a thin band, and pressing the rectum against the sacrum. Along with this latter section (Fig. 5) we should look again at Plate II, Fig. 1, in which the anterior parietal bone is bulging towards the vaginal orifice, and the plastic nature of the foetal skull is seen even in the altered shape of the two halves of the brain. The foetal head, therefore, in the second stage plugs the pelvis, bulging in the direction of least resistance; and the occipital bone dips below the parietal, but the relative levels of the bones at the other sutures are not affected. Further, the position of the caput succedaneum (see Plate II, Fig. 1) is immediately over the part of the scalp that is unsupported by the soft parts of the mother. This description of the alteration in the position of the bones in the moulding of the head does not correspond in all points with that usually given, but we must remember that the latter is that of the child's head *after* birth. This is the only foetus that has been described with the moulding actually going on during the process of birth. Thus, the parietal bones are generally described as overlapping. Now they are undoubtedly found to overlap after delivery, and it might quite well happen that after the head of this foetus had left the vulvar orifice, the parietal bones would take up an overlapping position. But this section shows that the head is reduced in its transverse diameter, not by the pushing-in of one parietal bone below its fellow, but through the plasticity of the head as a whole; the distortion towards the point where it is unsupported by the soft parts being most remarkable. There is, however, an actual dipping of the occipital bones below the parietals.

3. *Results from the Study of Oblique Sections.*—Figs. 7, 8, and 9 of Plate IV give us three sections in the oblique diameter of the pelvis, Figs. 7 and 9 being in the right oblique diameter, and Fig. 8 in the left. As the head of the child usually enters the pelvis in one of the oblique diameters, it is of importance to know to what extent the length of these diameters is affected by the soft parts—a point with regard to which we have as yet no data, and which only frozen sections can settle. Without burdening you with detailed measurements⁹, we find that, excluding the loop of intestines in Fig. 8, the thickness of soft tissues at both ends of the oblique diameters is in all about seven-eighths of an inch; and, therefore, we may say that in estimating the available space in the oblique diameters of the brim, we must deduct almost an inch from the measurement as made on the bony pelvis.

Below the brim the shortening becomes more marked, through the obturator internus and the cellular tissue of the broad ligament containing large venous sinuses. The oblique section made after delivery (Fig. 9) is of no use in giving data for the shortening of the diameters by soft parts, as the uterus and its cellular tissue have sunk down into the pelvis, but it brings out this very important point that the large venous sinuses which were so prominent a feature in Figs. 7 and 8, are in Fig. 9 so compressed as to be scarcely visible. Figs. 7 and 9 form, indeed, a most interesting comparative study when placed side by side, as showing the change in the appearance of a section running through an oblique diameter at these two periods, the carrying down into the pelvis of the cellular tissue of the broad ligaments with the sinking uterus, and the compression of the venous sinuses being the most striking features.

⁹ In Plate IV, Fig. 7, we have at the posterior end of the right oblique diameter $\frac{3}{4}$ inch of soft parts, 3 being uterine wall and 2 cellular tissue; and anteriorly there are about $\frac{3}{4}$ inch divided equally between the wall and the tissue outside of it. In Plate IV, Fig. 8, at the posterior end of the left oblique diameter, there are $\frac{3}{4}$ inch of soft parts, 2 being uterine wall, 2 cellular tissue, and 3 intestine with its mesentery, while at its anterior end we have $\frac{3}{4}$ inch, 2 being uterine wall, and 1 extrauterine cellular tissue. Lower down, anteriorly, we have in each case the obturator internus muscle encroaching on the pelvis, forming a muscular pad $\frac{1}{4}$ to $\frac{1}{2}$ inch thick. We may say, therefore, that at the ends of the oblique diameters we have anteriorly from $\frac{1}{4}$ to $\frac{3}{4}$ inch (in a former paper I estimated this from other sections at $\frac{1}{2}$ inch), and posteriorly from $\frac{1}{4}$ to $\frac{3}{4}$ inch, making in both cases about $\frac{3}{4}$ inch in all.

(To be continued.)

⁷ Bandl's classical monograph on *Rupture of the Uterus* first emphasised the thinning of the uterus before rupture. What we have called the retraction ring becomes "Bandl's ring" in pathological labour.

⁸ As Berry Hart has pointed out; *Edin. Obs. Trans.*, xiii, 113.

THE Hutchinson Prize at the London Hospital Medical School has been awarded to Mr. John Thomas, of the Infirmary, Cardiff.