infection, and (b) twenty cases, resistant to the older methods of treatment, receiving uleron at a late stage of infection.

3. The first group, after a period of observation including test of cure, gave 74 per cent. cures, 17 per cent. failures, and 9 per cent. relapses. The second group gave five "failures" out of the twenty cases.

4. With the dosage used in this series (11 grammes per three-day course) no alarming drug reactions were observed. Three complained of slight headache, four of slight nausea, and two developed a generalized urticaria, which cleared up in about a week on cessation of the drug. No polyneuritis was encountered.

I have pleasure in expressing my thanks to Mr. A. E. Robertson, a member of my staff, for the valuable assistance rendered in the investigation of my cases.

REFERENCES

- Bowie, F. J. T. (1938). *British Medical Journal*, **2**, 283.
 Cokkinis, A. J. (1938). *Ibid.*, **1**, 1151.
 — and McElligott, G. L. M. (1938). *Lancet*, **2**, 355.
 Felke, H. (1937). *Dtsch. med. Wschr.*, **63**, 1393.
 Fischer, C. (1937). *Fortschr. Ther.*, **13**, 553.
 Fuhs, H., and Volavsek, W. (1938a). *Wien. klin. Wschr.*, **51**, 586.
 — (1938b). *Derm. Wschr.*, **107**, 1057.
 Gennerich, W. and K. (1938). *Ibid.*, **106**, 541.
 Grütz, O. (1937). *Münch. med. Wschr.*, **84**, 1201.
 Hanschell, H. M. (1938a). *Brit. J. vener. Dis.*, **14**, 18.
 — (1938b). *British Medical Journal*, **2**, 375.
 Harrison, L. W. (1938). *Ibid.*, **2**, 90.
 McGregor-Robertson, J. G. (1938). *Glasg. med. J.*, **130**, 1.
 Mergelsberg (1938a). *Dtsch. Militärärztl. Z.*, **5**, 206.
 — (1938b). *Ibid.*, **8**, 363.
 Nicol, H. (1938). *Med. Pr.*, **197**, 168.
 O'Hanlon, O. J. (1938). *J. R.A.M.C.*, **70**, 183.
 Prebble, E. E. (1938). *British Medical Journal*, **2**, 591.
 Walsh, D. F. (1938). *Ibid.*, **2**, 215.

TREATMENT OF STERILITY

REPORT ON SIXTY-TWO CASES

BY

M. MOORE WHITE, M.D., F.R.C.S.

First Assistant, Gynaecological Department, Royal Free Hospital

This investigation has been undertaken because on reviewing a series of forty-one cases which had been admitted to the gynaecological wards of the Royal Free Hospital for the treatment of sterility in 1936 it was found, on inquiry one year to eighteen months later, that only five (12 per cent.) had become pregnant. A dilatation, insufflation, and curettage was performed in each case, seventeen receiving in addition injections of hormones.

It was felt that these figures did not justify treatment in the wards and that perhaps similar results might be obtained if sterility cases were investigated and treated in the out-patient department. Between August 1, 1937, and May 31, 1938, sixty-two consecutive women who attended Miss Dearnley's out-patient department for sterility have been investigated and treated; forty-seven complained of absolute sterility, nine of one-child sterility, one that she could not become pregnant with a second husband (although she had had four children by the first husband, and the present husband had healthy spermatozoa as demonstrated by the seminal test), and five that they had had one or more miscarriages.

Routine Investigation of Sterility

A careful history was taken, and the duration of the marriage, the use or not of contraceptives, and any menstrual abnormality were noted. A routine Wassermann test was made, being found negative in every

instance. Nothing abnormal was detected on pelvic examination except retroversion in three cases, a unilateral hydrosalpinx in one case, and an undersized uterus in four cases. In each case the husband was asked to send a specimen of semen for examination, and thirty-nine of the sixty-two consented to this. Five (13 per cent.) showed no spermatozoa, and three (8 per cent.) very few and distorted specimens. Thus in 21 per cent. of the examined cases the husband was sterile. On account of the long distances many of the specimens had to be sent for examination no regard was paid to absence of motility provided the specimen was in all respects normal.

Uterosalpingography

Partly as a means of diagnosis regarding patency of the Fallopian tubes, and partly as a therapeutic agent, lipiodol was injected into the uterus. It was thought that this procedure might separate fine adhesions in a non-patent tube and that in view of the possible antiseptic action of the oil the endometrium would be left in a healthier state. No anaesthetic was given. The week following the end of the period was in most cases the time chosen to perform the test.

Technique

The patient was asked to lie in the dorsal position with the knees bent on a sterile towel on the x-ray table; a sterile bivalved speculum was then passed and the cervix painted with iodine. A uterine sound was employed to ascertain the direction of the uterine cavity; the two smallest-sized dilators were passed, so as to stretch the internal os gently if this was found tightly closed; and finally the insufflator (a modification of Provis's model with an oval plug, made for me by Messrs. Hawkins) was introduced. No vulsellum was used, but with the plug firmly pressed against the cervix 10 c.cm. of lipiodol was slowly injected and a radiograph taken.

It has been found that, provided the insufflator can be introduced past the internal os and the cervix is not unduly patulous, the lipiodol enters the fundus and passes into the tubes. Pain may be felt similar to that experienced during a retrograde pyelography, but not uncommonly there is no discomfort at all. In one case only of this series has there been any infection. This patient had a low temperature for a few days, and according to her private doctor she felt some pain, and there was tenderness in the pelvis on vaginal examination. Most of the patients attended twenty-four hours later for a second radiograph in order to demonstrate whether lipiodol had escaped through the Fallopian tubes into the pelvis. In twelve cases (19 per cent.) no lipiodol had entered the peritoneal cavity, suggesting absence of patency, and in two others it was doubtful whether there was a trace after twenty-four hours, but in both of these doubtful cases pregnancy occurred.

Since it is perhaps not generally agreed that one negative insufflation or lipiodol test should be accepted as conclusive evidence of occlusion of the Fallopian tube, the twelve negative cases (Table I) will be considered in further detail. In no instance where non-patency was demonstrated radiographically has the patient become pregnant. In one case the findings were confirmed at operation, and three cases were negative also to insufflation, but one of these, judging from the kymograph tracing, was rendered patent on one side by this procedure. Of the remainder, in two cases x-ray examination showed a unilateral hydrosalpinx, and in three there is a trace of lipiodol in one tube only; one of the latter had had a salpingectomy for tuberculous salpingitis two years

ago. Extensive destruction of the right hip-joint, the sequel to an acute arthritis, was depicted radiographically in another. Clinically, of the remaining two, one had a small acutely retroverted uterus and the other appeared normal, so that in this series the radiograph probably revealed the true state of affairs.

It is interesting to note that, although in twelve cases there was absence of patency and in nine others failure of lipiodol to enter one tube, in only two was there a history of previous inflammation with removal of a tube, and one other case had a history of removal of an ovarian cyst on the side with non-visualization of the tube. It is clear, therefore, that factors causing tubal occlusion

TABLE I.—*Record of Findings in 12 Patients with Non-patent Fallopian Tubes*

Name	Age	Parity	Lipiodol in Uterus only	Lipiodol in Tubes	Insufflation	Remarks
B. J.	31	0	+		—	
E. S.	34	1	+			Small retroverted uterus
L. A.	22	0	+			Confirmed at operation : plastic reconstruction operation performed. Right now patent
M. W.	25	0	+		—	Small acutely anteverted uterus
M. T.	39	1	+			Radiographs show extensive destruction of right hip-joint
E. S.	39	0		L. + ; R. +		None in pelvis
M. G.	38	0		L. + ; R. —		Left hydrosalpinx
K. F.	31	0		R. + ; L. —		Right hydrosalpinx
R. W.	32	1		L. trace ; R. —		
F. E.	37	0		R. trace ; L. —		
E. G.	32	0		R. $\frac{1}{2}$; L. —		Left salpingectomy for tuberculosis two years ago
E. E.	31	0		R. + ; L. +	— +	Bilateral hydrosalpinx. Insufflated since lipiodol test, sudden pain experienced at pressure of 140 mm. Hg. and air heard to escape on right side

can operate apart from symptoms. Two radiographs showed some free fluid in the peritoneal cavity, as evidenced by the lipiodol collecting in the form of globules.

A follow-up x-ray examination has recently been made of five of the early cases: four were chosen in which the films showed a large quantity of lipiodol in the pelvis, and one in which there was a bilateral hydrosalpinx. In two no trace of residual opaque material was found, in two others there were traces in the pelvis and possibly a trace in one tube, and in the hydrosalpinx case traces were still present in both tubes. In no instance was there any evidence of residual lipiodol under the diaphragm or any symptoms referable to the presence of this substance in the body. In no case in this series did the radiograph show that lipiodol had entered the pelvic veins (Kilroe and Hellman, 1933; Coventry, 1934; Pujol y Brull, Vanrell, and Riera, 1929).

Hormonal Therapy

It was originally intended to give a course of injections of dimenformon (prognon-B oleosum) twice a week in the two weeks following the cessation of the menstrual period, over a period of three months, but either because of the inability of the patient to attend so often or on

account of the cost of the injections, only twenty-two received this treatment, and in many instances the course was incomplete. Of this number, nine (41 per cent.) became pregnant (Table II). Six women were given a

TABLE II.—*Record of Women Pregnant Following Lipiodol Injection plus Administration of Endocrines*

Case No.	Name	Age	Number of Children or Miscarriages	Years Married or since Last Child	Years Contraceptives used	Report on Husband's Semen	Patency of Tubes		Months elapsed before Pregnancy	Miscarried (Number of Months Pregnant)	Remarks
1	S. L.	22	0	9	0		+	7 spaced injections of 10,000 I.B.U. prognon over 2 mths.	3 $\frac{1}{2}$	2	
2	L. T.	27	0	2 $\frac{1}{2}$	0	+	+	8 twice-weekly injections of 20,000 I.B.U. dimenformon	6		Small anteverted uterus and scanty menstrual loss
3	A. Y.	37	0	7	0		?	6 twice-weekly injections of 50,000 I.B.U. dimenformon	3		Small anteverted uterus and scanty menstrual loss
4	K. R.	31	0	5	0	+	+ R.	8 spaced injections of 10,000 I.B.U. of prognon over 2 mths.	2	2	Periods irregular lately
5	E. L.	36	0	11	0	+	+	Hormonal shock therapy twice; 50,000 I.B.U. dimenformon	3		See text. Pregnancy followed second injection
6	V. S.	31	0	5	0	+	+	Hormonal shock therapy once; 50,000 I.B.U. dimenformon	1		
7	E. W.	37	0	8	0	+	+	2 injections of 10,000 I.B.U. prognon	5	1 $\frac{1}{2}$	Intolerance to injections. Low sugar tolerance. Given spaced carbohydrate diet
8	F. C.	28	1 mis.	3 $\frac{1}{2}$	0	+	+ L.	8 spaced injections of 10,000 I.B.U. prognon + hormonal shock therapy twice	2 $\frac{1}{2}$		Pregnancy followed second "hormonal shock" injection
9	D. B.	28	1 mis.	1 $\frac{1}{2}$	0	+	+ R.	Hormonal shock once. Not followed by pregnancy. Corpus luteum gr. x t.d.s. before the missed period	2		

large dose (50,000 I.B.U.) in the form of dimenformon on the twelfth day of the menstrual cycle, as this "hormonal shock" therapy is thought to ensure ovulation (Clauberg, 1935), and in four instances pregnancy followed. Two women with a history of short menstrual periods and scanty loss were given injections of dimenformon twice weekly—in the one case 10,000 I.B.U. for four weeks, and in the other 50,000 I.B.U. for three weeks. Both women became pregnant.

Additional Investigations

In selected cases the basal metabolic rate was estimated, and to two women in whom it was found to be low 1 grain of thyroid extract was given orally twice daily. Three women in whom the uterus was found to be small were given $\frac{1}{2}$ grain of thyroid extract and 1 grain of pituitary extract (whole gland) orally in addition to hormonal injections, when after two months they had failed to conceive. Two of these patients became pregnant.

When there was a history of miscarriages the patient was sent to the biochemical department and a glucose-tolerance test was made; if a lowered sugar tolerance was found, the carbohydrate intake was regulated and spaced. One patient who had been on a diet for only five weeks miscarried. A second who miscarried was found to have a lowered sugar tolerance. It is known that diabetic women become pregnant only with difficulty, unless under treatment, and readily miscarry. Minor degrees of deranged carbohydrate metabolism may therefore be operative in some of these patients (Williams, 1933).

Advice was given as to the most opportune time for intercourse: if the menstrual cycle was normal the twelfth to the seventeenth day, reckoned from the first day of the last menstrual period, was advocated, though individual cases in my series have shown that conception can occur just before a period. (One patient was injected with lipiodol seven days before the period, had intercourse two days later, and failed to menstruate; she was subsequently found to be pregnant.) In those cases in which the patient gave a history of an excessive mucoid discharge she was advised to syringe with a solution of bicarbonate of soda, one drachm to the pint, in the early evening. Four patients who did not become pregnant within the first two or three months were given vitamin E in the form of one capsule of wheat-germ concentrate three times daily. Three subsequently became pregnant, two of them miscarried.

Results

Between August 1, 1937, and May 31, 1938, sixty-two patients were treated, and were injected with lipiodol. Up to the time of writing (September, 1938) twenty-two patients have become pregnant—that is, 35.5 per cent. of the whole series, or 47.8 per cent. of those in whom pregnancy was possible, since sixteen patients either had non-patent Fallopian tubes or their husbands were known to be sterile. Of the twenty-two who became pregnant seven miscarried, but two are now pregnant again. (Cases where menstruation was delayed for four weeks or more in a patient previously regular and followed by a heavy loss were accepted as miscarriages.) Of the remaining twenty-four cases in which the patient has not yet become pregnant (and in the later ones of the series only three months have elapsed) we can assume that in 21 per cent. the husband is sterile, judging from the result of the thirty-nine whose semen was examined; but we are still left with a number of women in whom so far pregnancy has not occurred though all investigations on both sides have been favourable.

Discussion

Lipiodol *per se* has a therapeutic value in that it opens up the Fallopian tubes. It would seem to be of greater use than air insufflation, because the latter ceases to distend as soon as the pressure is reduced. Lipiodol, being a heavy oil, does not immediately drain back through the uterus if the tubes are occluded, but continues to exert a small degree of hydrostatic pressure that may be sufficient to open up an otherwise non-patent tube; for a uterosalpingograph taken a minute after injection may reveal lipiodol to the end of the tube only, whereas a picture taken twenty-four hours later shows traces in the peritoneal cavity. The results obtained, however, cannot be attributed entirely to the action of lipiodol, for Bécclère and François (1937) quote only 18 per cent. of pregnancies out of a series of sixty cases with unilateral or bilateral patency, and Volk (1936) 25 per cent. out of a series of 110 cases. These latter figures do not differ to any extent

from the results obtained by air insufflation, for 24.57 per cent. of the 1,620 cases quoted by Rubin (1932) in which some degree of patency was observed became gravid, as did 18.42 per cent. of thirty-eight quoted by Bonnet (1936).

In my series forty patients injected with lipiodol received no hormonal therapy, and thirteen became pregnant (Table III). There was tubal occlusion or sterility of the

TABLE III.—Record of Women Pregnant Following Lipiodol Injection Alone

Case No.	Name	Age	Number of Children or Miscarriages	Years Married or since Last Child	Years Contraceptives used	Report on Husband's Semen	Patency of Tubes	Months elapsed before Pregnant	Miscarried (Number of Months Pregnant)	Remarks
1	K. E.	26	1	3	0		+	Stat.		
2	L. H.	25	1*, 1 mis.*	3	0		+	Stat.		*By previous husband
3	M. H.	31	0	7½	0	+	+L.	Stat.		
4	I. H.	26	2 mis.	4	0		+	7		
5	V. W.	22	0	5	0	+	+	3		
6	J. G.	30	0	3	2	+	+L.	Stat.		Right hydrosalpinx
7	B. C.	26	0	2½	0	+	+	Stat.	3	
8	E. H.	28	0	5½	0	+	+	1	2	Low sugar tolerance found
9	M. P.	27	0	4	1	+	+	1		
10	M. K.	28	0	4½	3½	+	+	2		
11	M. M.	25	0	2	0	+	+	4	1½	Now pregnant again. Taking thyroid extract and pituitary whole gland
12	B. G.	31	0	6	0	+	+	Stat.	1½	Now pregnant again. Obese; dieted, taking thyroid extract
13	E. B.	35	0	4	0	+	+	2		

husband in fourteen of these forty cases, so that 50 per cent. success can be attributed to lipiodol injection alone. It is known that the failure of many patients to conceive is due to failure of ovulation, and it is claimed that ovulation is more likely to occur after suitable administration of hormones. Dihydroxyoestrin benzoate, in the form of either progynon-B oleosum or dimenformon, given in two weekly doses of 20,000 I.B.U. in the two weeks directly following the menstrual period, or as a larger dose of 50,000 I.B.U. about the twelfth day of the menstrual cycle, appears to have had some value. In cases where the uterus is undersized, nidation of the fertilized ovum is more likely to occur if further development is brought about by more regular and prolonged hormonal therapy. Failure of the fertilized ovum to embed is one of the causes of sterility, the fault either lying in the ovum itself, owing to some intrinsic deficiency (maternal or paternal), or being due to failure of the uterus to undergo the modifications necessary for embedding of the ovum, because of insufficiency of corpus luteum hormone. Further investigation on the pregnanediol (excretion product derived from the corpus luteum) content in the urine at the time of ovulation in sterile patients is being undertaken.

Insufficiency of this latter hormone, again, may be dependent on an insufficiency or an unbalanced output of anterior pituitary hormones. As Davis and Koff (1938) have recently indicated, there are probably two factors in the anterior pituitary gland responsible (*a*) for follicle ripening and (*b*) for luteinizing; and sterility may depend on an incorrect balance between these two. It is hoped that in time one may be able to isolate the hormones from

the gonadotropic substance present in the urine of pregnant women and inject them into sterile women. Thyroidal insufficiency (in both parties) may account for sterility, since Lawrence and Rowe (1928) have shown that 39 per cent. of a series of patients with some degree of thyroid insufficiency were sterile. The results of this series show that investigation and treatment of as many as possible of the existing causes of sterility, at the same time, raise the fertility level and render pregnancy more likely to occur.

Summary

1. Sixty-two women complaining of sterility have been examined, the patency or non-patency of the Fallopian tubes being confirmed by uterosalpingography, and hormone therapy has been administered in many cases.

2. Of the husbands examined 13 per cent. were absolutely sterile, and in 8 per cent. few and distorted spermatozoa only were observed.

3. In 17.7 per cent. of women examined no lipiodol entered the peritoneum, showing non-patency of the tubes, and in 19 per cent. of the remaining cases one tube only was patent.

4. Of the women who had been sterile from two to eleven years (with the exception of two cases of nine months and one year respectively) 47.8 per cent. have to date (September, 1938) become pregnant within a period varying from one to seven months after treatment was begun. (In all these cases the tubes were patent and the husbands fertile.) Of the whole series 35.5 per cent. became pregnant.

5. The average age of the patients in this series is 30 years.

6. The average number of years of sterility is 5.7.

I am greatly indebted to Miss Dearnley, who has allowed me to carry on these investigations in her out-patient department at the Royal Free Hospital. I am also indebted to Dr. Ulysses Williams for the interesting series of radiographs, to Dr. Pillman Williams for so kindly carrying out basal metabolism and sugar tolerance investigations, and to Dr. Rawlinson, director of the Institute of Public Health and Hygiene, for his careful examination and report on the semen specimens.

REFERENCES

- Béclère, C., and François, E. (1937). *Bull. Soc. Obstét. Gynec. Paris*, **26**, 568.
 Bonnet, L. (1936). *Bull. Mém. Soc. Chirurgiens Paris*, **28**, 275.
 Clauberg, C. (1935). *Dtsch. med. Wschr.*, **61**, 1189.
 Coventry, W. A. (1934). *Amer. J. Obstet. Gynec.*, **27**, 912.
 Davis, M. E., and Koff, A. K. (1938). *Ibid.*, **36**, 183.
 Kilroe, J. C., and Hellman, A. M. (1933). *Ibid.*, **25**, 152.
 Lawrence, C. H., and Rowe, A. W. (1928). *Endocrinology*, July-August, 410.
 Pujol y Brull, A., Vanrell, J., and Riera, V. C. (1929). *J. Radiol. Electrol.*, **13**, 38.
 Rubin, I. C. (1932). *Amer. J. Obstet. Gynec.*, **24**, 561.
 Volk, W. (1936). *Z. Geburtsh. Gynäk.*, **113**, 339.
 Williams, L. C. Pillman (1933). *Lancet*, **2**, 858.

I. Snapper and P. Formigne (*Nederl. Tijdschr. Geneesk.*, 1938, **82**, 1937) review the literature and record three cases in adults of the coexistence of aneurysm of the cerebral arteries and congenital cystic kidney. The first case was one of a polycystic kidney associated with a calcified renal cyst, the second showed a solitary renal cyst, and the third a polycystic kidney with multiple small cysts in the liver. The first patient recovered and the other two died. In the patient who recovered neurological examination suggested the presence of a subarachnoid haemorrhage, and radiographs showed the polycystic kidney with a calcified cyst. In the second case necropsy showed a ruptured aneurysm of the right internal carotid at its junction with the middle cerebral artery, and in the third case a ruptured aneurysm of the middle cerebral artery with subarachnoid haemorrhage was found.

AURICULAR FIBRILLATION AND THE THYROID

BY

E. G. SLESINGER, O.B.E., M.S., F.R.C.S.

Surgeon, Guy's Hospital

Thyrotoxicosis is the fourth commonest cause of auricular fibrillation, rheumatic heart disease occupying first place and myocardial degeneration and hyperpiesia the second and third. In the thyrotoxic cases, however, it is almost invariably possible to restore the patient to normal rhythm and to prevent relapse, whereas in the other types the prognosis is less favourable. The correct diagnosis and treatment of thyrotoxic fibrillation are therefore of great importance. Without adequate treatment invalidism, gross cardiac failure, and death will follow, all of which are preventable by prompt and proper treatment. In this connexion it is important to remember that thyrotoxicosis is not always as easy to diagnose as might at first sight appear. Clinically the cases can be conveniently divided into two types, which are termed primary and secondary.

In the primary type the goitre and the toxic symptoms appear at or about the same time. The whole epithelium of a previously normal thyroid is able to respond to the stimulus producing the thyrotoxicosis, and the resultant symptom-complex, usually termed exophthalmic goitre, is typical and usually readily recognized. The patients are in the main young adults and their hearts can stand the strain of continued tachycardia for a long time without developing arrhythmia, so that auricular fibrillation, while it occurs, only does so late in the disease. In secondary thyrotoxicosis, on the other hand, a goitre, usually of the nodular colloid type, has been present for a number of years before the onset of toxic symptoms. Only part of the epithelium of the gland can respond to the exciting stimulus, and the resultant symptoms are less marked and in part different. The patient, however, is older than in the primary case, and the heart is less able to withstand a prolonged tachycardia. It is in this secondary group that the greater number of cases of auricular fibrillation are met with whose thyrotoxic origin has not been diagnosed.

Diagnosis

If the patient is recognized as suffering from thyrotoxicosis and is noted to have arrhythmia it is nearly certain to be auricular fibrillation. In a rapidly beating heart extrasystoles are sometimes indistinguishable without an electrocardiogram, and such occasional extrasystoles are commonly found in thyrotoxicosis. On the other hand it is the case with very frequent extrasystoles that is most liable to be mistaken for fibrillation, and, as Campbell (1935) has pointed out, this is not a common condition in thyrotoxicosis. If, therefore, there is an established arrhythmia in a patient known to be thyrotoxic it can safely be assumed to be fibrillation, although this should, of course, be confirmed by an electrocardiogram. Paroxysmal auricular fibrillation, again, is not uncommon in hyperthyroid cases, and, although it may exist for some time, generally progresses to a state of established fibrillation if untreated.

Careful inquiry must be made into the history of attacks of arrhythmia. These are generally recognized by the patient. Both the frequency and the duration of attacks vary in different patients, but are usually fairly constant in the individual, especially as to duration; the patient being aware of when the attack starts and of