EXTRACTS FROM
THREE LECTURES
ON
TUBERCULAR DISEASES OF BONES AND JOINTS.

Delivered at the Royal College of Surgeons of England.

By W. WATSON CHEYNE, M.B., F.R.C.S.,

Hunterian Professor; Surgeon to King's College Hospital; and to the Paddington Green Children's Hospital, etc.

LECTURE II.—TUBERCULAR CHANGES IN BONE.

THE changes which occur in bone as the result of tubercular disease are very various, and to some extent merge into one another. The following are, however, the chief forms:—(1) Miliary tuberculosis of bone; (2) soft caseating deposits in bone; (3) tubercular deposits with sclerosis of bone and necrosis; (4) superficial tubercular disease of bone (caries) in connection with which we have to study the changes in the articular cartilage; (5) the condition termed "caries sicca"; (6) diffuse condensation of bone in connection with tubercular disease; (7) diffuse softening of bone and the formation of so-called "red marrow"; and (8) tubercular periostitis and tubercular osteomyelitis. In connection with these tubercular affections of bones we must also bear in the mind inflammatory processes which accompany them, and which play a most important part in the destructive changes.

1. *Miliary Tuberculosis of Bone.*—Miliary tuberculosis seldom occurs in bone except in cases of acute general tuberculosis. It is true that we not uncommonly find miliary tubercles in bone in the vicinity of a large tubercular deposit, but I here refer to diffuse miliary tuberculosis without any large deposit. Where miliary tuberculosis occurs in bone in the course of an acute general tuberculosis, the appearance of the tubercles and the changes associated with them are the same as in other organs, but, as a rule, the number of tubercles is not so great, and they are not uncommonly most numerous towards the extremities of the bone. For a long time the occurrence of miliary tuberculosis in bone was entirely denied by some authors, but instances of this kind have been published from time to time, and Lazarus has recorded five cases of acute general tuberculosis in which the bones were examined with positive results. In his cases, however, the tubercles were not present in all bones nor in all parts of the bone. This form, being only part of a fatal general disease, is, however, of very little practical importance.

Apart, however, from acute general tuberculosis or large deposits in bone, miliary tuberculosis of individual bones has been described, but I have not yet come across a case, and I have cut up several museum specimens which were said and appeared to show

it, and where there has been anything abnormal at all it has only been patches of simple osteitis.

It is also not uncommon to find tubercles, or what look to the naked eye like tubercles, in the sternum, ribs, and even the vertebrae, in *post-mortem* examinations of cases of phthisis, and that although there is no general tuberculosis. The only specimen of this kind which I have obtained was, however, spoilt in softening, and I cannot therefore say anything about the condition.

2. *Soft Caseating Tubercular Deposits in Bone.*—Soft and generally caseous tubercular deposits in bone are by no means uncommon as the primary condition in tubercular joint disease. These deposits vary much in size and situation, but they are as a rule small, and situated in the ends of the bones, usually close to the articular cartilage or the surface of the bone. They also vary in number in the same case, but in my experience they are most usually single, though in some cases there may be deposits in more than one of the bones entering into the formation of the joint or in more than one part of the same bone.

In this specimen (Fig. 5) we have an excellent example of one of these soft caseating deposits in bone. This is a section of the external condyle of the left femur of a child, aged 5. Two years before admission an abscess formed on the outer side of the left knee-joint and burst, leaving a sinus; the synovial membrane became much thickened, the knee flexed, and fresh abscesses formed. The condition of the joint as seen after operation was that the synovial membrane was much thickened, the articular cartilage of the patella was destroyed, the internal tuberosity of the tibia was much eroded, and there was a soft caseous deposit in the external condyle of the femur which communicated with the exterior, and to which the sinuses led.

On examining this deposit we see that it is more or less circular in shape, and entirely composed of soft tissue, the osseous trabeculae having completely disappeared. Towards the inner part of the deposit caseation is occurring; around this there is a condition of tubercular infiltration, and elsewhere a number of tubercles. Immediately around the deposit the trabeculae of the bone are seen to be

considerably thickened, and more numerous than at some distance from this spot. In the cancelli around the tubercular mass the normal fatty tissue has disappeared, and its place is taken by a swollen, somewhat fibrous, vascular material, not unlike that seen in gelatinous infiltration of the synovial membrane. Further away from the deposit the tissue in the cancellous spaces presents a fairly normal appearance. Towards the left hand side we see that the deposit has destroyed the surface of the bone, and made its way outwards leading to the abscess formation over it.

In this next specimen (Fig. 6) we have an example of a caseous deposit immediately beneath the articular cartilage of the tibia close to the spine in a child. I have no notes of the history of this case as it was not under my own care, but the knee-joint was excised, and the chief disease was in the external tuberosity of the

Fig. 5.

tibia and the external condyle of the femur, and had probably begun primarily in both of these situations. Here we have a similar appearance to that seen in the last specimen, namely, a central mass of caseous material without any bony spiculae in it, surrounded by soft granulation tissue containing tubercles. Surrounding the soft deposit we see a few sclerosed trabeculae as in the previous case, while further away the epiphysis is infiltrated with inflammatory cells, and the osseous trabeculae are thin and few in number; in fact, the whole epiphysis is in a state of rarefying osteitis. At one part it will be seen that the deposit has burst through the cartilage and opened into the joint. On each side of this point some fragments of cartilage are still present on the surface of the bone, but further away the articular surface is destitute of cartilage and carious. [Specimens were also shown of soft deposits in the neck of the femur, in the internal condyle of the femur, and in the head of the tibia.]

These specimens are sufficient to illustrate the state of matters in cases where soft deposits are present in the bone. These deposits generally undergo caseation in the centre, but in some instances this may not occur for a long time, and sometimes we find a mass of firmer fibrous tissue containing only a few tubercles scattered through it. Around the caseous part we find granulation tissue with tubercles. In these deposits the osseous trabeculae have usually been completely absorbed, though in some cases fragments of bone may still be seen. Immediately outside the tubercular growth it is not uncommon to find that the osseous trabeculae are thicker than normal, but this condition does not extend to any great distance beyond the deposit, and indeed is not always present. Generally also we find that a few of the cancellous spaces around the deposit, more especially those in relation with the thickened trabeculae, are almost entirely devoid of fat cells, and contain a peculiar swollen semifibrous material not unlike that found in the thickened synovial membrane in the neighbourhood of tubercular growth. Further away from these soft deposits we usually find a condition of rarefying osteitis. It is noteworthy, however, that here, as in caries, the rarefying osteitis is most marked at some distance from the deposit, the intervening tissue being less affected; and this rarefying osteitis not infrequently occurs in patches. Not uncommonly we find new osseous formation from the periosteum outside these deposits, but this is perhaps not so common as in the next form of tubercular bone disease.

The process probably progresses somewhat in this way. The tubercular virus is deposited at some part of the bone, and tubercles form, the tubercles causing irritation and the formation of a young granulation tissue around them, which attacks the trabeculae of the bone and leads to their absorption. The condition is illustrated in the accompanying photograph (Fig. 7), where we see the formation of tubercular tissue in neighbouring cancelli, and the lacunar absorption of the bone by the granulation tissue

around the tubercular mass. The tubercles multiply, this granulation tissue extends, and absorption of the trabeculae progresses, while caseation occurs at the oldest part of the growth. Further away from the tubercular mass the irritation is less, and a small amount of young swollen fibrous tissue is formed accompanied by sclerosis of the trabeculae. This fibrous material is again invaded by the tubercular growth and the granulation tissue, and the thickened trabeculae are again absorbed, and so the process goes on extending, aided no doubt by the rarefying osteitis which is occurring in the neighbourhood, till ultimately the surface of the

bone is reached. In some cases the tubercular growth is more rapid, and time is not afforded for total destruction of the trabeculae before caseation is complete, and hence we find in these cases portions of osseous trabeculae in the midst of the caseous material and the sclerosis of the surrounding bone may be absent.

The further history of the case is as follows: While the tubercular deposit is still enclosed in the interior of the bone, there may be only slight clinical evidence of its presence; but once it reaches the surface, various phenomena occur, differing according to the point where the deposit has opened. If the deposit opens through the articular cartilage and communicates directly with the joint, as has occurred in most of the specimens which I have shown, the infective material is distributed over the whole surface of the synovial membrane almost at the same time, and the first type of synovial disease rapidly develops, soon followed in most cases by the occurrence of chronic suppuration in the joint. In another set of cases the deposit reaches the surface at the point where the synovial membrane is reflected off the bone, and it is thus shut off, for a time at least, from the joint cavity. In this case thickening of the synovial membrane, usually of the third type, occurs, being most marked, especially at first, over the osseous deposit, but gradually extending and involving the whole of the synovial membrane.

In a third set of cases the deposit reaches the surface of the bone, quite outside the synovial membrane, and then a chronic abscess usually forms over it, in the manner which I shall describe in the next lecture, and the joint itself may never become affected. The specimen on the screen (Fig. 8) shows the sequence of events, when the osseous deposit bursts into the joint cavity. This is a complete

longitudinal section through the lower end of the radius, the carpus, and the end of the second metacarpal bone. We see at the posterior and carpal end of the metacarpal bone (towards the lower part of the figure) a caseous deposit which has opened into the carpo-metacarpal articulation. In this articulation the cartilage is almost entirely destroyed and the synovial membrane is thickened and villous; the disease has spread to the articulation between the two carpal bones, which is filled up with fibrous tissue containing tubercular tissue, and here also the cartilage has been almost entirely destroyed. The radio-

Fig. 6.

Fig. 7.

carpal articulation is not nearly so much affected, but thickening of the synovial membrane is occurring, and this thickened tissue is creeping over the surface of the cartilage of the radius in the manner which I shall describe when I come to speak of the mode of destruction of cartilage. We also see sections of sinuses in the soft parts with caseating walls, and leading to the joints.

3. *Tubercular Deposits with Sclerosis of Bone and Necrosis.*—In marked contrast to the foregoing processes we have those where there is sclerosis of bone and formation of sequestra. This process is very common in tubercular diseases of bone, varying much, however, in extent and result in different cases. The following specimens will illustrate the chief points.

The first specimen (Fig. 9) is a section of the patella from a case of disease of the knee-joint in a female, aged 16. There was much swelling of the knee and an abscess on the outer side of the patella; the disease had commenced spontaneously. This is an excellent example of this form of tubercular disease, and shows all the stages in the formation and separation of tubercular sequestra. With regard to this photograph I must, however, in the first place, point out that for some reason or other the thickened trabeculae in the deposit have taken the stain (which was a triple stain) somewhat differently from the trabeculae in the healthy part, and have not blocked the light to such an extent; and the consequence is that while in the healthy part of the bone the osseous trabeculae appear dark, in the photograph and the cancelli light, in the diseased part the condition is reversed, and the trabeculae are light and the caseous contents of the cancelli dark. Bearing this in mind, we see very distinctly the marked thickening of the trabeculae in the deposit, and we also see that the majority of the cancelli contain caseous material. Around the deposit, rarefying osteitis is occurring, and the trabeculae are becoming destroyed, though only in places; the tissue at these points contains numerous tubercles. In other parts at the margin the cancelli contain vascular fibrous tissue, and the trabeculae are becoming thickened. Nearer the centre tubercles appear in these cancelli, and the thickened trabeculae are again becoming eroded, and at the centre the contents of the cancellous spaces are caseous, and the erosion of the trabeculae has come to a standstill. I shall now show more highly magnified photographs from the line of separation of the sequester, from the outer part where sclerosis is going on, and from the central part where caseation has occurred.

If we examine the line of separation (Fig. 10) we see that at parts, as in this photograph, the trabeculae have completely disappeared, and the interval is filled up with granulation tissue containing tubercles; at other parts the trabeculae in the deposit are still continuous with the trabeculae outside.

In the photograph from the outer part of the sequester there were evidences of active sclerosis, the trabeculae being greatly thickened and presenting a laminated appearance, while the contents of the cancelli are young fibrous tissue. In the photograph from the centre of the sequester the trabeculae were somewhat thinner and a good deal broken up, and the contents of the cancelli were cheesy.

[A complete vertical section of the os calcis was also shown, in which the greater part of the bone—the posterior—formed a sequester composed of thickened trabeculae, with cheesy material in the cancelli. Between the anterior part and the sequester there was a layer of vascular, fibrous, and granulation tissue, with caseating tubercles; at that point separation of the posterior part of the bone as a sequester was occurring, and it showed very well one great characteristic of tubercular sequestra, namely, that for a long time the separation is incomplete, for at various parts trabeculae could be traced between the necrotic portion and the anterior part.]

These sequestra vary in density very greatly, being in some cases only slightly denser, or even less dense, than normal cancellous bone, while in other cases the trabeculae are extremely thickened, and the mass resembles rather the outer shell of a bone than cancellous tissue.

In the section of the patella, which I have just shown, we have an example of the intermediate, or what one may term the normal, density and character of one of these sequestra. [A specimen of a less dense form was shown from the lower end of the femur; one of extreme density from the acetabulum, where the cancellous spaces were very small, and in some parts almost absent; and one in the upper end of the humerus showing a sequester of varying density. In the last specimen the margin of the sequester was composed, in parts, of sclerosed bone, in the cancelli of which was vascular fibrous tissue, while in other parts of the circumference the cancelli were larger, and contained caseous material. The centre of the sequester was much less dense, and did not contain caseating tubercular tissue.]

These tubercular sequestra are usually larger than the soft tubercular deposits, but vary much in size and shape. They are generally quadrilateral, or irregular, in form, but they are sometimes wedge-shaped, the base of the wedge being directed towards the cavity of the joint. In the freshly cut bone they present a dense yellow appearance, and they are usually incompletely detached. In some cases the articular cartilage over them is destroyed, and they project into the joint cavity, and if the joint has still been used the surface of the sequester may be eburnated. They vary in density from that of the hardest bone to a friable consistency, but most commonly they are denser than the normal cancellous bone. In most joints they usually occur close to the articular ends of the bones, and frequently immediately under the cartilage; in other cases, more especially in the upper end of the femur, they are found just beyond the epiphyseal cartilage, or in the substance of the neck of the bone.

When we study these sequestra under the microscope, we find that they are usually composed of much thickened osseous trabeculae, and also of trabeculae of new formation. The density of the newly-formed bone varies in different cases, and this variation apparently depends on the rapidity and extent of the process. The meshes of the caseous network in these sequestra are filled with fibrous or granulation tissue, frequently containing tubercles, or more usually with caseous material. Where the sequester is very dense, the amount of soft material in the meshes of the bone is naturally very small. Frequently, however, even in very dense sequestra, many of the trabeculae have been a good deal broken up in places by the tubercular growth before caseation is complete, and thus a crumbling condition may be produced; and though many of the trabeculae are much thickened, parts of the mass may break down readily under the finger. Surrounding the sequester, we find a layer of soft, and often fibrous, tissue, generally containing tubercles or tubercular infiltration, and when the sequester is quite loose osseous trabeculae are, of course, completely absent from this limiting wall of soft tissue.

The process of sequester formation in tubercular disease is probably shortly the following: As the result of the deposit of tubercles in a certain part of the bone there is inflammatory reaction in the neighbourhood, and ultimately the formation of young vascular fibrous tissue in a considerable number of the cancelli of the bone. In these cancelli thickening and new formation of osseous trabeculae goes on, and this process continues to extend at the edge. While this fibrous formation and sclerosis of the bone is going on, the tubercular growth is also extending, and accompanying it there is reabsorption of the newly formed bone. Caseation of this tubercular growth occurs, and then the

Fig. 10.

Fig. 11.

Fig. 12.

Fig 8.

Fig. 9.

Fig. 13.

absorption of the bone ceases, and thus we come to have a central patch of sclerosed bone, with caseous material in the cancelli, surrounded by very vascular fibrous tissue and denser bone. By-and-by the process ceases to extend with the same rapidity, and then absorption of the connecting trabeculae of bone, and ultimately complete detachment of the sequestrum, may take place. The thickness of the osseous trabeculae in the necrosed fragment depends on the rapidity with which caseation of the tubercular deposit has occurred, and also on the course taken by that deposit. If caseation has only occurred slowly there may be considerable thinning and breaking up of the previously sclerosed trabeculae; while if it has occurred rapidly the thickness of the new bone is not much diminished. The tubercular growth may not, however, invade the whole of the sclerosed bone, and where it does not extend the sclerosis goes on and very dense bone is formed, this bone being ultimately cut off from the surrounding parts, and composing part, or even the whole of the sequestrum. In some cases the tubercular growth follows a circular or triangular course around bone which has been invaded by it, bone which has become much sclerosed but not invaded, and bone which has not been much affected, and thus the sequestrum may present the appearances shown in the last specimen. We thus see that in the formation of these sequestra we have two processes going hand in hand, namely, growth of vascular fibrous tissue with sclerosis of bone, and growth of tubercular tissue with rarefaction of bone; and the character of the sequestrum varies according to which of these processes is in excess, according to the rapidity with which the tubercular tissue extends or caseates, and according to the course taken by the tubercular growth. Formation of vascular fibrous tissue with sclerosis of bone is not uncommon in tubercular diseases of bones, but this sclerosis may go on to a great extent without ending in death. What determines the formation of a sequestrum is the invasion or encircling of this sclerosed tissue by the tubercular growth.

Various views have been put forward at different times as to the nature of these sequestra. Some authors have supposed that they were composed mainly or entirely of fibrous tissue, while others have held that their density is due to the deposit of calcareous salts. Without doubt the caseous material in the cancelli in these sequestra is very apt to become infiltrated with calcareous salts, and when this is the case the weight and apparent density of the sequestra are much increased; but, as I have shown, this is not at all the essence of the process. The view, however, which has been most generally accepted is that advocated by König, namely, that these sequestra arise as the result of embolism, and that, the vessel being blocked by tubercular material, the portion of bone supplied by it dies. König lays especial stress on the wedge shape and position of these sequestra as evidence of the accuracy of his opinion; and while he has not failed to observe that the osseous trabeculae in the sequestra are thicker than in the surrounding bone, he concludes that this is only an apparent sclerosis due to rarefaction of the surrounding bone. The most superficial glance at complete microscopic sections of the affected parts—examples of which I have shown on the screen—will, however, suffice to show the erroneous nature of these views. There can be no question that the portion of bone which has died has previously been the seat of active growth—a fact quite irreconcilable with the embolic theory—and we see similar evidences of active growth in the bone around the deposit. This being the case, it is quite clear that whatever influence the distribution of the vessels may exert on the direction of spread of the tubercular tissue, and consequently on the shape of the sequestrum, the formation of the sequestrum is not the result of embolism.

I need not repeat what I have previously said in connection with soft deposits as to the further progress of the disease when the tubercular process reaches the surface of the bone. The progress of events is practically exactly the same in both cases.

[Several specimens were then shown to illustrate what frequently happens when a case of tuberculosis of bone has become "cured." One specimen was a section of the lower end of the humerus from a case of disease of the elbow-joint, where the disease had been in abeyance for ten years. There was very marked sclerosis of the bone, and remains of tubercles in many of the cancelli; these remains consisted of amorphous material, in some parts traces of cells and giant cells and fragments of bone. In other cases the deposit may entirely disappear, and dense fibrous tissue be found in its place. In others, again, the tubercular de-

posit, especially when caseous, may become infiltrated with calcareous salts and encapsuled.]

4. *Superficial Caries of Bone and the Changes in the Articular Cartilage.*—The most common tubercular affection of bone is tubercular destruction or caries of the surface of the bone. The term caries was originally applied to all destructive or ulcerative changes of the surface of bone, but of late it has been restricted to those changes of the superficial layers of bone which occur in connection with tubercular disease, and restricted in this way, the term is a very convenient one.

Caries of the articular ends of bones is never, so far as I have seen, the primary tubercular affection; it practically always occurs secondarily either to a deposit in bone or to synovial disease. It is also a curious fact that, even in the case of osseous deposits, the growth of tubercles in the superficial cancelli of the bone in the neighbourhood of the deposit, and the destruction of the cartilage, only goes on to a comparatively slight extent, and in some cases not at all, till the synovial membrane has become affected or the deposit has opened into the cavity of the joint, after which the disease progresses with great rapidity. One might have expected that in the case of a deposit immediately beneath the cartilage, as in several of the specimens shown, the tubercular growth would have spread along under the cartilage, and caused its exfoliation from within. This, however, is not the case, at any rate not to any marked extent; the tubercular growth seems to be more or less confined within the deposit in the first instance, and not till it has formed a communication with the joint or reached the soft tissues does it seem to be liberated, and free to spread over the whole part. As I have previously pointed out, where a tubercular osseous deposit reaches the articular surface, erosion of the cartilage occurs over it, a communication is formed with the joint, the synovial membrane becomes infected, and then the cartilage is attacked, and it is not, as a rule, till destruction of the cartilage has progressed to a considerable extent that the surface of the bone is affected. We must, therefore, study the changes which occur in connection with the cartilage in the first instance.

The changes in cartilage as seen by the naked eye are well known, and consist chiefly in thinning and complete disappearance of portions of the cartilage; its detachment in the form of thin shreds often full of holes, or even of thicker masses; indeed, in some cases cartilage which shows very little alteration may be partially loose and movable on the subjacent bone. These changes generally commence and are most marked at points where the synovial membrane joins the cartilage, as at the edges of the cartilage, at the points of attachment of the crucial ligaments in the knee-joint, in the neighbourhood of the ligamentum teres, in the hip, etc. Its destruction may also begin in the neighbourhood of the openings of tubercular osseous deposits, and the thinning of the cartilage is sometimes well marked at points of pressure. The following specimens will illustrate the nature of these changes, and the mode in which they occur.

As I have said, the early changes in the cartilage are, as a rule, most marked at the points where the synovial membrane is reflected on the cartilage, and we not infrequently see that the cartilage is but slightly, or not at all, affected, except at the margins where soft tissue continuous with the synovial membrane is spreading over and destroying it from the surface. This was well seen in the section through the wrist-joint (Fig. 8) where the synovial membrane was spreading over the cartilage of the radius and destroying it first at the margins. [Other specimens were shown to illustrate this.] The first change which occurs is apparently the spread of fibrous vascular tissue over the surface of the cartilage, this tissue becoming thinner and thinner as we pass away from the edge. This is seen in this specimen (Fig. 11), which is from the margin of the articular cartilage of the hip-joint, showing commencing destruction of the cartilage from the surface. [Several specimens were then shown to illustrate the minute changes in the cartilage and the subsequent invasion of the new tissue with the tubercular growth.] The process then which takes place at the margin of the cartilage is the extension of the new tissue in a thin layer over it, followed by changes in the cartilage itself, which lead to its conversion into fibro-cartilage and ultimately into fibrous tissue. The new tissue so formed becomes subsequently infiltrated with the tubercular growth and undergoes caseation.

If now we study the changes which occur in the cartilage at a greater distance from the edge of the synovial membrane, we find that this extension of the soft tissue goes on for a considerable

distance, but, as a rule, in quite a thin layer, while here and there greater growth has occurred and the previously-mentioned changes in the cartilage are taking place, and thus we come to have islets of soft material filling depressions on the surface of the cartilage, separated by patches of but slightly altered cartilage, and connected by a thin layer of soft tissue on the surface of the cartilage. In this way the pitted or completely perforated appearance of the shreds of cartilage is produced. This is well seen in this specimen (Fig. 12), where a number of these depressions are present. [Other specimens were shown, illustrating in a more marked degree the result of the erosion of the surface of the cartilage, and also showing the presence of tubercular tissue in some of these depressions.]

The chief destruction of the cartilage occurs from the surface in the manner just described, but we not infrequently observe somewhat similar changes in the deeper part, especially at the margins. In some cases also osteitis, without formation of tubercle, occurs in the superficial cancelli of the bone over a considerable area, and a slight amount of soft tissue is formed between the bone and the cartilage, and this is what has usually occurred in those cases where the apparently unaltered cartilage can be moved over the bone beneath.

As a rule, when the deeper part of the cartilage becomes affected one of two things happens, namely, either from the margin of the cartilage, or from the bottom of one of these excavations on the surface the newly-formed tissue spreads in between the cartilage and the bone, destroying the deeper part of the cartilage, detaching it, and leading to exfoliation of pieces of as yet unaltered cartilage; or tubercles form in the superficial cancelli, destroy the superficial layer of the bone, and lead to erosion of the deeper part of the cartilage. This erosion of the cartilage from the deeper surface is, however, a comparatively rare occurrence, except at the margins where the process has begun at the surface, as previously described. [Specimens were shown illustrating these points.]

Another mode of destruction of cartilage is seen in this specimen (Fig. 13) which is a vertical section of the internal condyle of the femur from a case of disease of the knee-joint of six months' duration, in which the disease was primarily synovial, and in which there was great thickening of the synovial membrane. The cartilage was intact except at the margins and at one spot towards the centre and anterior surface of the internal condyle of the femur, where there was a small depression on the surface. This photograph is taken from a vertical section of the end of the bone through this spot. We see that the depression in the cartilage is due to destruction of the cartilage at this point, and before the celloidin was dissolved out one or two small detached fragments of cartilage were seen at the surface which have been swept away in clearing the specimen. We further see a number of flask-shaped spaces in the cartilage, which under a higher power are seen to be filled with young fibrous tissue. Some of these spaces communicate with the surface of the cartilage either freely or by narrow channels, while the majority, and perhaps all, are connected with the superficial cancelli of the bone by similar channels. At various other points along the deeper part of the cartilage we see similar flasklike projections communicating with the bone, and where these occur the most superficial cancelli show osteitis. I have not been able to find any tubercles in the bone.

Before leaving the subject of the changes in cartilage, I may refer to the important question whether cartilage can be the primary seat of the disease. As regards hyaline or articular cartilage, this may certainly be answered in the negative, but it is still an open question with regard to fibro-cartilage. In the case of the spine it was long thought, and is I believe still held by some, that the changes frequently commence in the intervertebral cartilages, but this is clearly not the case. In the knee-joint, however, Kocher has described, under the term "meniscitis fungosa," what he believes to be primary tubercular disease of the semilunar cartilages, but there is no evidence that the disease did not begin in the synovial membrane in the neighbourhood of the cartilage and spread on to it.

BALNEOLOGICAL CONGRESS.—The thirteenth Congress of Balneologists will be held at Berlin in March, 1891, under the presidency of Professor Liebreich. Notice of communications, etc., should be sent to the General Secretary, Dr. Brock, 42, Schmidtstrasse, Berlin.

THE BENIGN NEOPLASMS OF THE RECTUM.

Read in the Section of Pathology at the Annual Meeting of the British Medical Association, held in Birmingham, July, 1890.

By CHARLES B. BALL, M.Ch., F.R.C.S.I.,

Surgeon Sir Patrick Dun's Hospital, and University Examiner in Surgery.

MORBID growths are met with in the rectum and at the anus with great frequency, and, as compared with other portions of the intestinal tract, the fact that this part of the gut can alone be examined during life with facility renders the proportion met with in this region apparently large, yet it cannot be denied that the termination of the bowel is very much more frequently affected than the other portions by these growths. This is more particularly noticeable in the case of piles, or hæmorrhoids, which constitute by far the commonest tumours of the rectum and anus, but are unknown in the other portions of the intestinal tract. Owing to the more potent sources of irritation to which the lower bowel is subject, the greater liability to tumour formation can readily be accounted for, and in the case of hæmorrhoidal tumours there is a distinct anatomical reason for their existence here. During the passage of a solid mass of faeces along the great intestine, the contraction of the muscular tissue causes a considerable pressure to be exercised on the blood vessels. In the upper portions of the intestinal tube this is rather salutary than otherwise, as the vessels here are for the most part arranged at right angles to the axis of the bowel, so that the contraction of the tube empties the veins into the larger channels, and thus materially aids in the circulation of blood; but when the termination of the rectum is approached the conditions are materially altered, because the blood from the lower portion of the bowel passes up by the superior hæmorrhoidal veins, the tributaries of which originate at the junction of skin and mucous membrane. These vessels, instead of being arranged at right angles to the axis of the intestine as is the case higher up, are parallel to the axis, lying immediately under the mucous membrane, so that as the faecal mass is pressed onward by the contracting muscular wall, there is a tendency to force back the column of blood in the veins towards the tributaries, which are therefore liable to become dilated and varicose; indeed, a slight amount of varicosity of these veins is held by some authors to be normal. This liability is further augmented by the fact that the radicles of the superior hæmorrhoidal veins have but trivial anastomoses with the inferior and middle hæmorrhoidal veins, so that but little relief can be afforded to the distended veins by the passage of blood *via* the inferior and middle hæmorrhoidal veins into the general or caval circulation.

Various terms have been used to designate the forms of hæmorrhoids, external and internal, the former of which are subdivided into venous, cutaneous, and compound, while in the latter the venous, columnar, and nœvoid varieties are recognised. Now, although these varieties possess points which render them clinically important, it cannot be maintained that they are distinguished by marked pathological characteristics. When taken broadly, the pathology of all is practically identical, namely, increase and dilatation of the blood vessels, more particularly the veins with proliferation of the connective tissue. In the case of external piles the covering is the scaly epithelium of the anal canal, while internal piles are covered with mucous membrane.

I have elsewhere endeavoured to show¹ that extravasation of blood is not an important factor in the pathology of piles. In the simple venous external pile which appears apparently suddenly, what really happens is this: A varicose condition of one or more of the inferior hæmorrhoidal veins has previously existed without attracting the attention of the patient; it is only when thrombosis and perivascular inflammation supervene that the tumour is noticed, and the patient has what is known as "an attack of piles." That this is indeed an intravascular, and not extravascular, coagulation of blood can easily be demonstrated; if an incision be made into a simple inflamed external pile a blood clot can be turned out leaving a smoothly lined cavity in which an endothelial lining can be observed; moreover, as a result of stretching of the anus—now the prelude to almost all rectal opera-

¹ *The Rectum and Anus, their Diseases and Treatment.*