congratulate him upon his delay in publication until the patient's recovery has been well established.

ENTERECTOMY.

What would Astley Cooper or Abernethy, Lawrence, Syme, or Brodie, have said if a proposal had been made to cut away several inches of intestine, and, after uniting the divided edges of the upper and lower openings together, to return the intestines, and close the opening made in the abdominal wall. They all regarded a mere wound of intestine as almost certainly fatal, and thought life could only be saved by forming an artificial anus. How amazed any of these great surgeons, or any of their contemporaries, would have been if they had seen the specimen I now show you, which has been kindly lent me from the Museum of St. Thomas's Hospital. A boy, aged 14, was kicked by a horse one evening, and taken to St. Thomas's late at night. Next morning Mr. Croft, suspecting ruptured intestine and acute peritonitis, at once made an exploratory incision. He found faecal matter in the peritoneal cavity and a part of the ileum bruised and ruptured. The piece you now see was separated from its mesenteric attachment, and cut away. Makins's forceps, which are here, proved very useful in keeping the canal closed before the intestine was divided, and while the sutures were being passed. The mesenteric wound was closed by sutures, the cut ends of the intestine carefully adjusted, and fastened together by Lembert's sutures. About forty sutures were left in. The peritoneal cavity was carefully cleansed by warm boracic solution. No drain was used. The external wound was closed by silk sutures. Recovery was uninterrupted, and the boy was shown at the Clinical Society nearly a year afterwards in good health. The piece of intestine seen here inverted measures nearly three inches at the free border, but only about half an inch at the mesenteric border. The rupture is only about half an inch in diameter, but it had permitted free escape of faecal matter into the peritoneal cavity, and, if ever a life has been saved by a surgeon, this boy's was by Mr. Croft, and so will be many more if the practice he adopted is imitated.

Lembert's suture was employed in this and in other successful cases, but I cannot help thinking that Mr. Stanmore Bishop's is more perfect. He has described and illustrated it so fully in his paper published in the *Medico-Chirurgical Transactions*, that I need not now do more than point to the interesting preparations before you to show what accurate and complete union follows its application. The fine needles and silk Mr. Bishop uses are here, and his needle holder. I must, however, add a word of advice to those who may think of using them, that a good deal of practice in the study is necessary to enable anyone to use them with precision and at all rapidly.

My distinguished American friend, Dr. Senn, professor of surgery at Chicago, has made some valuable additions to the practice of enterectomy by teaching the use of absorbable plates of decalcified bone in facilitating the approximation and union of the two openings to restore the continuity of the intestinal canal. I have some of his plates here, preserved in an antiseptic solution as they are kept for use. They come away with the faeces about a week after the operation. They assist very much in securing complete apposition of the two serous surfaces, and Dr. Senn's method of slightly scarifying the two surfaces with the point of a needle doubtless adds to the rapidity and firmness of their adhesion. Similar scarification might be of use in the uterine suture, and I shall certainly give it a trial.

I can strongly advise all operating surgeons to study carefully Dr. Senn's important memoirs *On Resection of the Cecum for Carcinoma*, and on *The Diagnosis and Operative Treatment of Gun-shot Wounds of the Stomach and Intestines*. They abound in original suggestions and in records of experimental and operative work. But I cannot do more now than allude to two cases of intraperitoneal rupture of the bladder, which are brilliant examples of modern abdominal surgery.

Intraperitoneal rupture of the bladder has been almost always fatal, but Sir William MacCormac had two cases four years ago where he exposed the bladder by abdominal section, carefully closed the rent by silk sutures, washed out the peritoneal cavity by warm boracic solution, closed the external wound, and perfect recovery was the result. These are the first of such cases on record, and they are brilliant examples of modern abdominal surgery. Careful diagnosis and early skilful operation earned their just reward. Silk sutures were passed about a quarter of an inch apart through the serous and muscular coats only, twice on each side of the opening, so that when they were tied the mucous mem-

brane was inverted, and two serous surfaces were fastened together on the principle I have spoken of before. The peritoneal cavity was thoroughly cleansed by irrigation, a glass drainage tube introduced, and a catheter fixed in the urethra. These were removed after three days, and in three weeks the man was as well as before the accident.

In a second and similar case the same course was adopted, with the exception of drainage, and recovery was even more rapid than in the first case. Accurate closure of the wound in the bladder by sero-muscular sutures and thorough cleansing of the peritoneal cavity were doubtless the chief points in these two operations. As Sir William MacCormac urges in his very able report of these cases,² in cases of intraperitoneal ruptured bladder there is need for "earlier interference and bolder practice." The whole history of abdominal surgery points in this direction.

Here I am compelled by the hour to close this very hasty and imperfect sketch of what we, the surgeons of the present, have done, and are doing, and how we do it. And for our younger Fellows and Members—for the surgeons of the future—we may not be confident that with the energetic spirit of inquiry now awakened, with an enlightened determination to apply all the resources of modern scientific discovery to the perfecting of our art—with a conscientious aim at making it as truly conservative as is compatible with usefulness and progress—and with honourable feeling and highly cultivated judgment, directing hands delicately and expressly trained, we may augur for the surgeons of the coming time an influence supremely beneficent for mankind, and promise to its devotees the dignity and distinction justly earned by their life-giving and health-preserving work.

[The following corrigenda should be made in the portion of Sir Spencer Wells's paper published in the *BRITISH MEDICAL JOURNAL* of December 20th, page 1415, col. 1, paragraph 3, lines 3, 7, and 15: for "Bonilly" read "Bouilly." Same page and column, insert in paragraph 4, line 6, after "recovery," "of patients;" and in line 8, for "then" read "their."]

AN ADDRESS

OR

THE DIFFERENTIAL DIAGNOSIS OF PELVIC INFLAMMATIONS IN THE FEMALE.

Read before the East London and South Essex District of the Metropolitan Counties Branch of the British Medical Association.

By CHARLES J. CULLINGWORTH, M.D., F.R.C.P.,
Obstetric Physician to St Thomas's Hospital.

I HAVE chosen the subject of pelvic inflammation this evening mainly for two reasons: First, because it satisfies the conditions courteously imposed upon me by your secretary that I should address you on some topic of everyday interest to the general practitioner and that I should let my remarks be practical; secondly, because I am able to illustrate it from cases that have recently occurred in my own practice at St. Thomas's Hospital.

Those of us who occupy positions on the staff of a hospital ought continually to bear in mind that in accepting those positions we undertake great responsibilities. I have always been of opinion that one of the greatest of these is that of making such use of our exceptional opportunities for clinical observation and research as will best help to advance professional knowledge and give to our brethren who are engaged in the engrossing duties of general practice the benefit of our experience and results.

By pelvic inflammations I mean pelvic peritonitis and pelvic cellulitis, and not inflammations of the various viscera contained in the pelvis. These latter will only be dealt with so far as they are concerned in the pathological processes that lead to the affections I have just named.

Is it possible clinically to distinguish between inflammation of the pelvic peritoneum and inflammation of the pelvic connective tissue? I believe that in most cases it is. The diagnosis is undoubtedly beset with difficulties, and, in the present state of our knowledge, there are still some cases where it is impossible. Year by year, however, their number is diminishing. Our

² *Lancet*, December 11th, 1886.

knowledge of the anatomy of the female pelvis is becoming, chiefly from the study of frozen sections, more precise, while, owing to the advances in abdominal surgery, our opportunities of comparing the physical signs with the actual conditions are much more frequent than they ever were before. Exploration of the pelvis by abdominal section, now so frequently practised, is teaching us on the living subject, and therefore under far more favourable conditions, the lessons that until within recent years we could only learn from an occasional *post-mortem* examination, performed, not only when the disease had advanced too far to teach us anything of its earlier stages, but also when the parts had ceased, from their loss of vital tonicity, to hold the same relative position to each other and to convey the same impression to the touch that they did during life. The knowledge thus gained has revolutionised our ideas about pelvic inflammations. Until quite recently the teaching on this subject was vague and unsatisfactory. Whenever a hard mass was detected above the vaginal roof on one or both sides of the cervix uteri the case was pronounced to be one of pelvic cellulitis with exudation at the base of the broad ligaments. It is quite true that many years ago that excellent French observer, Bernutz, combated this view, and produced evidence to show that those hard irregular masses above the lateral fornices were generally due, not to cellulitis, but to peritonitis. But his words fell on deaf ears, for old beliefs die hard, and men were not prepared to throw over as an exploded error what had appeared to them almost the one pathognomonic sign of an otherwise obscure disease.

And so the teaching continued on the old lines until the subject was again taken up and it became clear from the accumulating evidence of the operating theatre and the *post-mortem* room that these lateral masses of exudation were sometimes due to pelvic peritonitis and sometimes, but much more rarely, to pelvic cellulitis, and that when they were due to pelvic peritonitis they generally consisted of diseased Fallopian tubes, surrounded by thickened peritoneum and involved in a mass of adherent viscera, including almost always the ovary and broad ligament and not infrequently pieces of omentum and one or more coils of intestine.

In the year 1886, Dr. W. M. Polk published a remarkable paper in the first volume of the *Transactions of the Association of American Physicians*, entitled *A Study of Periuterine Inflammation in its Relation to Salpingitis*, based upon Sixteen Cases of Extirpation of the Tubes and Ovaries occurring in Bellevue Hospital. After giving a brief sketch of the history of medical opinion on the subject of pelvic inflammations, and examining the question from the standpoint of analogy, he proceeds to inquire what evidence is furnished by the facts of the necropsy and operating rooms, which are, after all, as he says, "the pivots of our faith in medical science." With reference to the *post-mortem* evidence he states that, at Bellevue Hospital, it is rare, except in the case of patients who have died from septicæmia, to find any proofs of pelvic cellulitis, unless it be clearly secondary to a previous inflammation of the pelvic peritoneum. "On the other hand, nothing is more common than to find evidences of peritonitis about the ends of the tubes, and it is noticeable that in such cases the evidences of such inflammation diminish as you leave the extremities of the tubes." And he sums up this part of the subject by saying that the *post-mortem* evidences go to show that periuterine inflammation "is a product of salpingitis and not secondary to cellulitis, and that if cellulitis is present, it is either secondary to peritonitis, or else is a lesion of septicæmia, in common with those which belong to that disorder when originating in other parts of the body." He then proceeds to give the records of sixteen patients who presented the signs commonly attributed to pelvic cellulitis, "but in whom abdominal section showed salpingitis, periovaritis, and peritonitis. In two of the cases there was slight cedematous swelling of the cellular tissue in the broad ligament, just beneath the spot at which an inflamed tube had rested; in the remainder the most careful examination failed to detect the slightest induration or swelling in any part of the cellular tissue that lay about the uterus or between the peritoneal layers of the ligaments." Bimanual vaginal exploration of the uterus and its surroundings was employed in each case, both before and after the operation. "Before the operation indurated sensitive masses were present about the uterus in each patient, and the mobility of the organ was impaired to a greater or less degree. After the operation the masses could not be found in a single case, and the mobility of the uterus was restored to about its normal range in nearly every instance. During the operation two fingers were placed astride the broad ligaments, every portion

being thus readily explored. The entire pelvic floor was likewise examined by touch, and whenever the method seemed incomplete, the opposing fingers of an assistant were placed in the vagina. Between the two it was impossible for any induration or thickening to escape detection." I have quoted freely from this paper because its conclusions rest on a solid foundation of facts. If time permits I propose to present you with corroborative evidence from my own ward at St. Thomas's. In the meantime, I may say that my experience in abdominal surgery enables me to confirm to the fullest extent the propositions that Dr. Polk lays down.

One word as to the sequence of pathological events in these cases of salpingitis and pelvic peritonitis. Where does the inflammation commence and what is its etiology? In the great majority of cases the inflammation begins in the mucous membrane of the uterus, either from septic absorption, or the poison of gonorrhœa, or the infection of tubercle, or simple catarrh, as from getting the feet wet during menstruation. It spreads from thence along the mucous lining of the Fallopian tubes, just as inflammations of the mucous membrane of the respiratory tract or the urinary organs spread along their various portions. There being no ready means of exit for their secretions in the case of the Fallopian tubes, the discharges, whether of mucus or pus, accumulate, and by so doing aggravate the inflammatory process. The inflammation thus tends to spread still further outwards beyond the fimbriated extremity of the tube to the adjacent peritoneum, and a localised peritonitis is thus established, with its focus at the mouth of the tube. The result of this sometimes is promptly to seal the end of the tube, which then becomes distended by the secretions, and forms a more or less distinct tumour. In other cases the peritonitis appears to commence by the sudden pouring out of the morbid secretions from the fimbriated extremity into the pelvic cavity, setting up a sudden and violent pain, followed by all the signs and symptoms of an attack of acute pelvic inflammation. Up to this point the process has been probably painless. Endometritis and endosalpingitis are not characterised by pain, but the moment the peritoneum is reached pain begins; so that what may seem to be the starting point of an illness often is, in reality, merely an indication that the inflammatory process has passed from a mucous to a serous membrane. The inflammation of the uterine mucous membrane may by this time, especially if it be simply catarrhal, have subsided. The canal of the cervix usually allows free escape of the inflammatory secretions, so that an inflammation of the mucous lining of the uterus is ordinarily a much less serious affection than a similar inflammation in the Fallopian tube, where the readiest outlet is into the peritoneal cavity.

With regard to pelvic cellulitis, there is strong reason to believe that this disease, when it is the primary affection, is always septic, and that it is carried by the lymphatics through the uterine or cervical tissues themselves to the connective tissue immediately adjacent. It is, I know, commonly taught that pelvic cellulitis may be purely traumatic; but the results of antiseptic midwifery and antiseptic surgery show that if a laceration or other wound of the genital canal be guarded from septic infection, no cellulitis will occur. The old idea, that some patients are so susceptible that the mere passing of a uterine sound will set up a cellulitis, puts the blame on the wrong shoulders. Neither a clean sound nor a clean wound ever yet produced a cellulitis. One day a patient was admitted into my ward at St. Thomas's, whose case seemed to be an exception to the rule that a local absorption of septic matter is necessary for the production of pelvic cellulitis. The patient was a widow; her last confinement had taken place fifteen months previous to her admission, and she had made a good recovery. There had been no miscarriage since. Her occupation was that of a monthly nurse, and at the time she was taken ill she was attending a puerperal fever case. Her patient was a poor woman, and she had to wash the soiled linen as well as to administer vaginal injections. After a week's attendance her left middle finger became very sore, and three or four days later she was seized with shivering, abdominal pain, fever, and loss of appetite, and had to give up work. She was laid up at home for a few weeks, and then came into the hospital. All the signs of an extensive pelvic cellulitis were present. I heard the history read, but although it was amply sufficient to explain a septic infection of the finger, it appeared impossible to connect the infection of the finger with the cellulitis in the pelvis. I confessed to my class that here, for the first time since my attention had been called to the question, was a case in which there was a pelvic cellulitis with no history of local infection to account for it, and that the views I had long firmly held seemed in danger. Next

morning my resident told me he had elicited some further facts about the case. The woman had for years worn a vaginal pessary, which she was in the habit of removing every night, and replacing in the morning. Here, then, was the explanation. She had carried the septic infection to her vagina by means of her own finger; and so the case that seemed as though it were going to upset my belief, turned out to be in strict and even striking conformity with it.

I must not, however, lose sight of the object I set before myself in this paper, namely, to discuss the differential diagnosis of pelvic peritonitis and pelvic cellulitis. It will facilitate our study of the distinguishing characters of these two inflammations if we consider for a moment the anatomical distribution of the two tissues involved. The peritoneum in the female pelvis is reflected in front from the anterior abdominal wall on to the fundus of the bladder a little above the top of the symphysis pubis. It thence passes down the posterior surface of the bladder as far as the level of the os uteri internum, when it is reflected upwards and forwards on to the anterior surface of the uterus. This reflection forms the vesico-uterine pouch. After investing the fundus uteri the peritoneum passes downwards and backwards over the posterior surface of the body of the uterus, the posterior surface of the supravaginal portion of the neck of the uterus, and behind the upper fourth of the posterior vaginal wall. Here it is reflected on to the front of the rectum and the posterior pelvic wall, forming a pouch or sac, which is the most dependent part of the peritoneal cavity. This pouch is bounded laterally by two folds of peritoneum, called the utero-sacral ligaments or Douglas's folds, which pass one on each side from the lateral borders of the supravaginal portion of the cervix uteri, upwards and backwards to the sacrum, the upper margin of the folds reaching the sacrum at about the level of the second sacral vertebra. Returning to the uterus, the peritoneum, after investing the body of that organ, is prolonged outwards from its lateral borders in the form of a fold to the lateral wall of the pelvis on each side, forming with the uterus a sort of curtain stretching across the whole breadth of the pelvis. This fold is the broad ligament. It encloses along its upper border the Fallopian tube, which runs outwards to within a short distance of the side of the pelvis, and then leaves the broad ligament to terminate behind it, the last portion of its course describing a curve backwards, downwards, and inwards, so that the fimbriae embrace the outer surface of the ovary, which latter organ lies behind the broad ligament on a lower plane than the main portion of the tube. Immediately below the ovary the layers of the broad ligament separate, becoming reflected on to the anterior and posterior pelvic walls respectively. It must be remembered that, although we speak of the surfaces as anterior and posterior, in the erect posture and in the normal condition the so-called posterior surface is in reality directed upwards and backwards and the anterior downwards and forwards. The cavity of the pelvis being divided transversely by a curtain consisting of the uterus and the broad ligaments, the spaces behind the broad ligaments may be spoken of as the right and left posterior quarters of the pelvis, while those in front may be called the right and left anterior quarters of the pelvis. These latter are so shallow, and, except under very exceptional circumstances, so destitute of important contents that they are of little or no practical importance. On the other hand, the two posterior quarters of the pelvis, with Douglas's pouch between them, possess the highest clinical interest. The latter, because fluids in the peritoneal cavity tend to gravitate into it, and because, dipping down as it does behind the upper part of the vagina, it can easily be reached on vaginal examination; the former, because they contain the free ends of the Fallopian tubes and the ovaries, structures, many of the morbid alterations of which can be ascertained by the bimanual method with considerable accuracy.

Turning now to the distribution of the pelvic connective tissue, we find that, except perhaps over the fundus uteri, a layer underlies the entire pelvic peritoneum, parietal and visceral. The so-called ligaments of the uterus contain a greater or less quantity between the peritoneal folds of which they are composed, while in certain special situations the connective tissue may be said to be abundant, for example, around the supravaginal portion of the cervix, along the base of the broad ligament, and behind the symphysis pubis.

With these facts in mind, we may proceed to inquire—(1) What portions of this widely-distributed tissue would be first affected when cellulitis starts from the body and neck of the uterus respectively? and (2) In what directions would the inflammation

tend to spread? In the case of cellulitis originating from the body of the uterus, the inflammation is transmitted directly through the parenchyma, to the connective tissue between the layers of the upper part of the broad ligament, on one or both sides. One of the diagrams before you shows the exudation limited at first to that portion of the broad ligament adjacent to the uterus. Thence it travels outwards towards the lateral wall of the pelvis, or forwards beneath the anterior layer of the broad ligament to the subperitoneal connective tissue in the anterior abdominal wall. In the former case, having reached the lateral wall of the pelvis, the inflammation may pass either upwards along the psoas muscle to the region of the kidney, or, more commonly, downwards and forwards to the anterior abdominal wall, creeping along above Poupart's ligament, to the mass of fat and cellular tissue lying in front of the bladder, and above and behind the pubes. This same spot is reached by the inflammation if it travels forwards instead of laterally. Thence it may continue to spread upwards beneath the anterior abdominal wall, even as high as, or higher than, the level of the umbilicus.

In the case of cellulitis originating in the cervix, the cellular tissue immediately surrounding the cervix is the first to be affected. The inflammation then spreads along the base of the broad ligament to the lateral wall of the pelvis. It may also affect the vesico-uterine cellular tissues in front, or the utero-sacral ligaments behind, in which latter case the rectum will be surrounded with exudation on at least three of its sides, and in some instances will be completely encircled.

We are now in a position to discuss the differential diagnosis of pelvic inflammations.

1. *With Regard to their Etiology.*—Pelvic peritonitis, in the great majority of cases, is secondary to disease of the lining membrane of the uterus, and, for the most part, travels to the peritoneum along the Fallopian tubes. This mode of origin is found in all the cases due to gonorrhoea, and in all the cases due to catarrh. It is also found in most of the cases resulting from tuberculosis, and from septic infection. In a certain number of cases of septic peritonitis the inflammation spreads directly to the peritoneum through the parenchyma of the uterus, by means of the lymphatics or the veins or the connective tissue, but where this is the case there will be cellulitis as well, for the inflammatory process affects all the tissues *en route*. Pelvic cellulitis, on the other hand, when a primary disease, is always septic, and spreads directly outwards from the body or cervix uteri through the parenchyma. There is therefore no involvement of the Fallopian tube.

These considerations have considerable bearing on the diagnosis. Thus, if a case of pelvic inflammation be known to have a gonorrhoeal or a catarrhal origin, the tissue affected will be the peritoneum, if it occur after abortion or labour or operation, or the passing of a tent or a sound, that is, if it be known to be septic, it may be either a peritonitis or a cellulitis, or both. If it be tubercular, it will be peritonic and not cellulitis.

Of course pelvic peritonitis may arise from other than the causes just mentioned—as from the presence of tumours, the rupture of cysts, hæmorrhage into the peritoneal cavity, disease of the vermiform appendix, intestinal perforation, etc., but in none of these cases will there be associated disease of the Fallopian tube, except as a coincidence, or any of this hard irregular swelling in the lateral fornix, which is the main element of difficulty in discriminating between pelvic peritonitis and pelvic cellulitis.

2. *As to the Amount of Pain.*—It is generally held, and with truth, that the presence of acute pain points to the pelvic inflammation being peritoneal. Cellulitis, when uncomplicated, is a disease not attended with pain, at any rate, not with severe pain. But we must remember that salpingitis is also a painless affection, and that the sudden onset of severe pain in an attack of pelvic inflammation is not to be taken as marking the commencement of the inflammation, but as indicating that it has reached the peritoneum.

After the acute stage of pelvic peritonitis has passed, the pain is only felt after standing and walking. The tenderness, however, remains, and is apparent on vaginal examination and on coitus. Pain, however, is a most misleading symptom, for it is never so marked in cases of actual disease as it is in some neurotic patients who have no pelvic lesion of any kind.

3. *The Physical Signs.*—In the early days of an attack of acute pelvic inflammation physical examination tells us little or nothing. The vagina is hot and tender, and its vessels may be felt pulsating, and that is all. But in chronic inflammation and

in acute inflammation, when exudation has occurred, the physical signs are usually highly characteristic.

In cellulitis arising from the cervix we find either a uniform hardness and resistance in one or both lateral fornices, with depression of the vaginal roof and partial or complete obliteration of the cervix, or a collar of induration immediately in front, or immediately behind the supravaginal portion of the cervix. In pelvic peritonitis, on the other hand, there is no depression of the lateral or anterior fornices, for the exudation is within the peritoneal cavity, and if it encroaches upon the vagina at all, it does so posteriorly and there only. It will then be felt as a distinct swelling in Douglas's pouch, obliterating the posterior fornix and pushing the uterus forwards, if it is extensive, and forming an elastic tongue-shaped swelling behind the cervix and upper part of the posterior vaginal wall, if the amount of exudation is small.

A similar difference exists in the conditions found on rectal examination. In cellulitis affecting the tissues behind the cervix, including the utero-sacral ligaments, the rectum in that region will be felt to be surrounded wholly or partially with a hard belt of exudation. Whereas an exudation due to pelvic peritonitis will be confined to Douglas's pouch, and will be felt to be entirely in front of the bowel, though it may and generally does bulge into it. The swelling, moreover, will be less hard.

When the body of the uterus is the starting point of a cellulitis, and the broad ligament is the seat of the exudation, bimanual examination will reveal a hard, smooth, flattened tumour by the side of the uterus, sometimes displacing it slightly to the opposite side. This tumour can be moved within certain narrow limits backwards and forwards between the two hands.

Later on, when the exudation has reached the subperitoneal connective tissue of the abdominal wall, it will give rise to a smooth hard swelling beneath the abdominal muscles, occupying either the region immediately above Poupart's ligament, which forms its lower boundary, or the suprapubic region. No such swelling or sense of superficial resistance is found in peritonitis.

What, then, are the characteristic signs of pelvic peritonitis? There is a hard, tender, irregular swelling, felt, on bimanual examination, lying posteriorly in the pelvis, above the vaginal roof and not implicating or depressing it. This mass consists of thickened peritoneum surrounding the thickened Fallopian tube and the ovary, the two latter being matted together by adhesions to each other, to the back of the broad ligament and to other parts in the neighbourhood. The uterus is seldom pushed aside by this mass and does not form, as it were, a part of it, as in cellulitis of the broad ligament. If the uterus is displaced at all, it is displaced forwards unless, indeed, it have been retroverted or retroflexed to begin with, when it will have become adherent in its abnormal position. Very often the mass can be felt to extend behind the uterus, and to be adherent to its posterior surface. It is very seldom that any evidence of the presence of a pelvic mass of the kind just described can be obtained on examination of the abdomen. This is only possible when there is superadded an encysted peritoneal effusion, or, as in a case to be related presently, when some of the pelvic viscera have happened to contract adhesions to the abdominal wall.

As time goes on, the hardness and thickening of the peritoneum become, in favourable cases, much less marked; but the viscera once adherent are apt to remain so, and there is generally to be felt a soft irregular mass in the posterior part of the pelvis for the rest of the patient's life, with some amount of fixation of the uterus from adhesions. On the other hand, in cellulitis, the exudation entirely disappears, leaving, in those cases where the exudation was in the substance of the broad ligament, some contraction of the broad ligament, with drawing over of the body of the uterus to the affected side as the only trace. I have not time to go at length into the differential diagnosis of pelvic abscess, but I may just say that these abscesses are by no means all of cellulitic origin. For example, most of those which burst into the rectum are intraperitoneal, being situated behind the uterus, and limited by adhesions. Cellulitic abscesses point, for the most part, above Poupart's ligament, where intraperitoneal abscesses never do, or burst into the bladder, which is rarely the case with encysted collections of pus within the peritoneal cavity. Either form of abscess may burst into the vagina.

Let me illustrate the foregoing remarks by one or two typical cases:—

1. A woman, aged 27, was admitted on October 10th, 1889; she had been married three years and had no children. Six months after marriage she took cold whilst menstruating, and had an

attack of inflammation with suppression of the menstrual flow. She apparently recovered, and remained well until five months before admission, since which time she had suffered from pain, sometimes very acute, in the lower part of the abdomen, with occasional vomiting and faintness and a yellowish discharge from the vagina.

On examination, there was found an ill-defined sense of resistance and tenderness in the left iliac region, with dullness on percussion over an area 3 in. × 2 in. *Per vaginam* the uterus was found anteverted and fixed. On its left side was a hard, irregular, fixed mass, which could be traced from the uterus, running outwards and forwards, to a mass lying beneath the abdominal wall and inseparable from it. This mass corresponded with the area of dullness and resistance in the left iliac region. The patient declared that at this point a swelling occasionally appeared, of about the size of a hen's egg. On the right side of the pelvis a mass could be felt running outwards from the uterus for a short distance, then turning backwards and terminating high up behind the cervix uteri.

The diagnosis was pelvic peritonitis with salpingitis. The swelling on the left was made out to be the left tube enlarged and adherent, terminating in a mass of adherent viscera fixed to the under surface of the abdominal wall. From the patient's description, it was suspected that the mass in which the fimbriated end of the tube terminated would be found to include a coil of intestine. The swelling on the right was believed to be the right tube, thickened by inflammation, surrounded by adhesions, and bent back upon itself so that its fimbriated end lay behind the cervix, to which it had become adherent.

The patient improved so much that she was able to leave the hospital in a month, free from pain, and feeling quite well. She was told to come in for operation if the pain and other symptoms recurred.

In a fortnight she came back, and abdominal section was performed on November 28th. The left tube was found passing outwards and forwards to a large hard, irregular mass, adherent to the pelvic wall, immediately beneath the abdominal parietes and close to the edge of the psoas. The tube and ovary were dug out of the centre of the mass, the former being much thickened and its fimbriated end occluded, the latter healthy in structure but covered with shreds of adherent membrane. The remainder of the mass consisted of intestine and omentum. The right tube and ovary were found dipping down behind the cervix and firmly adherent to their surroundings. The tube was thickened and occluded, the ovary healthy. The tubes and ovaries were removed and the patient made a good recovery, leaving the hospital thirty-one days after the operation.

This case was a good example of pelvic peritonitis due to extension of a catarrhal inflammation of the Fallopian tubes to the peritoneum, the diagnosis being confirmed by the operation in every particular. The physical signs were precisely those that used to be considered distinctive of pelvic cellulitis, of which there was no trace. The treatment of these cases hardly comes within the scope of this paper; but as it may occur to some to inquire why the tubes were removed when the inflammation was no longer in an active state, I will briefly state the reasons. Tubes damaged and adherent to the extent they were in this case cannot possibly perform their functions again, and, as they are a constant source of irritation, interfering with a patient's walking and working powers, and liable on the smallest provocation to set up fresh inflammation in the form of recurrent peritonitis, with the result of increased damage and further adhesion, it appears to me that the proper course is to remove them. This should be done, as a rule, when the acute symptoms have subsided, that is, not during an attack of acute pelvic peritonitis, but as soon afterwards as possible. It is from the point of view of treatment that the differential diagnosis of pelvic inflammations derives its enormous importance. So long as all pelvic inflammations were treated much in the same way—by rest, poultices, and opium—a more precise diagnosis was not of much consequence. But now, when we know that the one form of inflammation can be advantageously dealt with by operation, while the other—except, of course, in the case of an abscess forming—cannot, the differential diagnosis has come to be a matter of the greatest moment. It was my strong conviction upon this point that, amongst other things, led me to bring the matter before you this evening.

The next case I wish briefly to relate is an instance of septic salpingitis following delivery. The inflammation was apparently roused into renewed activity six weeks later, by the patient get-

ting her clothing wet through, an attack of pelvic peritonitis supervening within a few hours.

2. The patient was a young widow aged 23, admitted May 17th, 1890. She had been confined October 8th, 1888, and had shortly afterwards been admitted to the surgical wards at St. Thomas's for breast abscess. On the day she left the hospital, it was raining heavily, and she got drenched. That night she was attacked with severe pain in the lower part of the abdomen, striking down the thighs. Three weeks later she was admitted into Adelaide Ward. There was then, in addition to a hard ridge of induration in front of the cervix, due to cellulitis, a mass in each lateral fornix, running outwards from the body of the uterus, and more marked on the left side. The patient improved quickly, and left the hospital January 19th, 1889. From that time up to April, 1890, she remained well, and able to do her work. In May she was readmitted to the hospital under my care. The uterus was then in its normal position. On each side of it there was an irregular hard swelling. In the midst of the mass, on the right, there was a softer and more circumscribed swelling. The right tube could be made out quite distinctly, bending backwards upon itself, and dipping down behind the uterus. The posterior fornix was very tender. A few days later there was observed a fluctuating swelling in the retrouterine pouch, tense and globular. On June 6th the fluid had disappeared, and the contents of the pouch of Douglas were felt to be hard and nodulated. The right tube could again be distinctly traced.

Operation was proposed to the patient, and her consent having been given, abdominal section was performed on June 6th. On putting the fingers into the pelvis, a quantity of serum escaped. Tracing the right tube from the cornu of the uterus, it was found thickened, bent upon itself, and adherent behind and to the right of the uterus. In the angle of flexion was the ovary, enlarged to the size of a pigeon's egg, and containing several cysts. The left ovary and tube were adherent, though apparently themselves unaltered. The ovaries and tubes on both sides were separated from their attachments with some difficulty, and removed. There were still some thickened masses in the retrouterine space, but as it seemed certain that they were merely portions of thickened peritoneum, they were left undisturbed.

On examining the parts removed, the right tube was found thickened, its diameter measuring half an inch; its external surface was covered with vascular shreds of adherent peritoneum. Its fimbriated extremity showed the fimbriae thrown back and adherent, the opening in the tube being contracted to the size of a mere pin hole. On section, the tube was apparently empty; its mucous membrane was pale, swollen, and cedematous; its muscular coat was white and dense, and from a quarter of an inch to three-sixteenths of an inch in thickness. There were two cysts beneath the peritoneal covering of the tube on its upper surface. The left tube was denser and firmer than natural, and of about the thickness of a quill; its outer end was much contracted, but still pervious. The mucous membrane lining the inner half of the tube was normal; that lining the fimbriated end was swollen and soft, and of a livid red colour. A section of the right tube was examined under the microscope, and showed small-celled infiltration, with a few loose pus cells in the lumen of the tube.

The patient made an excellent recovery, and was sent to a convalescent home at the end of a month, feeling quite well.

I could easily bring before you other similar examples, were there time to do so, but I must be content with a representative selection, and will proceed to give a brief summary of a very severe case of pelvic peritonitis following gonorrhœa.

3. A young woman, aged 28, with a worn, pale face, and wretchedly thin, was admitted September 10th, 1889, complaining of severe pain in the lower part of the abdomen, and with a temperature of $101\frac{1}{2}^{\circ}$. She had been married five years, but had been separated from her husband for three years on account of his intemperance and cruelty, and during this time had had to maintain herself and her two children by dressmaking. Only on one occasion since their separation had she and her husband cohabited. This act of intercourse took place twelve months before her admission. Very soon afterwards she began to have a profuse yellow discharge; it was evident that he had infected her with gonorrhœa. In a few weeks she became too ill to continue at her work, and had to give up her home and go into the parish infirmary with her children. She came out in three or four months, but soon had to return. She again took her discharge, and resumed her occupation. Her health, however, soon gave way. She suffered great pain in the lower part of the abdomen, in the groins and in the back, and

eventually sitting became so difficult and painful that she had to relinquish her employment, and for some weeks before admission had subsisted on the generosity of friends. The pelvis was filled with a hard, irregular mass, especially to the right of the uterus and behind it. The uterus was of normal size, pushed over to the left of the middle line and fixed. There was no depression of the lateral fornices.

Abdominal section was proposed, and readily agreed to. The operation was performed on September 14th, 1889. There was no general peritonitis, but the pelvis was occupied by a mass of adherent viscera, difficult to recognise and separate. The uterus, of normal size, lay in front and to the left. The left tube, much thickened and distorted (Fig. 1), lay mainly behind the uterus, where it

Fig. 1.—*a*, Purulent salpingitis, the result of gonorrhœa. A bristle has been passed into the uterine end; at a little distance beyond it the tube suddenly expands and turns sharply downwards. The fimbriated end is open. The mesosalpinx, or adjacent portion of broad ligament, is seen to be much thickened from secondary cellulitis. *b*, Transverse section of same, actual size. Walls of tube enormously thickened; little or no dilatation of the canal; secondary cellulitis in the mesosalpinx.

was firmly adherent. On removing it, a small cavity was exposed, containing pus and broken-down blood clot—evidently a suppurating intraperitoneal hæmatocele. The broad ligament, and especially the mesosalpinx, were swollen and friable from secondary cellulitis, so that the ligatures tore through the tissues. The right tube was less thickened, but much distorted, and very firmly adherent to the pelvic wall and all the surrounding parts; it terminated behind the uterus. The cæcum and appendix were so deeply involved in the adhesions, that the tube was not removed. Adhesions were, however, separated sufficiently to open another intraperitoneal abscess of a similar character to that on the left side. The pelvis was flushed with hot boracic acid solution, and a drainage tube was left in. The operation lasted an hour and a half.

The patient had a severe attack of pneumonia after the operation, and there was a good deal of suppuration through the drainage tube before the abdominal wound entirely closed, but she eventually made an excellent recovery, gaining flesh, and looking quite bright and cheerful, and was able to be sent to a convalescent home on October 29th. Before going out, she complained of a vaginal discharge, which, on examination, proved to be due to a purulent inflammation of the urethra and nymphæ, and cervical catarrh, for which she underwent the usual treatment before she left the hospital. The gonorrhœal origin of the pelvic inflammation was thus abundantly confirmed. This was a typical example of the class of cases that until recently were regarded as cellulitic, and beyond the reach of surgical treatment.

The series would be incomplete without an example of tubercular inflammation of the pelvis. I have records of several such cases, but must limit myself to one well-marked instance.

4. The wife of a country stationmaster, aged 30, was sent to me by her doctor in December, 1889, on account of severe paroxysmal pain in the lower part of the abdomen and back, with great depression of spirits and considerable emotional disturbance. Her symptoms were thought to be due to hysteria. Her history was briefly as follows: At the age of 11 she was very delicate, and was declared to be threatened with consumption. After a visit to Brighton, however, she improved greatly, and at 17 she weighed 10 stone. Her health remained good up to her marriage in April, 1889. In July she had a miscarriage, without any unusual symptom. At the end of a week she left her bed, feeling fairly well. Three days afterwards she awoke at 2 in the morning with acute pain in the lower part of the abdomen and in the back. This lasted some hours, then passed off, returning the following night, and again disappearing during the day. After the first week the pain returned occasionally, but less violently, until a month or six weeks had elapsed, when it again became very severe. About six weeks ago she had another severe attack, thought, as before, to be neuralgic. All this time menstruation was irregular, the intervals prolonged, the periods profuse. She was sent away for change of air, and seemed better, but on returning home she again had attacks of pelvic pain, and also pain in the back of the neck and right arm and shoulder. She had by this time lost flesh considerably. Lately she had suffered from night sweats, but had no cough or expectoration.

On examining her chest I found slight dulness at the right apex, with harsh breathing, prolonged expiration, and increased vocal resonance. There were no moist sounds or other evidence of active disease. *Per vaginam* the uterus was normal in length and direction; its mobility was impaired. The sound, on withdrawal, was stained with purulent mucus. High up on the right side a resistant mass could be felt, in which the right tube could be distinguished, swollen, tender, and tortuous, passing behind the uterus and the ovary, rather large and tender, surrounded by the tube

Fig. 2.—Tubercle of the Fallopian tube, showing general enlargement of the tube, both in length and breadth, with irregular dilatations, corresponding to deep ulcers on the inner surface, filled with caseating tubercle. Miliary tubercles are seen on the peritoneal surface, chiefly near the fimbriated end. The uterine end of the tube is much twisted. The fimbriated end is nearly occluded by purse-string contraction.

There was no depression of the lateral fornix. Nothing abnormal was detected on the other side. The diagnosis I ventured to make was: Tubercular disease in an early stage at the right apex; chronic inflammation of the right tube and pelvic peritoneum, due either to septic infection or tuberculosis, more probably the latter.

I suggested abdominal section, which was agreed to, and the patient was accordingly admitted to St. Thomas's Home, December 21st, 1889, and operated upon on the 24th. The pelvis was entirely occupied by thickened peritoneum and adherent viscera. The peritoneal surface of the uterus, Fallopian tubes, and adjacent coils of intestine were studded with miliary tubercles. The uterus was pushed forwards by a mass behind it, consisting of a small cystic ovary with the enlarged and thickened right tube curving round it, the whole so densely adherent that nearly an hour was occupied in the separation. The left tube was exceedingly tortuous, much thickened, and universally adherent, the adhesions being more recent than those on the right side. The left ovary contained a large cyst, but was not enlarged. It was entirely surrounded by adhesions, and its external covering was thickened and opaque.

On examining the parts removed, the portion of right tube was 3 inches long and $1\frac{1}{2}$ broad. The walls were like cartilage, and measured $\frac{1}{8}$ to $\frac{1}{4}$ of an inch in thickness. The mucous membrane was much swollen. The free end of the tube was buried amongst the adhesions. The portion of left tube (Fig. 2) measured, while still unstretched, $2\frac{1}{2}$ inches. It was twisted at its uterine

end like a corkscrew. At the centre there was a dilated portion $\frac{3}{4}$ of an inch long, softer than the rest of the tube. The mucous membrane was deeply ulcerated, showing deep pits full of caseating tubercle. A water-colour drawing was made of the left tube while still perfectly fresh, and an excellent photograph was taken of its appearance when laid open by longitudinal section, showing the rugged ulcers in which the caseous material was found. I am glad to be able to show you both the photograph and the drawing, and also a beautiful drawing (Fig. 3) by

Fig. 3.—Tubercular salpingitis. A, Transverse section of tube under a power of $\times 50$; and B, section through a tubercular nodule under a power of $\times 250$, showing two giant cells. a, Peritoneum; b, tubercular nodules in subperitoneal tissue; c, longitudinal muscular coat of Fallopian tube; d, blood vessels; e, circular muscular coat; f, hypertrophied mucous membrane, showing numerous tubercular nodules containing giant cells—the lining epithelium still remains in places; g, lumen of tube; h, remains of ciliated epithelium; i, giant cells; k, epithelioid cells; l, space lined with columnar epithelium.

my late resident, Mr. C. H. James, of the microscopic appearances of a section of a tube, in which the characteristic giant cells are seen in abundance.

Unfortunately the patient died from the effects of the operation. I never conceal from my patients or from myself that this operation, though generally successful, entails a grave immediate risk.

I do not know that I need detain you by much comment on this case. It was a good example of pelvic peritonitis on both sides, dependent on tubercular inflammation of the Fallopian tube, that on the right side being of old date, that on the left more recent. Though the operation in this particular case was unsuccessful, I do not for a moment regret having advised it. It is evident that the patient could not have lived very long as she was, and the removal of the parts principally affected afforded her the only possible chance of recovery.

These four cases illustrate very well the usual mode in which pelvic peritonitis originates, namely, from salpingitis; they also illustrate the four varieties of this later affection, namely, the catarrhal, the septic, the gonorrhoeal, and the tubercular. There are other classifications of salpingitis besides this etiological one, but none so free from objection or so practically useful.

It must not be assumed, because abdominal section was performed in all these cases, that I recognise no other treatment in these affections. I have merely chosen them for my purpose this evening because they are pathologically complete. The state of matters in the pelvis has in each case been actually demonstrated, and the diagnosis has been lifted out of the region of probability into that of certainty. The work of abdominal sur-

gery has the same effect as that of the *post-mortem* room; it makes one more minute and careful in one's diagnosis, and gradually also more and more accurate.

My paper would, no doubt, have been more complete if a case of pelvic cellulitis had been added, but I have already detained you too long. I trust that, incomplete and otherwise imperfect though this paper undoubtedly is, it may, nevertheless, help to show the direction in which advances are being made in our knowledge of the diseases that occur in the female pelvis, and to shed some little light upon a subject with reference to which we have still much to learn.

MEMORANDA:

MEDICAL, SURGICAL, OBSTETRICAL, THERAPEUTICAL, PATHOLOGICAL, Etc.

FRACTURE OF THE ANTERIOR SUPERIOR SPINOUS PROCESS OF THE ILIUM.

I wish to place on record a fracture which I believe to be of very rare occurrence *per se*. Whilst acting as *locum tenens* for the "visiting surgeon" at the Chester Infirmary, I was summoned to see the following case.

The patient (who was a powerfully-made bricklayer about 35 years of age) gave a history of having slipped on the kerbstone, and fallen on his side. On examination I found that the anterior superior spinous process of the right ilium was fractured and very freely movable. He had sustained no other injuries. In three weeks' time perfect union had taken place and he was able to resume his work.

Dublin.

J. N. EUSTACE, M.B., B.Ch.

SUDDEN BRONCHIAL OBSTRUCTION.

I WAS summoned in great haste to see a woman, aged 35, who for some years had suffered from phthisis. A severe hæmorrhage occurred one year ago. She is an active woman, and professes to enjoy good health, beyond a slight pain in her chest and cough if she catches cold. From previous examinations I knew she had a large cavity in the upper part of the right lung, almost dry, while the left lung was sound. On her way to church she had been suddenly seized with a violent fit of coughing, which lasted in a paroxysmal fashion for several minutes; she returned home in a very breathless condition. On my arrival in about ten minutes from the beginning of the attack, I found her suffering from what at first sight looked like an asthmatic attack; her breathing, hurried and laboured, was accompanied by much wheezing. From inspection and percussion, I could gain no information, but, on auscultating over the right side of the chest, I was struck by hearing loud snoring *râles*, a new feature, which had a definite point of maximum intensity in the second interspace, about four inches from the mediastinal line. The *râles* accompanied both inspiration and expiration, and were not removed or altered in any way by frequent spasms of coughing. The left lung was perfectly free from any morbid physical signs. Feeling sure of some obstruction, I gave a simple stimulating expectorant. This state of things continued till the evening of the next day, when I was again sent for; the patient, after a violent cough, had brought up a substance which she thought was bone; it was an elongated body, of irregular shape, weighing three grains, and composed of carbonate of calcium and pus. I now found her breathing quite comfortably, and without the slightest feeling of discomfort; she obtained immediate relief on coughing out the obstructing body. The stridulous breathing and rhonchi I found had quite gone, and the lung appeared in its former condition of chronic phthisis. The features of interest are: 1. The sudden onset from no apparent cause. 2. The instantaneous relief on bringing away the foreign body. 3. The composition of the foreign body,

Truro.

F. HAROLD CARLYON, M.B., C.M.

COEXISTENCE OF ERUPTIVE FEVERS.

IN THE BRITISH MEDICAL JOURNAL of November 8th is a short note by Mr. Moore, of Falmouth, of the coexistence of measles and scarlatina in the same patient. Cases of two different kinds of eruptive fever in one patient at the same time are not, I think, uncommon

Some years ago a case of small-pox and scarlatina in a child came under the care of two practitioners in this city—Dr. Giles and Mr. Weaving, both men of experience, neither of them now living.

Mr. Weaving was asked to see a child who had been ill for some days. On making his visit he found the child under an attack of scarlatina, well marked in every respect, and he told the mother that the child had scarlet fever. He did not see the child again. Some days afterwards Dr. Giles was asked to see the child. The mother told him that the child had been seen by Mr. Weaving, who pronounced the case to be scarlet fever. Dr. Giles saw that the child was suffering under the characteristic eruption and other symptoms of small-pox; there could be no mistake about it. After giving directions for the treatment, he called on Mr. Weaving to tell him, as a friend, of the mistake he had made in diagnosis. Mr. Weaving, however, would not acknowledge that he made a mistake. He said the symptoms of scarlatina at the time of his visit could not be mistaken by any practitioner. They went together to see the child, and Mr. Weaving then agreed with Dr. Giles that the case was clearly one of small-pox. They visited the child together during the illness, and as the symptoms of small-pox passed off, those of scarlatina again came out. The two practitioners attended the child throughout the attack of scarlatina as they had done through the small-pox. The child recovered favourably.

Oxford.

E. L. HUSSEY.

LATERAL SUPPORT OF THE PERINEUM.

NOT having noticed in print any mention of a way of supporting the perineum in labour which has been a great comfort to myself, I desire to draw the attention of practitioners to it, so that they may give it a trial, if they have not already done so. Leaving the schools with no definite idea how to avoid laceration, and with a prejudice against support, many like myself looked forward to the passage of the head with anxiety, and an unpleasant feeling of helplessness. "Doing nothing" was intolerable, and "support" had been condemned by our teachers as bad practice. The only resource was to give chloroform, and by relaxation of the perineum, hope to avert disaster. In this frame of mind I resorted to the following plan, which has not yet disappointed me. The hand is partially closed; so that the tip of the thumb presses on the right side, and the tips of two or three fingers on the left side of the perineum. The thumb and fingers are separated by a space of three or four inches, and no pressure whatever is exerted on this area, but firm pressure reaches the fetal head through the more fleshy tissues on each side and directs its course forward. More pressure can thus be safely applied than by inserting the finger into the rectum and pushing the head forward—a somewhat questionable proceeding. The advantages derived are:

1. Control over the downward course of the head.
2. More or less influence in preventing the recession of the head between the pains.
3. Pains are excited by the firm pressure on the perineal nerves.
- 4, and not the least, the sense of support to the patient at a spot where, and at a moment when most needed.

The plan aims at directing the head forwards, and is totally different from the old method of "supporting the perineum," with the palm of the hand, which if done efficiently tends to cut, as with a knife, the thin and friable tissues in the middle line.

HOLLAND J. COTTON.

Sloane Street.

CHOLERA IN JAPAN.—The epidemic of cholera which is at present raging in Japan is one of extraordinary severity. It is said to have followed an outbreak of influenza which commenced in February last, and reached its climax in April, dying away towards the beginning of summer. As soon as the influenza disappeared, however, cholera seems to have been imported from China, and, as the *Sei-i-Kwai Medical Journal* of October 25th says, "furiously invaded with tremendous rapidity." The terrible nature of the visitation may be judged from the fact that up to October 2nd, 33,863 persons had been attacked, of whom 22,560 died. Owing to the vigorous efforts of the sanitary authorities the epidemic is now, according to our contemporary, "rather rapidly fading away."