THERAPEUTIC TRIALS OF DIETHYL-STILBOESTROL

BY

ALFRED A. LOESER, M.D. Berlin, L.R.C.P. & S. Ed.

The product diethyl-stilboestrol, synthesized by Dodds, Golberg, Lawson, and Robinson (1938), possesses oestrogenic properties which have been described by Dodds, Lawson, and Noble, and later by Folley and Watson. Investigations on human subjects were undertaken by Guldberg. In the light of the evidence so obtained it appeared well worth while to try this preparation in gynaecological practice.

Tests for Harmlessness

I first tried the preparation on myself. On the first day I took one tablet of 0.1 mg., on the second day one of 1 mg.; on the third day I gave myself an intramuscular injection of 1 mg., and on the fourth day an injection of 1 mg. in oily solution. There were no unpleasant symptoms or any pain following the injections. Trials were then carried out on three women with regular menstruation and normal genitalia. Two of them were given injections totalling 5 and 12 mg. diethyl-stilboestrol, respectively, immediately after menstruation: 1 mg. of the substance was given daily by intramuscular injection. The period started in one case five days and in the other six days before it was due. In the third woman the injections were begun on the eleventh day following menstruation, 2 mg. being injected daily for seven days. Here again the period started one week too early. After the period these injections were continued, and a total amount of 33.2 mg. was given. The next menstruation began one week late. The following symptoms were observed in connexion with the treatment in this case. As soon as the individual dose was increased to 3 mg., nausea occurred several hours after the injection. During treatment the breasts became sensitive, the nipples and some small warts on the patient's arm and back became more deeply pigmented, and the clitoris swelled, though only very slightly; libido was increased, and also the general feeling of well-being. When the injections were stopped all these effects disappeared within a week.

Trials for Efficacy

Patients with minor degrees of ovarian insufficiency, such as every gynaecologist sees daily during consultations, were treated. More serious cases of ovarian disturbances were not used for these trials, as it is very difficult to judge a new preparation which only has a substitution effect in complicated endocrine cases.

(a) *Four Cases with Climacteric Disorders.*—The main complaints of these patients were frequent attacks, day and night, of flushing and palpitations and excessive sweating. One of them still had occasional menstruations. They received one 1-mg. tablet daily. The symptoms diminished as early as the fourth or fifth day of treatment, and they gradually disappeared altogether. If the treatment was interrupted the attacks returned. One of these patients, who still had occasional periods, suffered from severe nausea about eight to ten hours after taking the tablets.

(b) *Four Cases of Dysmenorrhoea.*—Three of the patients were young girls, and one, 24 years old, was married. They all suffered from severe premenstrual pains. Treatment was instituted either by mouth or by injection fourteen to eighteen days before the period was expected. Three patients received between 20 and 30 mg. diethyl-stilboestrol. These three found their severely marked premenstrual pains definitely reduced

and made tolerable, while the fourth was entirely free from pain, and was surprised when her period started one week before the expected time. Two patients stated that they experienced nausea some hours after taking the tablets.

(c) *A Case of Hypo-oligomenorrhoea and a Case of Secondary Amenorrhoea.*—The patient suffering from hypo-oligomenorrhoea received a total amount of 28 mg. The period following this appeared after thirty days, and lasted for three days instead of barely one day as usual. The patient suffering from secondary amenorrhoea merits more detailed mention. For years she had had irregular periods, often menstruating only two or three times a year. The last period occurred six months ago. The uterus is very small, rigid, and infantile, and the adnexa are not palpable. A stroke curetting revealed poorly developed glandular structure with marked degenerative change of the lining epithelium. The sound length of the uterus was two inches. A course of 11 mg. of diethyl-stilboestrol was given by intramuscular injections, 1 mg. being injected seven times and 2 mg. twice. On the tenth day after treatment was begun another one-stroke curetting was performed. Histologically a more normal structure of the glandular elements was now noted: there was absence of the marked degeneration observed previously.* The patient always experienced severe nausea several hours after the injections and on one occasion had unquenchable thirst (after 2 mg.). Then treatment with 1-mg. tablets was continued until a total dosage of 22 mg. was reached, and prolution was also given, without menstrual bleeding being brought about. The uterus grew half an inch. The vulva and vaginal tissues were looser than before, and more cyanotic. Drawing pains were felt in the abdomen at the time when menstruation should have taken place. This failure is in contrast to the results obtained by Guldberg, where vaginal bleeding was achieved in the case of a castrated woman patient by means of 20.25 mg. followed by synthetic progesterone.

(d) *A Case of Ovarian Insufficiency following Supravaginal Amputation of the Uterus.*—A 30-year-old patient, who in 1937 had been operated on by me for supravaginal amputation owing to numerous large uterine fibromas incarcerated in the pelvis, started to have distressing flushings and sweatings one year after the operation, when both ovaries, which showed cystic changes to some extent, had been left. Prompt relief of these attacks was obtained, first of all with injections, then with tablets of diethyl-stilboestrol; the latter, however, invariably caused severe nausea and sometimes vomiting in this case. Even after only 5 mg. diethyl-stilboestrol had been given, relief of the flushings was noticed.

(e) *A Case of Pruritus Vulvae.*—The prompt cure of a woman aged 24 was achieved by means of diethyl-stilboestrol tablets—one 1-mg. tablet daily for ten days, without any other treatment. The pruritus reappeared on cessation of the treatment, but a successful result quickly followed when treatment was renewed.

The synthetic preparation ethinyl oestradiol produced gastro-intestinal symptoms which were too severe to enable therapeutic trials to be carried out with that substance.

Conclusions

Diethyl-stilboestrol shows such marked oestrogenic effects that it has a good substitution action in various forms of ovarian insufficiency. In six of the fifteen cases for which it was used unpleasant secondary effects such as nausea or vomiting were produced.

REFERENCES

- Dodds, E. C., Golberg, L., Lawson, W., and Robinson, R. (1938). *Nature*, **141**, 247.
— Lawson, W., and Noble, R. L. (1938). *Lancet*, **1**, 1389.
Folley, S. J., and Watson, H. M. S. (1938). *Ibid.*, **2**, 423.
Guldberg, E. (1938a). *Ugeskr. Laeg.*, **100**, 854.
— (1938b). *Zbl. Gynäk.*, **62**, 2584.

* For the histological observations in this case I am indebted to Dr. W. A. Oliver of St. Bartholomew's Hospital.