the cavity has begun, the air must be replaced by oil. This "anti-symphysis" oleothorax sometimes works, preventing the further shrinking in of the cavity. If it does not, then one must try to establish collapse by means of thoracoplasty.

Conclusions

Extrapleural pneumothorax is an "easy" operation for both doctor and patient. But this statement must be qualified by pointing out that a variety of complications are possible. The "optimistic" literature at present available is based on but brief experience. Having had nine years' experience, I believe that much greater caution must be urged. *Thoracoplasty and its various modifications and plombage are superior to extrapleural pneumothorax.* When it is possible to apply them, extrapleural pneumothorax should not be thought of. It should only be considered when pneumothorax fails as a result of widespread adhesion, when the extent of this does not allow of plombage, or when the general condition or the state of the other lung does not permit of a thoracoplasty. That was approximately the conclusion I reached in my first publication in 1931, and after many digressions I have now returned to it. Within these narrow indications extrapleural pneumothorax is the only possible method of collapse therapy. It may be employed even though one cannot expect much from it. But the rare permanent results gain in value when one considers the hopeless fate of the patients for whom this form of therapy is recommended.

A doubtful indication is in recent cavities with delicate pleural adhesions, where a thoracoplasty seems unsuitable on account of the risk of mediastinal flutter, and a plombe because of the likelihood of its slipping down. These cases are relatively rare. We now perform extrapleural pneumothorax for them only so long as we can assume a sufficient degree of pleural thickening. Then we replace it by thoracoplasty (apical thoracoplasty). Extrapleural pneumothorax here plays the part of a preparation for rib resection.

REFERENCES

- Nissen, R. (1931). *Ther. d. Gegenw.*, **72**, 433, 497.
 — (1932). *Über die neuere Entwicklung der chirurgischen Behandlung der Lungentuberkulose*, Urban and Schwarzenberg, Berlin.
 — (1933). *Med. Klinik*, **29**, 767.
 — (1937). *Schweiz. med. Wschr.*, **67**, 334.

ANAESTHESIA IN THORACIC SURGERY

BY

J. T. HUNTER, M.R.C.S., L.R.C.P.

Anaesthetist to the Brompton Hospital, and the Prince of Wales's Hospital, Tottenham

In no other branch of surgery is the anaesthetist faced with such difficulties as in thoracic operations: each operation presents one or more problems that are found in these cases alone. The procedures are not only extensive, but they take a long time and cause a good deal of haemorrhage and shock. Added to these features is the ever-present question of the correct method of dealing with a case that has sputum in any quantity.

Beyond the technical difficulties, there is the fact that the surgeon is practically always operating on a patient with a disease of long standing and of a nature which causes a degree of toxic absorption.

Preparation of Patient

Generally speaking, the preparation of the patient is just the same as for any other operation of any magnitude. Rest, increased fluids, and glucose, with attention to oral hygiene and the intestinal tract, are essential, and should be carried out as thoroughly as possible, provided the patient will not suffer by any delay. Where there is any sputum a course of postural drainage is invaluable in reducing the quantity at the time of operation, and in the intrapleural operations an artificial pneumothorax is necessary on the affected side, and should be induced some three or four days before operation. Premedication should be of the lightest. Perhaps the most useful combination is omnopon 1/3 grain and scopolamine 1/150 grain, given three-quarters of an hour before operation. No drug should be used that will diminish the cough reflex to any extent. Basal narcosis may suit nervous patients, but drugs of transient action that are quickly and easily detoxicated should be employed. Evipan and sodium pentothal are the most useful in this group, and of the two sodium pentothal is probably the better. Though avertin is an excellent basal narcotic, its action is prolonged, and where a quick return to consciousness is desirable, as in these cases, it is to be avoided.

Local Anaesthesia

This is used extensively on the Continent, but comparatively rarely in the British Isles. The infiltration of the area to be operated on is carried out with different preparations of novocain in varying strengths, but where an experienced anaesthetist is available there is no doubt that other methods are preferable. Infiltration takes a long time and needs multiple skin punctures. Surgeons in this country seem to prefer, where major operations are concerned, to operate on an unconscious patient—and, moreover, the average British patient would rather be unconscious of the operation if it is to take any length of time. A number of operations can, however, be carried out easily and successfully by this method. These will be classified later.

Spinal Analgesia

High spinal analgesia has proved excellent in lobectomy and thoracoplasty. One of the first papers on this subject was published by Shields (1935) of Toronto, and it

The following medical promotions in and appointments to the Venerable Order of the Hospital of St. John of Jerusalem were announced in the *London Gazette* of January 3: as Knight, Dr. L. W. Dryland; as Commanders, Dr. A. G. Osborn (since deceased), Colonel G. C. E. Simpson, O.B.E., T.D., Hon. W. H. Kauntze, C.M.G., M.B.E., M.D., Dr. A. G. S. Broughton, Lieutenant-Colonel W. Blackwood, D.S.O., Captain F. H. R. Heath; as Officers, Dr. J. A. Carman, Dr. J. A. Kennedy, Dr. H. M. Palmer, Dr. D. E. Brown, M.B.E., Lieutenant-Colonel W. R. Stewart, C.I.E., I.M.S., Colonel H. Stott, O.B.E., I.M.S., Colonel W. Archibald, Dr. F. W. M. Palmer, Colonel C. H. S. Redmond, T.D., A.M.S., Lieutenant-Colonel F. W. W. Dawson, Dr. R. Y. Stones, O.B.E., M.C., Dr. J. B. Mitchell, O.B.E., Captain D. C. Low, Lieutenant-Colonel G. C. Lawson, Dr. Sybil C. Bevan; as Serving Brothers, Dr. W. F. J. Whitley, Dr. C. R. Corfield, Dr. R. W. Jubb, Captain R. P. Cormack, Dr. Francis Liddell, Dr. W. B. Grant, Dr. W. Wood, Dr. J. A. Hendry, Dr. J. B. Aickin, Dr. S. M. Vassallo, Dr. D. G. Hunter; as Associate Serving Brother, Mr. D. D. Variava, O.B.E.; as Serving Sisters, Dr. Alice M. Benham, Dr. Gladys E. Wilkinson, Dr. Mildred M. E. Y. Edwards.

describes a number of cases done under spinal procaine. Shields found that local infiltration was unsatisfactory, for various reasons already mentioned, and that cyanosis and trouble with sputum were a drawback in nitrous-oxide-oxygen anaesthesia. These facts led him to try high spinal analgesia. He chose the space between L. 3 and 4 as the site of injection and, using procaine as his agent, obtained the correct level of anaesthesia by using 200 mg. dissolved in 10 c.cm. of the patient's cerebrospinal fluid. The injection was made with the patient lying on his affected side. The solution was run in slowly, taking two seconds for each cubic centimetre. This usually gave him anaesthesia to the level of the fourth rib. In some cases it had to be supplemented by local infiltration in the region of the upper end of the incision. Nearly all the cases were thoracoplasties. He reported very little fall in blood pressure—much less than in upper abdominal surgery under spinal anaesthesia. Ephedrine $\frac{1}{2}$ grain was injected intramuscularly at the time of the introduction of the spinal needle.

Magill (1936a) describes a modification of the Howard Jones technique, using procaine 1 in 1,500, and reports very favourably on the method. The technique for injection was as described by Howard Jones (1930), and then, with the patient in the ventral position, the head and feet were dropped so that the highest point was about the region of the third dorsal vertebra. This posture was maintained for six minutes. In a series of cases in which the method was carried out by myself the patient was kept in that position for at least ten minutes. After this he was put into a slight Trendelenburg position, and then placed on his side ready for operation. A further modification was described by Harris and Rink (1937). In their series of cases the injection was made according to the Howard Jones technique, with the affected side uppermost. The shoulders were kept slightly higher than the pelvis, with the head dropped so that the highest point was as before, in the region of the third dorsal vertebra. This method presents one or two advantages: first, it is not necessary to turn the patient; secondly, a unilateral analgesia is produced, leaving the accessory muscles of respiration unimpaired on the sound side; thirdly, there seems to be even less tendency for the blood pressure to fall in these cases.

Yet another technique is used; this is a modification of that described by Etherington-Wilson (1934). The patient sits up on the table and 14 to 16 c.cm. of percaïne solution 1 in 1,500 is injected between L. 2 and 3. His head is depressed forwards, so that again the highest point is the third dorsal vertebra, above which the level cannot rise. If this posture is maintained, the time the patient is kept in it is not as important as in the original technique described. Lately it has been found possible to produce perfect anaesthesia with as little as 8 c.cm. of fluid. Weaker solutions of percaïne—that is, 1 in 2,000—have been tried, but the results have not been so good.

The advantages of spinal anaesthesia may be stated as follows:

1. Bleeding is reduced to a minimum.
2. The cough reflex is quite unimpaired.
3. In spite of the fact that one pleural cavity is opened and that no oxygen is being administered under positive pressure, respiration seems to be undisturbed and there is no sign of cyanosis.

The disadvantages are:

1. As in spinal analgesia in other branches of surgery, there is a tendency to post-operative headache.

2. A fall in blood pressure occurs. This can be treated by the intravenous injection of ephedrine in $\frac{1}{2}$ -grain doses, and other therapeutic measures.

3. The surgeon is operating on a conscious patient, but by the administration of omnopon $\frac{1}{3}$ grain and scopolamine $\frac{1}{150}$ grain, combined with two or three grains of nembutal injected intravenously, all memory of the operating theatre is destroyed and, indeed, the patient will probably sleep peacefully throughout the whole operation. Alternatively, a light cyclopropane anaesthesia may be combined with the spinal analgesia.

General Anaesthesia

This is most commonly used in this country for thoracoplasty, apicolysis, for the establishment of extrapleural pneumothorax, and for lobectomy and pneumonectomy. It is proposed to consider the relative advantages of each agent in turn.

Ether is disregarded because of its irritating effect on the mucous membrane and its tendency to cause secretion of mucus. Incidentally, the increased respiratory movement arising out of its stimulating propensities may inconvenience the operator.

Chloroform will produce quiet peaceful anaesthesia, but its toxic effect on the liver is a factor that cannot be overlooked in these operations. The lengthy administrations necessary cause liver dysfunction and acidosis, which definitely militate against smooth convalescence. However, chloroform is a useful adjunct to gas-and-oxygen when given in minute doses in difficult cases.

Divinyl ether is a comparatively new agent which also forms a useful adjunct to nitrous-oxide-oxygen and does not possess the toxic properties of chloroform.

Nitrous-oxide-Oxygen.—The many advantages of this anaesthetic mixture have been described by Langton Hewer in papers written in 1930 and 1935. Hewer still regards nitrous-oxide-oxygen as the best anaesthetic in thoracic cases. Briefly, its advantages are that it is unirritating and non-toxic and is speedily eliminated, leaving very little in the way of after-effects. Nausea and vomiting are reduced to a minimum. It does, however, present certain disadvantages. The chief is that the amount of oxygen it is possible to give with the nitrous oxide is comparatively small, varying from 15 to 20 per cent. It is often difficult to keep a steady anaesthesia without adding small quantities of chloroform or divinyl ether. Then, again, muscular relaxation is very poor.

Ethylene and Acetylene.—Ethylene is quite a useful agent, but is employed very little in this country because of its explosive properties. Incidentally, its unpleasant odour adds to its unpopularity. Acetylene is used on the Continent, but I have had no experience with it.

Cyclopropane is the latest member of the group of inhalational anaesthetics, and has proved itself invaluable in thoracic surgery. The advantages are as follows:

- (a) It is given with as high a proportion of oxygen as 80 to 85 per cent. This, in cases where there is extensive disease in the lungs with resultant low vital capacity, solves a great problem.
- (b) Respiratory movement is reduced to a minimum.
- (c) In spite of quiet anaesthesia a cough reflex is easily maintained.
- (d) It is rapidly excreted—almost as rapidly as nitrous-oxide-oxygen.
- (e) The induction, even in cases where no barbiturate is used, is rapid and pleasant, because of the high percentage of oxygen and because of the fact that the odour is not objectionable.

(f) Cyclopropane, during extensive experiments carried out by Waters (Waters *et al.*, 1934) of Madison, U.S.A., proved to be absolutely non-toxic to the liver and kidneys.

(g) It is absolutely non-irritating to the membranes lining the air passages.

Thus it will be seen that in spite of a few disadvantages cyclopropane very nearly approaches the ideal inhalational anaesthetic.

An Efficient Machine

The machines used in thoracic surgery anaesthesia are numerous and complicated, but the essential features are:

1. The machine should be fitted to administer nitrous oxide and oxygen with some attachment for the adding of chloroform or divinyl ether if necessary.

2. It should be of the closed-circuit type, with a carbon-dioxide-absorbing apparatus. This is, of course, essential with cyclopropane; and in any other form of anaesthesia it keeps the respirations smooth and gentle, helping to maintain throughout that tranquillity which helps so much in intrapleural procedures.

Choice of Method and Agent

The final points to be considered are the best method of anaesthesia and the best agents to be used in various operations:

Operations requiring Novocain.—The following three operations—(a) induction of artificial pneumothorax, (b) resection of rib for empyema, (c) operations on the phrenic nerve—may be grouped together as requiring novocain 1 per cent. infiltration. A small injection of pentothal is kindly and useful in the last-mentioned operation, when the nerve is avulsed. Rib resection is carried out under local infiltration with 1 per cent. novocain, requiring at the most four skin punctures.

Thoracoplasty.—Hewer (1935) in his paper on anaesthesia in thoracic surgery states that he considers nitrous-oxide-oxygen the best anaesthetic for this operation. Over a period of eight years or more at the Brompton Hospital a routine administration of nitrous-oxide-oxygen combined with certain therapeutic measures gave excellent results, and it is doubtful if a change over to cyclopropane would make any improvement. Morphine and hyoscine, or omnopon and scopolamine, are given as premedication. Sometimes the patient is actually put to sleep with a few cubic centimetres of evipan or sodium pentothal. Anaesthesia is maintained with the highest possible percentage of oxygen. Careful watch is kept on the blood pressure and pulse pressure. When the rib resection is actually started 1½ litres of tap water with glucose are run into the rectum. In these cases the added tranquillity and peaceful anaesthesia obtained with cyclopropane do not quite compensate for the increased bleeding and the lengthening of the operation time.

Apicolysis and Extrapleural Pneumothorax.—These are of a similar type of operation, and the essential feature is that quietness of respiration and the minimum of movement must be aimed at when the separation of the pleura is being carried out. In these cases cyclopropane has proved to be vastly superior to nitrous-oxide-oxygen. Extra bleeding with the former is a disadvantage, as is also the fact that the use of diathermy for haemostasis is contraindicated, both of which facts go to increase the length of the operation time. But these disadvantages are outweighed by the smoothness of the anaesthesia obtained.

Thoracotomy.—This group of operations includes the removal of one lobe or the whole of the lung for abscess, bronchiectasis, and carcinoma, and also operations for mediastinal tumours, etc. Where there is very little or no sputum, and where the patient's psychological condition permits, one or other of the techniques described for the production of spinal analgesia is excellent. Provided an artificial pneumothorax has been induced successfully on the affected side, respirations are not impaired and the patient is not at all embarrassed when the pleural cavity is laid open. If, however, postural drainage has not been entirely successful and there is much sputum, then measures must be taken to deal with the accumulation of material in the trachea and bronchi. In the less serious cases anaesthesia may be conducted just using an ordinary face-piece and allowing the patient to cough up the sputum or blood. This is easily carried out with nitrous-oxide-oxygen or light cyclopropane. If, however, the sputum is copious, tracheal or bronchial intubation must be resorted to. The simplest method is to intubate the patient with an ordinary Magill tube by the nasal route. Then, when quiet anaesthesia has been established, a gum-elastic catheter is passed into the trachea beside the Magill tube and attached to a sucker. Continuous or intermittent suction can then be applied. This may tend to upset the anaesthesia in closed-circuit administration, so that it is probably best to use a machine of the intermittent-flow type such as the McKesson, with which machine the pressure can be maintained without altering the proportions of the mixture. The pharynx should be packed off with gauze soaked in liquid paraffin. Intra-bronchial intubation is more difficult technically. First of all an extensive cocaineization of the pharynx and larynx should be carried out. This will be described later in dealing with bronchoscopy. A special introducer designed by Magill (1936b) carries a tube fitted with an inflatable rubber cuff into the bronchus on the sound side. The cuff is inflated, shutting off the bronchus of the affected side and the lower part of the trachea. This gives very smooth anaesthesia, but in a certain number of cases of lobectomy collapse of the remaining lobe ensued, which may have been due to the prolonged closure of the bronchus of the affected side. Anaesthesia may then be carried out, using either nitrous-oxide-oxygen or cyclopropane. Rovenstine (1935) of New York, and all the anaesthetists at the Lahey Clinic, Boston, use cyclopropane exclusively in their cases, chiefly on account of the advantages enumerated above. They have written papers on this subject, which have been published from time to time in the *Current Researches in Anesthesia and Analgesia*.

Bronchoscopy.—This is always carried out under local anaesthesia at the Brompton Hospital. The nose and pharynx are sprayed with 20 per cent. cocaine, and finally the pharynx and pyriform fossae are swabbed with the solution, allowing the passage of an ordinary Magill's laryngoscope in order to spray the larynx and trachea. With the use of omnopon and scopolamine beforehand the operation is usually performed quite smoothly. In nervous patients pentothal may be injected.

Conclusion

I have touched but briefly on this extensive subject, for a whole paper might be written about one individual operation and one individual method. Results all tend to show that improvements in anaesthesia technique have done great things to help the thoracic surgeon to carry out the formidable operative procedures of the present day.

REFERENCES

- Etherington-Wilson, W. (1934). *Brit. J. Anaesth.*, **11**, 43.
 Harris, T. A. B., and Rink, E. H. (1937). *Guy's Hosp. Rep.*, **87**, 127.
 Hewer, C. Langton (1930). *Proc. roy. Soc. Med.*, **23**, 771.
 — (1935). *Post-grad. med. J.*, **11**, 11.
 — (1936). *Proc. roy. Soc. Med.*, **29**, 652.
 Jones, W. Howard (1930). *Anesth. & Analges.*, **9**, 218.
 Magill, I. W. (1936a). *Proc. roy. Soc. Med.*, **29**, 643.
 — (1936b). *Amer. J. Surg.*, n.s., **36**, 450.
 Rovenstine, E. A. (1935). *Anesth. & Analges.*, **14**, 270.
 Shields, H. J. (1935). *Curr. Res. Anesth. Analges.*, **14**, 193.
 Waters, R. M., et al. (1934). *Anesth. & Analges.*, **13**, 56.

INGUINAL LYMPH GLAND BIOPSY IN THE DIAGNOSIS OF TUBERCULOUS DISEASE OF THE KNEE

BY

H. J. SEDDON, F.R.C.S.*Resident Surgeon, Royal National Orthopaedic Hospital,
Stanmore, Middlesex*

(WITH SPECIAL PLATE)

One of the gratifying features of the practice of orthopaedic surgery to-day, at any rate in the London area, is that most patients with tuberculous joint infection are sent to us for treatment at a very early stage. I take this opportunity of expressing my admiration for the vigilance of those general practitioners, tuberculosis officers, and school medical officers to whom this state of affairs is largely due. The result is that the well-known clinical picture of tuberculous joint disease is now rarely seen, and one is often faced with patients who complain of little more than a slight limp and some joint pain, and show the minimum of signs, either clinical or radiographic. The finesse of diagnosis has increased immensely; now one often sees cases so early that it is at first impossible to say with any certainty what form of joint infection is present. Of no joint is this more true than the knee.

I do not propose to discuss the signs that are of value in the diagnosis of tuberculous disease of the knee, but I am sure we agree that there are many cases in which all the available evidence (short of biopsy) is insufficient to justify even a provisional diagnosis. Certain surgeons are content to await events, trusting that some sign will ultimately appear to dispel the uncertainty. This often means waiting many months; and sometimes, as with others who once sought after a sign, "there shall no sign be given." If the patient is a child, treatment is instituted on the assumption that the joint is tuberculous—which retards or prevents the development of characteristic features. This is an unsatisfactory state of affairs, leading to waste of time and hospital beds, unnecessary confinement of the patient, and needless apprehension in the minds of all concerned. That is why exploration of the joint has become increasingly frequent. But this involves opening the largest joint of the body, and perhaps a prolonged search for an obscure juxta-articular focus. Although I have not encountered any unfortunate sequelae, such as sinus formation, in any of the joints that I have explored, I would hesitate to recommend the operation. Minor disasters sometimes occur: I have seen the results of them.

Two and a half years ago my friend F. P. Fitzgerald suggested that examination of an inguinal lymph gland on the affected side might give us valuable information, and this was when our investigation began. At that time the pioneer work of José Valls (1933) of Buenos Aires was unknown to us. To him is due the credit for describing this aid to diagnosis.

Forms of Tuberculous Infection of the Inguinal Lymph Glands

Tuberculosis of the inguinal lymph glands is known to occur under the following conditions: (1) As part of a generalized tuberculous adenitis—a rare condition (Battigelli, 1935). (2) As part of a generalized tuberculous adenitis associated with disease in the lungs—a rare condition (Benda, 1936). (3) As part of a generalized tuberculous adenitis associated with multiple bone foci—again a rare condition (Jungling, 1920; Jacobsen, 1936). (4) Secondary to tuberculous infection of the anal region or vulva. (5) Secondary to tuberculous infection of some part of the lower extremity: (a) skin (Carter and Smith, 1935); (b) tendon sheaths of the foot (one personal case—unpublished); (c) tarsal bones with infection of the soft parts (an unpublished case under the care of my colleague Mr. R. Y. Paton). The point I wish to establish is that there is no evidence that tuberculous inguinal adenitis occurs as an apparently isolated lesion. If it does it must be exceedingly rare. Therefore if it can be shown that a patient with arthritis of the knee occurring as an apparently isolated lesion also has tuberculous foci in the inguinal lymph glands on the same side, it is reasonable to presume that the lesion in the joint is of the same nature.

Although the existence of subsynovial lymphatic vessels in diarthrodial joints has long been known, the lymphatic drainage of human joints has still to be investigated. J. A. Key (1926, 1932) has carried out valuable work on the fate of indian-ink particles injected into the knee-joints of adult rabbits. A small proportion of the particles were carried to the popliteal lymph glands, but none to the inguinal.

It is curious that we have been so slow in arriving at the diagnostic possibilities of examination of lymph glands draining a suspected joint; for palpable enlargement of the external iliac glands has long been recognized, particularly in France, as a feature of tuberculous disease of the hip.

Technique

For removal of the gland local anaesthesia is employed, except in small children. If the glands are palpable, size and inferiority of position should guide the surgeon in his choice: a large gland situated low in the femoral triangle is promising. If no gland is palpable, the incision should be made about one inch below the site of the saphenous opening. Two glands may be removed if easily accessible, but it must be remembered that the operation is a biopsy and nothing more. A fragment of gland about 10 × 5 × 5 mm. is set aside for bacteriological examination (guinea-pig inoculation or culture), and the remainder is examined histologically.

Results

The results in the eighteen cases of tuberculous disease in which our information is now sufficiently complete are set out in the table. In a few of these cases inguinal biopsy was not required to establish the diagnosis, since material from the knee was available from the first; how-