BRITISH MEDICAL JOURNAL

LONDON SATURDAY JANUARY 21 1939

THORACOLYSIS

A CONSERVATIVE AND SELECTIVE OPERATION FOR THE TREATMENT OF CERTAIN CASES OF PULMONARY TUBERCULOSIS

BY

LAURENCE O'SHAUGHNESSY, M.D., B.S., F.R.C.S.

London

AND

GEORGE MASON, M.B., B.S., F.R.C.S.

Newcastle-upon-Tyne

(WITH SPECIAL PLATE)

The importance of obtaining selective effects with the various forms of collapse therapy now used in the treatment of pulmonary tuberculosis is becoming increasingly apparent. The reasons for this include the necessity for performing minimal interventions on patients in poor condition and the desirability of avoiding the excessive sacrifice of healthy pulmonary tissue. Such an object is, of course, aimed at every day in the employment of artificial pneumothorax and even in such radical measures as the apicolysis of Semb. There are, however, cases with cavity systems limited to the middle or lower zone of one lung in which an artificial pneumothorax is unattainable and where a standard total thoracoplasty seems needlessly drastic both from the point of view of the severity of the operation and the curtailment of lung function; for these, the development of some conservative and selective, yet efficient, operation is desirable.

Developments in Thoracoplasty

In the early days of thoracoplasty favourable results were occasionally obtained with modified rib resections carried out in different parts of the chest wall; but the lack of radiological guidance and the consequent "blind" performance of these measures made them, in the majority of cases, unsatisfactory. Subsequently the standardized paravertebral extrapleural thoracoplasty of Sauerbruch became, and remains, the operation of choice in the treatment of carefully selected unilateral cases of known chronicity. Its application both to less extensive and to more recent lesions was also frequently successful, but as the original conditions for operation were relaxed the incidence of disappointing results increased.

The search for this maximal collapse effect led to the development of paravertebral thoracoplasty and its modern modifications, but meanwhile the therapeutic possibilities of less radical rib resections elsewhere were neglected. However, the evolution of apical thoracoplasty, and its later modifications such as that of Semb, demonstrated the possibility of providing a degree of collapse sufficient to permit of local healing and yet of conserving the function of a maximum amount of healthy lung tissue.

A rib being a curved structure, concave inwards, the removal of even a small portion diminishes the capacity of the thorax, especially in the vicinity. Scar tissue within the lung, adherent to the parietes, subjacent to such a costectomy, is consequently able to retract. Monaldi in 1934 described his favourable experiences with antero-lateral thoracoplasty, and André Maurer (Maurer, Tobé, and Davy, 1935; Maurer and Rautureau, 1936) elaborated this idea still further with an account of his work on "La Thoracéctomie Élastique de Détente et d'Attente."

Maurer emphasized the local selective nature of these minimal interventions. Whether any localized rest factor occurred which might initiate "productive" response in a hitherto "exudative" lesion is not known. Monaldi, however, interpreted the absence of breath sounds at the site of operation as evidence of local rest. We consider the time opportune to recall this work and to add our own observations on a small series of similar cases in which it became obvious that healing of the lesions was improbable without some form of collapse therapy. It was considered desirable, after consultation with the physicians responsible for their care, to attempt to promote healing by some more selective and less radical operation than by one of the orthodox forms of thoracoplasty.

We have been concerned with that group of cases in which the disease—usually of a fibro-cavernous type—has mainly involved the middle or lower zone of one lung, and in which it has been impossible to obtain an adequate collapse by artificial pneumothorax, and where healing has not followed diaphragmatic paralysis. For this type of case the alternative to the operation described below was a "total" thoracoplasty with the consequent functional sacrifice of an unduly large area of healthy lung tissue. This was considered an operation of undesirable severity in view of the general condition of the patients.

Treatment by Thoracolysis

Healing was initiated by carefully designed selective resections of relatively small portions of ribs in the vicinity of the lesions—that is, just where relaxation of scar tissue in the underlying lung was required. Thoraco-

lysis was the term originally applied to the operation whereby removal of the precordial ribs, etc., made that part of the chest wall more yielding. We have used the term for the procedure under discussion as it consists essentially in making the chest wall more yielding in some desired situation and so permitting retraction of any scar tissue in the underlying lung. The use of the word has the additional advantage of avoiding confusion between this conservative procedure and the more radical type of operation commonly known as thoracoplasty.

The portions of ribs which it is usually necessary to remove in this operation are easily accessible through direct incisions; little retraction being necessary, tissue trauma is reduced to a minimum. Consequently the operation is one which can be employed for patients in relatively poor general condition. For similar reasons it is a comparatively easy operation to carry out under local anaesthesia, should this be desirable, although for our cases we have not done so. Comparative freedom from marked post-operative disturbances characterized the cases in the series.

Whether or not it is an operation which might be used in the preparation of patients for more serious interventions such as a major thoracoplasty is a matter which requires further experience; but it is obvious that even if it fails to achieve a cure in any particular case there is nothing to preclude a subsequent thoracoplasty of the standard type. It also seems likely that this procedure could be applied with some likelihood of success to those cases where there is a previous contralateral thoracoplasty. It is conceivable that its application to certain apical lesions might prove beneficial. It also seems likely that it might be employed as a bilateral procedure. Accurate radiological determination of the type, extent, and localization of the disease is an essential preliminary to the employment of this method. In addition to the ordinary antero-posterior views, lateral, tomographic, and stereoscopic views are essential. Of these we have found tomographic and lateral the most important.

Rib regeneration after resection occurs as a rule so rapidly that this operation would have little chance of success were it not for the means which are now available to retard it—cauterization of the periosteum with Zenker's solution, with 10 per cent. formol, or by diathermy. We prefer to use strips of gauze soaked in Zenker's solution, placing them in the periosteal bed, from which the rib has been removed, for at least four minutes. In this connexion we are opposed to ablation of the periosteum: ultimately restoration of bony continuity in the chest wall is desirable on mechanical grounds. For the same reason—so that localized paradoxical movement may be prevented, and so that regeneration of the ribs may occur with the greatest possible diminution of their internal concavity—some form of pad may be applied direct over the area concerned, until such time as regeneration of the ribs is completed.

If there is clinical and radiological evidence that the operation has favourably influenced the lesion but has not healed it by the time the ribs have regenerated, the operation—which itself may have been done in stages adapted to the condition of the patient—may be repeated, the chest wall being made yielding once again, so that the underlying scar tissue may further retract.

The clinical and radiological picture after operation was that of progressive improvement, such as is seen with cases in which other forms of collapse therapy have been successfully employed, the pulmonary lesions—as dis-

cernible by radiographs—disappearing as the general condition improved. As this improvement took place the sputum diminished in quantity, becoming free from tubercle bacilli and ultimately ceasing entirely.

Case I

A man aged 21 was admitted to the sanatorium in August, 1935, having suffered from pulmonary tuberculosis for three years following an attack of influenza. He had twice had periods of ordinary sanatorium treatment elsewhere—one for eighteen months and again for six months—without material improvement. On two occasions there had been a haemoptysis of 3 to 4 oz.

On admission the general condition was fair, the weight being 9 st. 12 lb. and the blood sedimentation rate 17. Tubercle bacilli were present in the sputum. X-ray examination of the chest revealed slight, probably quiescent, infiltration at the right apex, and on the left side a considerable infiltration and mottling of the infraclavicular and middle zones. Artificial pneumothorax could not be induced. The left phrenic nerve was effectively crushed, but despite this and three months' "bed-rest" the weight remained stationary, and the sputum (2 oz. a day) remained positive. Moist sounds were still audible in the left lung. Tomographic examination revealed the presence of a series of cavities in the posterolateral part of the middle zone of the left lung (Fig. 1, Special Plate). Although the temperature was normal, the pulse rate—absolute rest—ranged between 88 and 104. The blood sedimentation rate still remained low—14—and the vital capacity was 2,860 c.cm.

Thoracolysis was advised, and in May, 1936, under general anaesthesia (oxygen and ether), posterior portions of ribs four to eight were removed from the left side, the periosteum being cauterized with Zenker's solution. An uncomplicated convalescence was marked by little appreciable post-operative disturbance. Subsequent tomograms showed a progressive fibrosis and pleural thickening with contraction of the cavities, while the sputum diminished to half a drachm daily, but remained positive. Accordingly, in October, 1936, the diaphragm having become reactive, the left phrenic nerve was again crushed. In November, 1936, under general anaesthesia (oxygen and ether), the regenerated portions of the fourth, fifth, sixth, and seventh left ribs were again removed and the periosteum cauterized with Zenker's solution.

Convalescence was again smooth, and three months later the sputum became negative not only to ordinary examination but also on culture and guinea-pig inoculations. Tomographic (Fig. 2) and radiographic examination (Fig. 3) in February, 1938, indicated that healing had occurred with a minimal loss of functioning lung tissue. This satisfactory progress was maintained, and in September, 1938, the general condition was excellent, the weight 12 st., the B.S.R. normal, and the vital capacity 2,300 c.cm. The patient was almost free from sputum, which was persistently negative. He was able to work regularly for four and a half hours daily in the wood-graining department of the settlement.

Case II

This patient, a miner aged 32, was admitted to the sanatorium in September, 1936. He had suffered from cough and sputum for five years; haemoptysis had occurred four years previously. He had been discharged from the sanatorium only three months previously, in an apparently quiescent state and with negative sputum, following a nine-months period of routine sanatorium treatment.

On admission the general condition was poor; the weight 10 st. 7 lb.; tubercle bacilli were present in the sputum; there was thickening of both vocal cords; the pulse rate ranged between 70 and 90 per minute, and the temperature between 97° and 98.6° F. The x-ray examination showed an extensive infiltration of the lower part of the right lung, in which there was a fair-sized cavity (Fig. 8).

The right phrenic nerve was crushed on November 17, 1936, and a month later, under general anaesthesia (oxygen and ether), the postero-lateral segments of ribs five to nine on the right side were removed, and the periosteum cauterized with Zenker's solution. There was no post-operative disturbance, and he continued to improve with routine bed rest and sanatorium treatment until July, 1937, when he was discharged. At that time, although he had a trace of sputum, no tubercle bacilli could be found in it, his laryngitis was quiescent, the temperature range normal, and an x-ray examination indicated that the lung condition had apparently healed. Since December, 1937, he has worked in the sectional building workshop of the settlement, and apart from an occasional cough he has remained free from symptoms and has gained 12 lb. in weight. In July, 1938, an x-ray examination showed that no relapse had occurred (Fig. 9).

Case III

A man aged 20, a factory hand, was admitted to the sanatorium in October, 1936, with a history of pleurisy dating from the previous August. The temperature varied between 97.4° and 99.4° F. He weighed 7 st. 9 lb., and the pulse rate was between 80 and 100. Although there was only a trace of sputum, it contained tubercle bacilli.

X-ray examination showed diffuse infiltration of the lower and middle zones of the left lung, with a cavity in the vicinity of the hilus (Fig. 13). The phrenic nerve was crushed on December 12, 1936, but although the patient was kept completely at rest the lesion had not healed four months later. Accordingly, on April 6, 1937, under general anaesthesia (oxygen and ether), postero-lateral portions of ribs four to six were removed in the region of the cavity, and the periosteum was cauterized with Zenker's solution. There was no post-operative disturbance, and an uneventful convalescence resulted, the patient being discharged in February, 1938. At that time the temperature and pulse rate were normal, the B.S.R. was 4.5, the weight 8 st. 10½ lb., and the sputum negative on direct examination. The cavity is no longer visible in the radiograph (Fig. 14), and the patient, whose health is well maintained, has been working for three months as a machinist in a hosiery factory.

Case IV

A girl aged 18 was admitted to the sanatorium in July, 1936, after an attack of haemoptysis. Three years previously she had suffered from pneumonia. The sputum contained tubercle bacilli. X-ray examination showed diffuse infiltration in the lower zone of the right lung, but no definite cavity. The general condition improved on routine sanatorium treatment and she was discharged with a negative sputum in January, 1937; but in April of that year, following a cold, relapse occurred. At this stage there was radiological evidence of a diffuse infiltration in the lower zone of the right lung with a small cavity. A pneumothorax could not be induced. The right phrenic nerve was crushed in April, 1937, but the cavity persisted and thoracotomy was decided upon.

Accordingly, on October 10, 1937, segments of the sixth, seventh, and eighth ribs were removed and the periosteum cauterized with Zenker's fluid. There was some pyrexia following operation, and on two occasions fluid was aspirated from the right pleura; it was not until December 19, 1937, that the temperature and pulse returned to normal. The patient has improved since operation, gaining about 2 st. in weight, and is working as a sanatorium orderly. The sputum is negative.

Case V

A woman aged 31 was admitted to the sanatorium on November 3, 1936, with a history of cough and sputum dating from the previous August. Neither temperature nor pulse rate was elevated. The sputum contained tubercle bacilli. Despite routine sanatorium treatment the condition did not greatly improve, the sputum remaining positive. X-ray

examination showed diffuse infiltration in the middle zone of the left lung, and with a lateral view a small cavity was seen near the hilus.

The phrenic nerve was crushed in August, 1937, but the sputum remained positive. It was decided to perform a thoracotomy. Accordingly, on November 23, 1937, under general anaesthesia (oxygen and ether), segments of the sixth and seventh ribs were removed. Unfortunately the lung was not completely adherent in the area chosen for resection, and during the operation a free pleural space was entered; hence this was a case where thoracotomy should probably not have been employed. There was no immediate distress, but a large effusion rapidly formed, and with the appearance of a secondary pyogenic infection of the effusion the patient became gravely ill. For a time the empyema was treated by aspiration, but on January 1, 1938, a rib resection and drainage had to be done. Subsequently a paravertebral thoracoplasty (ribs one to nine) was carried out, and the patient is in good general condition. Although a negative sputum has been repeatedly recorded since operation, on a recent occasion (October, 1938) it was positive.

Case VI

This patient, a boy aged 16, is a member of a family with several sufferers from open tuberculosis. Tuberculous glands were excised at the age of 9, and six years later he had been ill with dry pleurisy. A year before admission, occasional "streaking" of the sputum having occurred, he was taken into an institution, but failed to improve, and was transferred to a sanatorium.

Ordinary radiographic and tomographic examination revealed extensive infiltrative disease with cavitation of the lower middle zone of the right lung (Figs. 4 and 5). Artificial pneumothorax could not be induced as the right phrenic nerve was crushed. No material improvement occurred with sanatorium treatment, and thoracotomy was decided upon. At that time, five months later, not only was his condition poor, his temperature between 99° and 100° F. each evening, and his pulse rate correspondingly elevated, but his weight was only 7 st. 9 lb. About ½ oz. of greenish pus was coughed up daily. The vital capacity was only 2,100 c.cm., but the B.S.R. had improved from 23 to 12. Tomography had localized the cavity as being close to the ninth right costotransverse articulation. Accordingly, under local anaesthesia—supplemented by intravenous evipan at the moment of cutting the bone—two and a half inches of the ninth rib and one inch of the eighth rib were resected periosteally, and Zenker's solution applied for five minutes to the periosteum.

Convalescence was uneventful. The phrenic nerve was again crushed three weeks after the thoracotomy, and the patient has been kept at rest since. Marked improvement in the general condition has occurred, the pulse and temperature having become normal, and the sputum having almost disappeared and having become negative both to culture and to guinea-pig inoculation. The weight has increased by 28 lb., the B.S.R. is 7, and the vital capacity has actually increased to 2,800 c.cm. Radiographs and tomographs six months after operation demonstrated that healing had occurred (Figs. 6 and 7).

Case VII

This patient was a motor mechanic aged 29. Five years previous to admission to the sanatorium in November, 1937, he had pneumonia and pleurisy, but made a good recovery and remained well for three years. At the end of that time he developed a cough that has persisted, accompanied by a small amount of sputum. He was in another institution for a year. During that time, as an artificial pneumothorax could not be induced, the left phrenic nerve was avulsed (May, 1936). Haemoptysis had occurred on one occasion, about a year previously.

The general condition was fair; temperature and pulse rate were not elevated. The weight, which on admission was 9 st. 4½ lb., fell slightly in five months to 8 st. 10 lb. About one ounce of sputum containing tubercle bacilli was

expectorated daily. X-ray examination revealed cavitation in the left lower lobe—near its upper part posteriorly. Bed rest and general treatment failed to produce any improvement; indeed, the x-ray appearances after three months were, if anything, rather worse (Figs. 10 and 11). Accordingly thoracotomy was advised. On March 31, 1938, under general anaesthesia (intratracheal oxygen and chloroform), the posterior ends of ribs six to ten were resected subperiosteally and their periosteal beds cauterized with Zenker's fluid. The immediate post-operative period was uneventful.

It was recorded in the notes on May 12, 1938, that since operation the patient had done amazingly well: the sputum had decreased from one ounce to three spots daily, and on that day he had had none for the first time for years. X-ray examination showed the cavity diminishing in size. A few bacilli were found in one specimen of sputum about the middle of September, after it had been repeatedly negative for about two months.

On October 31, 1938, the general condition was excellent; a slight but steady increase in weight having taken place since August, the weight now was 10 st. 7 lb., and the sputum was so scanty that a specimen obtained with the utmost difficulty was negative. The x-ray appearances are correspondingly improved and suggest that healing has occurred (Fig. 12). The patient is now being allowed up, and sputum is negative on culture and animal inoculation.

Case VIII

An upholsterer aged 25 was admitted to the sanatorium on November 16, 1937, nine weeks after the first onset of his illness, which was regarded as a bronchopneumonia of the right base.

Slight improvement followed routine sanatorium treatment, but in February, 1938, a large basal cavity was seen in the x-ray picture of the right lung. Accordingly an artificial pneumothorax was induced, but as this was inadequate it was supplemented on April 6, 1938, by crushing of the right phrenic nerve, and the cavity became a little smaller. On June 30, 1938, there was still a little sputum containing tubercle bacilli, and an occasional slight elevation of temperature to 98.8° F. The pulse ranged between 75 and 100 per minute and the weight was 9 st. 10 lb. The cavity appeared in the x-ray picture to have increased (Figs. 15 and 16). Accordingly thoracotomy was advised. On July 7, 1938, under general anaesthesia (intratracheal oxygen and chloroform), posterior portions of ribs six to eleven were resected subperiosteally and their periosteum was cauterized with Zenker's fluid. The immediate post-operative period was marked by an elevation of the temperature, starting some days after the operation and persisting for a few days until the contents of a large haematoma were expelled from the wound during a fit of coughing.

The general condition steadily improved; the sputum, becoming negative, ceased, and x-ray examinations showed the cavity to be diminishing (Fig. 17) and then finally to disappear (Fig. 18). Although there was a slight loss of weight immediately after operation, three months later the patient was more than a stone heavier than immediately before it, and he was being allowed up for about four hours daily. The sputum is negative on culture and animal inoculation.

Summary

Attention is drawn to the possibility of treating successfully certain cases of pulmonary tuberculosis, especially those in which a middle or lower zone of one lung is involved by fibro-cavernous disease, in which an adequate artificial pneumothorax cannot be established, and in which phrenic paralysis fails to effect healing.

The underlying principles of the method and the possibility of extending its scope are referred to, and a series of eight cases in which it was employed with success are recorded.

It is possible, with the operation described and in suitable cases, to obtain healing with the minimum diminution in the functional capacity of the chest and often even with an active diaphragm.

We wish to acknowledge our indebtedness to our colleagues, the medical superintendents and their assistants at the sanatoria—Preston Hall, Ransom, and Wooley—where these cases were treated, for their co-operation in the selection and post-operative management of these, as well as for their interest in the problems involved and help in providing access to their records. We are similarly grateful to Dr. Philip Ayre and Dr. J. K. Hasler for their skilled administration of the anaesthetics required in these cases.

REFERENCES

- Maurer, A., Tobé, F., and Davy, P. E. (1935). *Rev. Tuberc.*, 5 ser., 1, 904.
— and Rautureau, R. (1936). *Ibid.*, 5 ser., 2, 642.
Monaldi, V. (1934). *Atti mem. Soc. Lombardia Chir.*, 11, Seduta del 25 Novembre.

CRITICAL REVIEW OF EXTRAPLEURAL PNEUMOTHORAX

BY

R. NISSEN, M.D.

Professor of Surgery in the University of Istanbul

In 1929—at first without knowledge of the case treated by Meyer fifteen years previously—I used extrapleural pneumothorax for a considerable number of patients, and have since reported thereon (Nissen, 1931, 1932, 1933, and 1937). Numerous publications during the last two years, personal communications from surgeons and tuberculosis doctors, and my own observations while visiting clinics and sanatoria show that extrapleural pneumothorax has become a very frequently used method. I was the first to reintroduce this treatment, and now, after nine years' experience of it, think that a stand should be made against the present tendency to extend the indications for its use. I will not go into these in detail now, but will first discuss the disadvantages and complications of extrapleural pneumothorax. The operation is an adventure into the unknown—to a lesser extent the same objection may be raised against the classical artificial pneumothorax, and unexpected complications may disturb the placid course of the medical treatment of phthisis—but, as will be shown, in extrapleural pneumothorax the mechanical conditions created at operation are immediately subject to modifications, for the most part beyond our control.

Its Disadvantages and Complications

EMPHYSEMA OF THE MEDIASTINUM

One possible complication is that after completion of the operation air escapes as a result of bouts of coughing, causing general emphysema and threatening to involve the mediastinum. In one of my cases a very serious emphysema of the mediastinum occurred the night after the operation, and I believe that only incision into the jugular notch prevented a fatal outcome.

DIFFICULTY IN REFILLING

When air has escaped from the extrapleural pneumothorax it may be difficult to find the right tissue space again in attempting to refill. It is true that the exudate usually gives the right clue. But everyone with experience of artificial pneumothorax refills realizes how awkward it is to introduce air without knowing just where one is. There is no manometric indication, for after extrapleural pneumothorax the pressure oscillates round about zero.

LAURENCE O'SHAUGHNESSY AND GEORGE MASON: THORACOLYSIS

FIG. 1.—Case I. Tomogram showing large cavity system in left middle zone.

FIG. 2.—Case I. Tomogram showing healed lung fifteen months after thoracolysis.

FIG. 3.—Case I. Radiograph showing minimal diminution in capacity of hemithorax after thoracolysis.

FIG. 4.—Case VI. Cavity and infiltration in lower part of right lung.

FIG. 5.—Case VI. Tomogram before thoracolysis.

FIG. 6.—Case VI. Tomogram six months after thoracolysis, showing disappearance of cavity.

FIG. 7.—Case VI. Radiograph showing healing six months after thoracolysis; very slight diminution in capacity of hemithorax.

FIG. 8.—Case II. Cavity and infiltration in lower part of right lung.

FIG. 9.—Case II. Soundly healed, nineteen months after thoracolysis.

LAURENCE O'SHAUGHNESSY AND GEORGE MASON: THORACOLYSIS

FIG. 10.—Case VII. Cavity system in lower middle zone of left lung.

FIG. 11.—Case VII. Oblique view indicating left paravertebral localization of disease.

FIG. 12.—Case VII. Soundly healed, eight months after thoracotomy, without marked diminution in capacity of hemithorax.

FIG. 13.—Case III. Cavity system near hilus of left lung.

FIG. 14.—Case III. Healed two and a half months after thoracotomy with little diminution in capacity of hemithorax.

FIG. 15.—Case VIII. Large cavity localized in the right paravertebral region before thoracotomy.

FIG. 16.—Case VIII. A.-P. view showing cavity before thoracotomy (unsuccessful artificial pneumothorax still present).

FIG. 17.—Case VIII. Cavity healing one month after thoracotomy.

FIG. 18.—Case VIII. Cavity completely healed three months after thoracotomy, with minimal diminution in capacity of hemithorax.