further relatives or friends are usually available without much delay. It is a great deal more satisfactory from every point of view to secure a donor from the family circle than from a transfusion service. Unfortunately, however, it is not always possible to do this, and an emergency list of donors should therefore be at hand. As a final resort, in case of extreme emergency the doctor himself may be a suitable donor; but how many medical men are aware of their own blood group?

Conclusion

It is perhaps advisable again to make it clear that blood transfusion is not being advocated as necessarily the method of choice in the treatment of pregnancy and gynaecological anaemia. It is advocated, however, for the severe haemorrhages which occur from time to time in every clinic and many private practices. Severe antepartum and post-partum haemorrhage, the extreme collapse due to extensive internal bleeding in certain ectopic gestations, the exsanguination that is at times associated with abortion or hydatidiform mole, and the collapse in severe haemorrhage of the newborn are all examples of those accidents which at the present time account for over 300 lives each year in England and Wales alone. It is for this type of haemorrhage that the wide adoption of emergency transfusion is urged as the most effective method of saving life.

REFERENCE

Solomons, B. (1938). *J. Obstet. Gynaec. Brit. Emp.*, **45**, 644.

THE NATURE OF GROWING PAINS AND THEIR RELATION TO RHEUMATISM IN CHILDREN AND ADOLESCENTS

BY

J. C. HAWKSLEY, Ph.D., M.D., M.R.C.P.

Assistant Physician to University College Hospital

While the phrase "growing pains" is apt to provoke the comment that there is no evidence that the process of growth is painful, it is still widely used as a diagnosis for cases of pain in the limbs of growing children and adolescents. Its use and abuse have often provoked controversy because of vagueness, particularly as regards the aetiology of the pains. A definition of what is meant clinically by growing pains must vary with the observer's own individual taste, but in most cases it would probably be found to indicate pain, sometimes recurrent, felt in the limbs or back by a growing person in whom there was no evidence of acute or subacute rheumatism, trauma, or local infection. The condition thus defined is common throughout school age, and may be seen for several further years beyond this: growth may be directly responsible for skeletal pain in a number of cases by causing an increased strain through weight-bearing on some postural deformity such as flat-foot, knock-knee, or scoliosis.

For many years it has been felt that growing pains were perhaps to be regarded as direct manifestations of rheumatism, and therefore of a disease capable of provoking carditis; but again there has been no uniformity of medical opinion. There are those who consider the two conditions to be quite unrelated, and also those who regard cases thus diagnosed as potential cases of rheumatic carditis, to be treated with periods of rest, absence from school, and curtailment of physical exercise. This state of affairs is very unsatisfactory, since it is clear that

if there is in fact a danger of carditis insufficient care is often being exercised, while if there is not, a large number of young people are having their activities curtailed and scholastic careers impaired by prolonged periods of treatment by rest, and are being educated as partial invalids to no purpose.

A series of approximately 1,000 children and adolescents has been investigated in London and Birmingham over the last ten years (Hawksley, 1938), the object being to ascertain whether or not rheumatism and growing pains are related. The conclusions reached were that growing pains are not symptoms of rheumatism and are not associated with or followed by carditis; that growing pains are associated with a number of clinical abnormalities which can be adequately diagnosed on clinical grounds; and that treatment directed towards the underlying cause is usually successful in stopping the pain. The supposed association between growing pains and carditis is largely dependent on the layman's tendency to talk loosely of limb pains in children as growing pains from ignorance of the symptoms of true rheumatism, and also on the tendency of physicians to accept a history of skeletal pain in a child as possible evidence of rheumatism without adequately ascertaining the site and type of the pain and its association with other features in the patient's history and clinical investigation.

Sheldon (1936) collected a series of 189 cases of growing pains at the Hospital for Sick Children, Great Ormond Street. After a lapse of four years he found that only two had developed rheumatism with carditis, and none chorea. He stresses the absence of relationship between growing pains and sore throats, and also mentions the fact that there is no evidence of the histological change of acute rheumatism to be found in patients with growing pains. A number of debilitating diseases are associated with pain of the typical growing-pain type. Bray (1931) quotes a series occurring in cases of asthma in childhood.

Frequency of Occurrence of Growing Pains

During the year 1928 a series of 505 children between the ages of 4 and 14 was examined at University College Hospital and an inquiry was made into the number who had growing pains: it was found that 33.6 per cent. had suffered from them. A series of 206 children seen in 1930-1 at the Children's Hospital, Birmingham, revealed a somewhat higher percentage. Accepting the lower figure as more appropriate for London children, one child in three is at one time or another afflicted with growing pains, and therefore in all probability one child in three that develops rheumatism will also have had growing pains. Many mothers will have regarded the pain complained of as something that the child has had before and will have failed to appreciate that a new type of pain has arisen. An inquiry by the doctor at this stage would reveal differences, and a clinical examination would show that the pain was now definitely articular, that the child was ill and possibly anaemic, that there was fever and also tachycardia. The further investigation of the case in hospital would probably demonstrate changes in the blood and the electrocardiogram, indicative of active disease, that are not seen in association with growing pains. It must also be remembered that primary cardiac rheumatism is a clinical entity: Findlay (1931) found that, of 489 children in Glasgow with cardiac disease other than of congenital origin or septic nature, 8.6 per cent. had nothing in their previous history to account for it. For every three cases of primary cardiac rheumatism one will probably have had growing pains. Another feature to which the frequency of occurrence of growing pains

draws attention is that many children are suffering from pain of considerable severity which is all too often being inadequately treated. No observations have been made on the social distribution of growing pains, but the investigation of 505 children in 1928 revealed that damp housing accommodation was just twice as common in the growing-pain group as in the control group, a fact that is in agreement with Sheldon's findings concerning the effect of damp weather on the incidence of growing pains.

The Nature of Growing Pains

A series of 115 cases of growing pains was collected at the Hospital for Sick Children in 1934 for more detailed study of the clinical features presented. The cases were referred from the out-patient clinics by different physicians and clinical assistants, so that any personal factors in diagnosis were as far as possible ruled out. A number were being watched in the rheumatic clinic, and were referred from this. The following were the chief clinical findings:

The pain was predominantly muscular: it was so in ninety-eight of the 115 cases (85 per cent.); and in those who complained of joint pain this, on closer examination, was usually found to be near, rather than in, the joint. The actual sites of the pain were most commonly the front of the thighs, the calves, and behind the knees; less commonly in the backs, shoulders, arms, and instep; the groin is also sometimes affected. There was no tenderness of joints, no pain on movement, no swelling, and no redness; it was rare to find tenderness of affected muscles, except in one or two cases which were diagnosed as fibrositis. The nature of the pain is usually quite different from that of rheumatism, being of a very nagging and severe type: inquiry of children who have had both conditions reveals that the pain is subjectively quite different.

It is rare to see a child with rheumatism in tears with the joint pain, and a number of experienced ward sisters who have been questioned state that they have seldom known it to happen. In contradistinction to this no less than 45 per cent. of the 115 patients with growing pains had at one time or another been reduced to crying with the pain; many were 11 or 12 years old at the time, and it takes a lot of pain to reduce London children of that age to weeping. A feature noted by many parents was the tendency of growing pains to make their children sulky and irritable. There was a preponderance of girls in the 115 cases in the ratio of 70 to 45.

A classification of the cases into various types for the purpose of observing whether coexisting abnormalities were in any way connected with the limb pain had the following result: a frequent cause of the pain is a postural or orthopaedic defect such as flat-foot, knock-knee, scoliosis, or bad stance; treatment of the cause in these cases nearly always gives relief. During the last eighteen months a number of such cases, many of which had for a time been ineffectually treated for growing pains, have been referred to Drs. N. Kellgren and A. S. Wesson of University College Hospital for physical treatment for correction of postural defects; these patients included some who were seen in 1935 and had since then reached adolescence without losing the pains. The success of the treatment has been most satisfying, and the majority have rapidly ceased to complain of pain.

Vague ill-health is a common finding, and thirty-nine of the cases had frequent respiratory catarrh, the pains usually coinciding. Many adults complain of dull aches in the limbs when they have a cold. Successful treatment of the catarrh certainly diminishes the frequency and severity of the pain.

Fibrositis is a rather uncommon cause, and the only one in which tenderness can be demonstrated. Psychological maladjustment was unexpectedly frequent, resulting in unhappy school or home life; whether this has any relation to the frequency of pain is debatable, but Neustatter's work (1937) suggests that it may have. Tiredness, specially associated with postural defect, was common, as was also inadequate sleep, a feature noted by Seham and Hilbert (1933). Adequate rest and a holiday were usually effective in banishing the pain in these cases. Only a few cases failed to fall into one of the above groups, and it is possible that more adequate investigation would have diminished these. The common associations of the pain were postural and orthopaedic defects, vague ill-health, emotional strain, and physical and mental tiredness; not infrequently more than one factor was present. Treatment of the cause was usually successful in banishing the pain, and failure suggests inadequate diagnosis.

In contradistinction to active rheumatism—where anaemia is common and in acute cases a leucocytosis may occur—twenty-five complete blood counts failed to reveal any characteristic abnormality. The sedimentation rate was estimated in nearly all cases, and over several months in a large number. The labile nature of this test renders it unsatisfactory as a routine means of differentiation between growing pains and rheumatism. Thirty-nine cases had a raised sedimentation rate on at least one occasion, and though it would probably be safe to assume that a normal sedimentation rate indicates that active rheumatism is not present, the converse cannot by itself be held to be true—a fact that became obvious from further examination of the cases mentioned.

There is a curious tendency towards vasomotor instability in many cases: the result of pricking a finger to obtain blood resulted in vasovagal syncope severe enough to put the child on the floor in over 10 per cent. of the 115 cases seen. I have pricked hundreds of children, and never seen one faint on any other occasion. Keith (1938) found a lengthened P-R interval to be a common early change in the electrocardiograms of early cardiac rheumatism. Sir Thomas Lewis kindly investigated twenty cases with active growing pains, and found no evidence of this phenomenon.

It has been previously pointed out (Hawksley, 1931) that, whereas Hill and Allan (1929), using the colour of eyes and hair as a means of classification, found no evidence for believing in the existence of a rheumatic type of child, growing pains occur more frequently in dark brown-eyed children than in Nordics; Jewish children seem to be very prone to growing pains.

Relation of Growing Pains to Rheumatic Carditis

After a lapse of four years sixty-four unselected cases from the 115 seen in 1934 were re-examined; none showed any evidence of cardiac involvement. Adding these cases to Sheldon's, it will be seen that of 253 children followed up for four years at the Hospital for Sick Children only two have developed evidence of rheumatic heart disease, or less than 0.8 per cent. The incidence of cardiac rheumatism in 301,392 L.C.C. school children examined in 1936 was 0.8 per cent.*

Conclusions

Growing pains and rheumatism resemble each other only to the extent that both produce pain in the limbs. Careful case-taking and examination will enable a differential diagnosis to be made in nearly every case. The impor-

* L.C.C. School Medical Officer's Report, 1937, p. 41.

tance of making a correct differential diagnosis cannot be overestimated; for while it is bad to fail to treat rheumatic cases with potential carditis by rest, it is equally bad to treat children with growing pains by making invalids of them, by depriving them of their freedom, and by curtailing their instruction at school. Most cases of so-called growing pains can be treated effectively in accordance with the cause of the pain as found on clinical examination. The following clinical features serve to distinguish the two conditions.

1. The site, nature, and intensity of pain are different. Rheumatism affects joints and is usually accompanied by tenderness on pressure and movement. There may be redness, swelling, or effusion. In growing pains, muscles, ligaments, and tendons are usually the site of the pain; tenderness is rare, and there is no swelling, effusion, or redness. The pain is subjectively different in the two diseases, and is more nagging and more likely to cause crying in the non-rheumatic case.

2. In growing pains a cause of the pain is often found in postural or orthopaedic defect. Tiredness and vague ill-health are common to both diseases, but patients with growing pains are usually sulky and unsociable; this, in the author's experience, is uncommon in rheumatism.

3. Nodules do not occur except in rheumatism.

4. There is no significant change in the temperature in cases of growing pains (Seham and Hilbert, 1933). There is also no tachycardia.

5. Cases of growing pains as a rule show no anaemia; rheumatic cases frequently do.

6. Cases of growing pains have no antecedent sore throat (Sheldon, 1936).

7. Cases of growing pains show no lengthening of the P-R interval on electrocardiography.

8. Evidence is brought forward to show that the incidence of carditis subsequent to growing pains is not greater than would be expected to occur in the child population generally.

Summary

The meaning of the term "growing pains" is discussed.

The incidence of the pains in children and adolescents is surveyed.

The nature of growing pains and the possibility of their relationship to rheumatism are discussed.

The clinical features that distinguish these two conditions are recorded.

I am very grateful to the physicians of the Hospital for Sick Children for permission to examine and follow up their cases, also to Dr. Nabarro for permission to carry out the pathological tests done in 1934 in his laboratories. I am also indebted to the physicians of the Birmingham Children's Hospital for allowing me access to their cases. I have received much help from the lady almoner's department at the Hospital for Sick Children, particularly Miss Salmon, Miss Adams, and Miss Lewis. Dr. Sheldon very kindly gave me access to unpublished results, which I have quoted more fully in another paper (Hawksley, 1938). The investigations at the Hospital for Sick Children in 1934 were carried out whilst under tenure of the Sebag-Montefiore Research Fellowship, and the medical unit of University College Hospital has defrayed the expenses of the follow-up of sixty-four cases.

REFERENCES

- Bray, G. W. (1931). *Recent Advances in Allergy*, London.
Findlay, L. (1931). *The Rheumatic Infection in Childhood*, London.
Hawksley, J. C. (1931). *Arch. Dis. Childh.*, **6**, 303.
— (1938). *J. roy. Inst. Publ. Hlth. Hyg.*, **1**, 798.
Hill, N. G., and Allan, M. (1929). *British Medical Journal*, **2**, 499.
Keith, J. D. (1938). *Quart. J. Med.*, **31**, 29.
Neustatter, W. L. (1937). *Guy's Hosp. Rep.*, **87**, 8.
Seham, M., and Hilbert, E. H. (1933). *Amer. J. Dis. Childh.*, **46**, 826.
Sheldon, W. (1936). *Diseases of Infancy and Childhood*, London.

CONGENITAL OCCLUSION OF THE POSTERIOR NASAL CHOANAE

BY

DEREK BROWN KELLY, M.B., F.R.F.P.S.

Assistant Surgeon, Throat, Nose, and Ear Department,
Victoria Infirmary, Glasgow

Congenital occlusion or atresia of the posterior nasal choanae was at one time considered to be a relatively rare anomaly. From a search of the literature Fraser (1910) could gather records of only 115 cases. During the past few years, however, more cases have been reported, due no doubt to the fact that not only rhinologists but also obstetricians and practitioners are on the look out for the condition. This form of nasal obstruction really deserves even wider recognition, for the symptoms in a baby are sometimes alarming, and their cause can be a source of puzzlement to the physician. A newborn infant with bilateral choanal atresia suffers from considerable difficulty in breathing. Its instinctive attempts to breathe through the nose result in severe dyspnoea and even cyanosis, which is relieved only when the mouth is opened to cry. A cycle of attacks of dyspnoea alternating with fits of crying and relief of symptoms is thus established, and forms a typical clinical picture. Sucking, and consequently nutrition, are also interfered with.

Unilateral cases are rarely recognized early in life. It is only when the patient is old enough to realize that one nasal passage is blocked, and there is a troublesome discharge of mucus, that relief is sought. Mistakes in diagnosis are not infrequent, the obstruction and discharge being ascribed to sinusitis or adenoids. Operations for the relief of these conditions have been performed without the real state of affairs being recognized.

The obstruction is caused by a failure of absorption of the bucco-nasal membrane. It may be unilateral or bilateral; in structure membranous, bony, or partly membranous and partly bony. Onodi has recorded the only known instance in which cartilage entered into the construction. Females are affected more often than males.

The Cause of Choanal Atresia

Before the embryology of the nose was fully understood, several theories as to the cause of choanal atresia were put forward. It may be of interest to mention briefly the chief of these. Luschka (1860) maintained that the obstruction was due to an upward and backward growth of the horizontal plate of the palatine bone. Kundrat, on the other hand, considered that it was the vertical plates of the palatine bone which extended inwards to meet the inner wall of the choanal orifice (vomer). An original view, taken by Bitot, was that the occlusion consisted of a separate independent bone. The occasional presence of sutures in the bone was put forward in support of his theory.

It is now generally accepted, however, that the obstruction is due to the persistence of a vestigial epithelial structure—the bucco-nasal membrane. This is a partition between the mouth and the nose, closing the posterior end of each olfactory pit in the embryo. It is formed by the apposition of nasal and stomodaeal epithelium, and ruptures about the fifth week, thus establishing the primitive choanae, or openings between the olfactory pits and the stomodaeum. In cases of posterior choanal occlusion this membrane fails to absorb, and the extent of