AETIOLOGY OF GLAUCOMA*

BY

PAUL WEINSTEIN, M.D.

(From the First Ophthalmological Clinic of the Peter Pázmány University and the Albert Apponyi Polyclinic, Budapest)

From the time investigators first described glaucoma increased ocular tension has been regarded as its chief characteristic. But this increase in tension is really only a symptom, the causes of which are manifold.

The current classification of glaucoma into inflammatory, primary, and secondary corresponds to clinical experience. This is confirmed by a series of more than 3,000 cases operated on at the First Ophthalmological Clinic. The common factor in the various forms of glaucoma is increase of ocular tension, but as these forms themselves differ in their clinical manifestations they necessarily differ also in pathogenesis. The earlier investigators believed the increase of tension to be due to an occlusion of Schlemm's canal at the so-called angle of the anterior chamber, by which stagnation is produced. The tendency of hypermetropes to develop glaucoma was similarly explained. But forms of glaucoma were discovered in which the depth of the chamber was normal (simple glaucoma and haemorrhagic glaucoma), this being proved by measurements taken by the slit-lamp and goniometer. Statistical data furthermore revealed that of the glaucoma patients 23 per cent. were emmetropic and 15 per cent. myopic, a proportion sufficiently large to convince investigators that general causes, as well as local alterations, must be concerned.

It seems superfluous to detail the investigations which aimed at determining the influence of the acid-alkaline balance of the body, the internal secretion, the vegetative nervous system, and blood pressure on glaucoma development. Hertel's, and especially Imre's, observations concerning the influence of hypophyseal, thyroid, and ovarian secretion were very important; so was another aspect—that of the relation of glaucoma to the circulatory system.

Relation of Glaucoma to Blood Pressure

Our own research work was undertaken to investigate the possible relation of glaucoma to blood pressure. At the time our investigation began there was a considerable difference of opinion in the literature concerning this question. According to Kümmel, the systolic pressure increases in those suffering from glaucoma; but Elschnig was unable to find any difference in blood pressure when comparing 100 cases of glaucoma with 100 cases of cataract. The investigations were conducted by means of Plesch's tonograph, a device by which information is obtained regarding the state of the vessel walls. The examination revealed the fact that the systolic blood pressure of 100 sound persons ranged from 120 to 200 mm. Hg, and the diastolic from 60 to 95 mm.; while the corresponding values in 150 persons suffering from glaucoma were 120 to 230 mm. and 60 to 115 mm. respectively. The difference is not sufficient to suggest the existence of hypertonia in glaucomatous persons. Attention was next directed to the small vessels. Changes in the general circulation are synchronized to the circulation of the eye through the

precapillaries and capillaries. It has long been known that the oscillations of a tonometer placed on the eye correspond to those of systolic and diastolic pressure. These oscillations were even photographically recorded by means of complicated optical devices. We ourselves used Recklinghausen's tonograph, the reading from which was absolutely identical with the oscillogram taken from the brachial artery.

Pressure within the small ocular vessels can be determined by applying Bailliart's tonometer and that of Seidel. By the former the pressure of the central retinal artery, by the latter that of the anterior ciliary vessels, are determined. Under normal conditions the diastolic pressure of the central retinal artery is half that of the brachial artery. Generally, the pressure within the anterior ciliary vessels varies with that of the central artery. According to the data procured from the application of Bailliart's instrument these pressures are in ratio to the ocular tension: if the ocular tension increases, the diastolic pressure of the central retinal artery increases correspondingly, seemingly in a passive manner regulated by an automatic mechanism. But the values determined within the anterior small vessels are normal, these being independent of the ocular tension, since their pressure is determined before they enter the eye. So far, then, no values that are characteristic of the tension of glaucoma could be determined, either within the large vessels or within the precapillaries. It was found, however, that when the ocular tension increases the pressure within the intra-ocular vessels increases also, but that of the extra-bulbar vessels does not.

O. Müller and Parrisius discovered by capilloscopy that the vasomotor instability of glaucomatous persons was a characteristic peculiarity. Mészáros, Tóth, and I myself came to the same conclusion. We procured a similar valuable result by the use of Herzog's skin-capillary measuring apparatus. By this means is determined the pressure which just suffices to prevent the filling of the arterioles, precapillaries, and capillaries of the skin from the supply of the digital arteries. This pressure Herzog calls "dermal tension," and he calculates its amount to be normally around 40 to 50 mm. Hg. The dermal tension according to Mészáros and Tóth is under 40 mm. Hg, and this figure was confirmed by our own investigation; we furthermore found that the reactive hyperaemia (the filling up of the bloodless finger with blood, normally within one to two seconds) is prolonged. All these data seem to prove that the capillary pressure increases in glaucomatous persons.

During investigations by means of the tonograph we observed that there is a difference between the amplitude of the oscillations in glaucomatous hypertonics and that in sufferers from thrombosis of the central retinal vein. The tonographic reading in glaucomatous persons at 200 mm. Hg pressure shows a normal oscillation of 8 to 10 mm., against over 15 mm. in thrombotic individuals, proving that there is a decrease of elasticity in the walls of the vessels. According to our data inflammatory glaucoma rarely occurs in persons suffering from grave arteriosclerosis (nephrosclerosis and diabetes), but their capilloscopic reading, as well as the tonogram and the dermal-tension value, differs from that found in cases of glaucoma. According to our view the acute capillary dilatation observed during glaucomatous attacks is possible only with an elastic wall of the vessels, and not in the rigid brittle wall of arteriosclerotic persons.

Of the 3,254 glaucomatous patients collected from the material of the First Ophthalmological Clinic during the

* Paper read before the assembly of the Royal Budapest Medical Association on November 18, 1938. Research conducted by a grant of the Széchenyi Scientific Association.

thirty years 1904 to 1933—and of the similar, though smaller, series from the ophthalmological department of the Polyclinic—60 per cent. were females, and 80 per cent. of the acute attacks befell women at the time of climacteric. This ratio explains the relation of glaucoma to the endocrine system and renders the vasomotor instability of glaucomatous persons easily understood, as during the climacteric women are disposed to congestion, perspiration, and similar vasomotor reactions. Furthermore, the frequency of glaucomatous attacks in winter time is conspicuous. This connexion with meteorological conditions points again towards vasomotor instability of the capillaries.

Primary Glaucoma

These data are characteristic of inflammatory glaucoma. Quite another group is that of primary glaucoma. This type is indicated clinically by the slow gradual weakening of vision and concomitant narrowing of the visual field. Externally, conditions seem to be unaltered; ocular tension has hardly increased at all, nevertheless the optic nerve is inevitably destroyed. The capillaries and the dermal-tension values of these individuals are the same as those of persons suffering from inflammatory glaucoma; their tonographic reading is rather that of arteriosclerosis; and according to statistics 56 per cent. of the cases concern men, whereas inflammatory glaucoma is more frequent in women. In one of our cases—that of a man suffering from primary glaucoma of one eye, under our care for years—the eye had to be removed on account of the glaucoma becoming absolute. Histological investigations showed the thick-walled central retinal artery responsible for the tonographic reading of large amplitude. It has to be supposed, consequently, that primary glaucoma, in contrast to the inflammatory type, develops because of arteriosclerotic alteration, and accordingly does not show any great increase in ocular tension. The destruction of the optic nerve seems to be the sequel to circulatory defect and to the resulting alimentary deficiency.

Glaucoma following Retinal Vein Thrombosis

The third type of glaucoma is that following thrombosis of the central retinal vein. Fifty cases of central retinal vein thrombosis were observed by us, and in twenty-three of them glaucoma developed in a few weeks. Pathologically the question why glaucoma follows thrombosis of the central retinal vein in a certain percentage is quite unanswered. First it should be decided in which cases this especially occurs. As a rule the victims are people of advanced age whose blood pressure is high and the amplitude of whose tonographic reading is large. Patients with thrombosis without glaucoma possess relatively satisfactory sight, a sign that the thrombosis has been partial, whereas those who develop glaucoma have previously lost even a sense of light in general, denoting that the thrombosis is situated within the main branch. Histologically such cases display a central retinal artery in which the wall has thickened, the endothelium has proliferated, and hyaline degeneration has set in. All these are types of alteration that are often followed by grave defects of eye metabolism. We know that in whatever organ thrombosis may be located pathological products such as carbonic acid, lactic acid, citric acid, and hydrochloric acid accumulate, and this must necessarily be the case within the eye.

It is probable that under the influence of these acid substances the vitreous body, which essentially is of

colloidal structure, slowly begins to swell up and gradually increases the ocular tension. This assumed swelling of the vitreous body is discussed in the literature. I investigated the matter by applying Krogh's osmometer, and found that the vitreous body shows an osmotic pressure of 98.5 mm. of water as an effect of physiological substances, while with a 1 per cent. solution of lactic acid the pressure increased to 166 mm. These experiments proved without doubt that the vitreous body is capable of increasing the ocular tension when affected by pathological products of metabolism, and to cause development of glaucoma in cases of thrombosis of the central retinal vein, probably by an alteration in the hydrogen-ion concentration.

Commentary

Summarizing the above observations, the part played by the circulatory system in all forms of glaucoma becomes evident—a fact that is proved by the results of the so-called stressing tests, which are applied to ascertain whether the suspicion of a possible glaucoma, suggested by the patient's complaints, is justified. The ocular tension in persons disposed towards glaucoma is increased by several millimetres compared with the normal when they drink a larger amount of liquids or after taking substances containing caffeine (coffee, tea), or if they are put into a position in which the head hangs down. This phenomenon is also explained by the alteration in the circulatory system. The ocular tension of glaucomatous persons considerably increases likewise after inspiration of amyl nitrite, whereas the ocular pressure of arteriosclerotics does not increase at all, or only to an insignificant degree, in similar conditions.

It is evident, therefore, that the endocrine dysfunction of glaucomatous persons exerts its effect by altering the reactive capacity of the vessels and capillaries, or, rather, by disturbing the balance between the osmotic pressure of the blood proteins in the capillaries and arterioles and the ocular tension. Therefore the efforts of various workers to influence the endocrine function of glaucoma patients, chiefly by application of folliculin—and thus in my opinion by the effect on the circulatory system—seem justified, when used in conjunction with either conservative treatment or surgery.

Preparations for the fourth World Conference of Workers for Cripples, which is to be held in London at Bedford College for Women next July, are going forward very satisfactorily. At the official opening on Monday afternoon, July 17, the Minister of Health has agreed to give an address. The meetings will be held in the big hall at Bedford College, and many of the delegates will be accommodated in the college residences, others being housed in hotels near by. Intending delegates should send in their completed application forms as early as possible to the organizing secretary, Conference of Workers for Cripples, 34, Eccleston Square, S.W.1. The conference is officially recognized by the Ministry of Health, the Board of Education, the Scottish Office, and the Ministry of Home Affairs for Northern Ireland, and delegates serving under local authorities should apply through them so that they may claim their expenses. At the first session on July 18 the subject "Preventive Orthopaedics in Childhood" will be opened by Mr. R. C. Elmslie. On July 19 the main subject will be "The Education, Vocational Training, and Subsequent Employment of the Cripple." The morning of July 21 will be devoted to "The Industrial Cripple." On July 20 the delegates will spend the day at Oxford, with a visit to the Wingfield Morris Orthopaedic Hospital. Among other arrangements for July 22 is an excursion to the Heritage Craft Cripple Schools at Chailey.