SURGICAL PROCEDURES IN GENERAL PRACTICE

This is one of a series of articles contributed by invitation

HAEMORRHOIDS

BY

C. NAUNTON MORGAN, M.B., F.R.C.S.

(Concluded from p. 401)

Injection and Operative Treatment

Though the injection method, first introduced by Blackwood in 1866, has an important place in the treatment of haemorrhoids, it has definite limitations. Selection of suitable cases is important, since the results may otherwise be disappointing and dangerous complications occur. Injection treatment is most suitable for small uncomplicated haemorrhoids when the main symptom is bleeding (first degree). Sometimes severe haemorrhage is encountered, which would appear to indicate immediate operative treatment; a well-placed injection, however, will control the bleeding at once. Haemorrhoids which prolapse on defaecation or on proctoscopic examination (second degree) may be injected, since chemical inflammation from high submucous injection above the haemorrhoid will fix the mucosa and contraction of fibrous tissue will draw up the pile into the rectum. It is impossible in cases of prolapse to promise the patient that the condition will be cured by injection; the larger and more easily prolapsed the haemorrhoid the poorer the response. When haemorrhoids are of the third degree, or of the intero-external variety, injection is not only useless but inadvisable. Very occasionally, when only one haemorrhoid remains permanently prolapsed, especially if operation is contraindicated, high injection above the haemorrhoid may be tried, but the patient should be kept recumbent and under observation for a few days after injection, as the haemorrhoid may swell and become strangulated and infected.

It is dangerous to inject haemorrhoids which are thrombosed, infected, or ulcerated. Associated infection such as an abscess, fistula, or a chronic fissure or proctitis is also a contraindication to injection.

Technique of Injection

The instruments necessary are a 10 c.cm. Record syringe with a long needle which has a collar $\frac{1}{8}$ inch from its point, and which is attached to the syringe by a locking device (this is necessary owing to the viscosity of the solution injected); a two-inch proctoscope; a good light; a pair of long plain dissecting forceps; and an artery forceps. The angular needle illustrated (Fig. 4) has a simple locking mechanism and enables a clear view to be obtained whilst injecting. An illuminated proctoscope is also useful (Fig. 5). For those not conversant with this treatment the knee-elbow position is the easiest, since when the proctoscope is inserted the ampulla of the rectum is distended with air and the landmarks are more obvious. When experience has been obtained the left lateral position, with a small pillow under the left buttock, is much more comfortable for both patient and operator.

There are two methods of injection, both of which have their place: (a) the submucous injection of 5 per cent. phenol in almond oil, with the addition of 2 grains of menthol to the ounce; (b) the injection of 20 per cent. phenol in glycerin and water, equal parts, into the centre of the pile itself.

THE SUBMUCOUS METHOD

The proctoscope is slowly and gently inserted into the rectum, its obturator removed, and the instrument slowly withdrawn until the upper limit of the ano-rectal ring is seen. The correct site for the first submucous injection is just above this ring, into the highest portion of the internal haemorrhoidal plexus (Figs. 2 and 2a). The haemorrhoids themselves are not seen when the proctoscope is in this position (compare Figs. 3 and 3a). The injection is as a rule made at points corresponding with the position of the primary haemorrhoids, but because

FIG. 4.—Syringe and angular needle.

FIG. 5.—Illuminated proctoscope.

the solution spreads readily in the submucous space around the whole circumference this is not absolutely necessary. The needle is inserted through the mucous membrane into the submucous layer and its point felt and seen to move freely in this layer before injection is carried out. The solution is injected slowly and the mucous membrane will be seen to be lifted up, and as injection is continued a swelling appears, on the surface of which tiny capillaries are visible running on a background tinged slightly yellow by the almond oil. Should a dead-white area begin to appear during injection the solution is being placed too superficially, and ulceration will result if injection is not immediately stopped. When the previously described swelling is seen the needle is slowly withdrawn, some of the solution being injected along the needle track during withdrawal. Bleeding from the needle puncture, should it occur, usually ceases on removal of the proctoscope, but should it not do so, or if it is profuse, application of an artery forceps for a minute will control the bleeding. The amount of solution injected depends upon the laxity of the mucous membrane. Generally 2 to

4 c.cm. is necessary at each point of injection, but sometimes a large haemorrhoid will require 6 to 8 c.cm.

One injection only is given at the first visit, as rarely a general reaction may occur due to sensitivity to the almond oil, since it does not arise after injection of the stronger solution, which contains glycerin and water. At subsequent visits two or three injections at different sites are given. The second injection is made about ten days later. If a good reaction from the first injection has been produced, the mucous membrane will feel tough and leathery both to the palpating finger and to the needle. Injection above the two other haemorrhoids is now carried out, and if there is no reaction at the place of the first injection another injection is given at the same site. Subsequent submucous injections are always made through *soft mobile* mucous membrane at intervals of ten to fourteen days. When a good reaction has been obtained at the site of the highest injection an additional submucous injection is given at a slightly lower level. Usually three or four are required for each haemorrhoid.

Injection into the Haemorrhoid.—Sometimes, although there is a good response to submucous injections, the lower portion of the haemorrhoid still remains prominent. In these circumstances a final injection of 20 per cent. phenol in glycerin and water—equal parts—is given deeply into the centre of the pile mass itself. This solution should never be used for submucous injection. Five to seven minims only are injected, using a Graeme-Anderson syringe and needle.

Injection treatment is ambulatory, and as it is given through insensitive mucous membrane it is as a rule painless. If the solution is injected quickly, discomfort or a sensation of distension may be experienced. A few hours after injection a fullness in the rectum may be felt. Acute pain on injection indicates that the injection has been given too low and into the wrong layer. Violent exercise should be avoided for at least twenty-four hours following injection. The response to injections is variable, and it is impossible to be sure of the length of time their effect will last. A large proportion of cures may be expected with haemorrhoids of the first degree, and bleeding can always be stopped, but when haemorrhoids prolapse recurrence of symptoms occurs after two years or longer, depending on the amount of the prolapse.

Operative Treatment

Internal haemorrhoids which remain continually prolapsed or which prolapse easily after replacement should be treated by operation. If injection fails to control the prolapse, and if it recurs on walking or straining or at work, non-operative methods are usually futile. It has been stated that haemorrhoids frequently recur after operation. This is true, but the reason for these recurrences is that in the majority of instances the internal haemorrhoids had only been partially removed. Pre-operative treatment consists in giving an aperient two nights before the operation, a soap-and-water enema the evening before, and a saline wash-out three hours before. General anaesthesia, using a basal narcotic with cyclopropane or gas-and-oxygen, may be employed, but the most satisfactory method is a low spinal anaesthesia, since the prevention of post-operative vomiting and coughing minimizes the pain and there is no oedema of the perianal skin. Local anaesthesia, using 1 per cent. procaine, supplemented by pentothal intravenously, is especially suitable when other methods are contraindicated.

The patient is placed in the lithotomy position and the perianal skin and anus are cleansed with spirit. The left

forefinger is inserted into the rectum and gently hooked around the ano-rectal ring. The skin behind the anus is painted with iodine and 10 c.cm. of warm procaine (an oily solution containing anaesthetic substances having a prolonged action) is injected through a needle puncture in the midline one inch behind the anus (Figs. 6 and 7).

FIG. 6.—Method of injection of anaesthetic solution.

The injection is given into the external sphincter and deeply under the skin around the anus in a fan-like manner; strict asepsis must be observed during the injection. The inserted forefinger, which acts as a guide to the position of the needle, is not removed until the injection has been completed. The perianal tissues are then gently massaged to ensure even distribution of the oily solution. Oil-soluble anaesthetics, if properly injected, considerably lessen post-operative spasm and pain. *The injection should not be given at the end of the operation.*

FIG. 7.—Showing depth of injection; also the method of infiltration for removal of an anal haematoma.

It is unnecessary to dilate the anus. A tag of skin at the anal margin is pulled outwards from the anus with dissecting forceps. This will deliver the corresponding internal haemorrhoid through the anal orifice. Sometimes, of course, the internal haemorrhoids are already protruding. The internal haemorrhoid is grasped by artery forceps, which is handed to an assistant, who retracts the haemorrhoid radially from the anus. The two remaining haemorrhoids are delivered in a similar manner and held by an assistant. When the three primary haemorrhoids are thus retracted, three longitudinal folds will be seen passing upwards into the rectum from the haemorrhoids. These folds are the pedicles of the haemorrhoids, and must be included in the ligatures to ensure radical excision. Each of these folds is seized with artery forceps, and the

three previously applied artery forceps are removed. When the forceps on the pile pedicles are retracted radially, folds of pink rectal mucous membrane will be seen passing between each pedicle, forming a triangle. Until this "triangle of exposure" is visible the haemorrhoids are not properly exposed (Fig. 8).

FIG. 8.—Showing the three primary internal haemorrhoids fully exposed. Note triangle of exposure.

The whole area is now cleansed with ether soap and with dettol, 1 drachm to the pint. Starting with the left lateral haemorrhoid, the tip of the left index finger is placed at the level of the subcutaneous external sphincter and gentle outward pressure exerted. The forceps holding the pedicle of the haemorrhoid is then placed in the palm of the same hand. The skin covering the corresponding portion of the external haemorrhoidal plexus is grasped in another artery forceps, which is also placed in the palm of the left hand and gently pulled inwards. A V-shaped cut is now made with a pair of scissors through the skin of the anus, each limb of the V being on either side of the forceps retracting the skin (Fig. 9). This

FIG. 9.—Showing the skin cut.

portion of skin, together with the underlying external haemorrhoidal plexus, is dissected cleanly off the subcutaneous external sphincter, which is thus laid bare. This wedge of tissue is further dissected inwards until the inner border of the subcutaneous external sphincter is exposed. At the inner border of this muscle the longitudinal muscle fibres of the anal canal will be seen passing downwards into the tissues now freed by the dissection.

The dissection is now complete, and the internal haemorrhoid, together with the corresponding external plexus, is held only by the pedicle of the internal haemorrhoid and the longitudinal muscle. Traction is exerted by an assistant on both the pedicle and the skin forceps, and the pile pedicle transfixes with No. 2 chromic catgut on a round-bodied needle and ligatured above the forceps

holding the pedicle of the pile (Fig. 10). The other two primary haemorrhoids are dissected and ligatured in a similar manner. For the right posterior haemorrhoid the artery forceps are held in the palm of the left hand. A strip of anal skin and mucous membrane should always be left between each haemorrhoid. Secondary haemorrhoids, when present, are either included with the ligatures of the primary piles or, if associated with a skin tag, are

FIG. 10.—Showing position of skin cut and ligation of haemorrhoid pedicle. (C. J. C. Siggers.)

dissected in a manner similar to that used in dealing with the primary haemorrhoids. The ligature, however, is placed no higher than the level of the triangle of exposure, as these piles have no pedicles.

The haemorrhoids are now removed, the ligatures cut short, and the pedicles returned into the anal canal. It will be found that the ligatured haemorrhoids will not retract far inside the anal canal, so that large raw areas will not be left below them. Redundant skin is removed from the edges of the three wounds at the anal orifice, leaving them flat and somewhat triangular. Three squares of gauze soaked in crude cod-liver oil are now inserted into the anal canal and laid flat on the radiating wounds, and a good pad of dressings applied and held in place by a firm tight T-bandage.

The external dressings are changed daily and gauze moistened with eusol is reapplied. Liquid paraffin and a normal diet are given from the second day onwards, and on the evening of this day an aperient is administered. On the third morning, when the patient has a desire for defaecation, the bowels are moved with an olive oil and gruel enema. After the bowels have acted the patient is placed in a hot bath, and on return to bed the external wounds are gently irrigated. The inserted dressings are, as a rule, extruded during defaecation. Hereafter, moist antiseptic dressings are applied twice daily following baths.

It is important to pass a finger into the anal canal on the ninth, eleventh, and twelfth days after operation to anticipate any constriction. The patient leaves the hospital on the twelfth day, but should be seen at the end of a week, and again in a fortnight, for the passage of a finger, since though constriction does not follow a properly performed operation, it is wise to anticipate its formation. For further details of the operation the reader is referred to an article by Milligan *et al.* (1937).

Correct diagnosis, gentleness, and attention to detail in pre-operative, operative, and post-operative technique are essential for successful results in the treatment of haemorrhoids.

Our thanks are due to Messrs. Vann Bros., London, for the loan of blocks of instruments illustrated in this article.

REFERENCE

Milligan, E. T. C., Morgan, C., Naunton, Jones, L. E., and Officer, R. (1937). *Lancet*, 2, 1119.