bution arising from the axillary basal (described by Ewart as being its only origin) or from the posterior basal. As stated above, it is additional to all other constant bronchi in the lungs of man and of the animals we have examined. The reason for its occasional absence from the common site in the left lung of man and its irregular appearance within the limits mentioned is not at present clear. It is possibly becoming vestigial in man, as its variations would suggest. We have named it the subapical, and in metal casts and by single bronchial fillings with lipiodol and by inflation have shown its distribution to be to the mesial, posterior, and lateral areas of the lower lobe between the apical and posterior basal segments.

Commentary

With the realization of its varied location and the recognition that the other segmental bronchi are constants, the bronchial tree can be described with the surety of ordinary anatomical description.

This dissertation cannot be suggested as detracting from the work of Ewart, who must have devoted many years to his investigations. Alone, and with crude instruments and materials, he described fully the bronchial tree even to its minute branches. Increasing experience of later investigators has confirmed the general accuracy of this master of bronchial anatomy.

BIBLIOGRAPHY

- Aeby, C. T. (1880). *Der Bronchialbaum der Säugethiere und des Menschen*, Leipzig. (Quoted by Ewart.)
 Ewart, William (1889). *The Bronchi and Pulmonary Vessels*, J. and A. Churchill, London.
 Kramer, Rudolph, and Glass, Amiel (1932). *Ann. Otol.*, **41**, 1210.
 Neil, J. Hardie, Gilmour, W., and Main, W. W. (1936). *Proceedings of Second Congress, Pan-Pacific Surgical Association*.
 — Gwynne, F. J., and Gilmour, W. (1937). *Med. J. Austral.*, **2**, 165.
 — — — Main, W. W., and Fairclough, W. A. (1937). *Ann. Otol.*, **46**, 338.
 — — — — — (1938). *Austral. New Zeal. J. Surg.*, **8**, 118.
 Nelson, H. P. (1934). *British Medical Journal*, **2**, 251.

The eighth annual report of the National Birth Control Association (president, Lord Horder), with which is now incorporated the Birth Control International Information Centre, the Birth Control Investigation Committee, and the Society for the Provision of Birth Control Clinics, points out that the amalgamation last spring has left the N.B.C.A. as the only central birth control organization apart from the Society for Constructive Birth Control. The chairman and secretary gave evidence before the Interdepartmental Committee on Abortion, a memorandum having been previously sent in. The medical subcommittee, under the chairmanship of Mr. A. C. Palmer, completed another year of active work, and among leaflets issued was one entitled "Services to Private Practitioners by the Medical Subcommittee of the N.B.C.A." A Scottish Advisory Committee was formed in 1938 and a number of new branches were set up in England and Scotland; some of them have already opened clinics. Early last spring a move was made to new and larger headquarters at 69, Eccleston Square, London, S.W.1. A conference of all branches, held in London before the annual meeting, was so successful that this will now be an annual event. Out of the 409 maternity and child welfare authorities in England and Wales 246 have now taken action of one sort or another under the Ministry of Health memoranda. Of these 246, seventy-nine now have post-natal gynaecological or birth control clinics; 113 refer cases to voluntary or municipal clinics, to hospitals, or to private practitioners; and thirty-eight make grants or lend premises, or do both, to voluntary clinics. Some take action in more than one of these ways.

TWO CASES OF PICK'S DISEASE STUDIED WITH THE AID OF KYMOGRAPHY

BY

G. L. S. KONSTAM, M.D., M.R.C.P.

*Physician to the West London Hospital, with Charge of the
Cardiographic Department*

AND

FRANKLIN G. WOOD, M.B., D.M.R.E.

*Radiologist to the London Chest Hospital and the
German Hospital, Dalston*

(WITH SPECIAL PLATE)

The following two cases of Pick's disease (chronic constrictive pericarditis) were considered worthy of publication, not only because of their comparative rarity, but because opportunity was afforded of studying them by kymography. The first case is of particular interest, as kymograms were taken before and after operation.

Our knowledge of this syndrome and its treatment by pericardial resection has been considerably extended by Paul D. White (1935), whose St. Cyres Lecture summarized his study of fifteen personal cases. The salient feature met with is fibrous or callous thickening of the pericardium, which by gripping the heart in an unyielding envelope prevents normal diastolic filling and leads to what has been termed "inflow stasis." The syndrome occurs in both sexes, and its onset is more usual in early than in middle life. The aetiology is often obscure, but there may be a past history of pneumonia, tuberculosis, polyserositis, pleurisy, and acute pericarditis. It is noteworthy that in none of White's series was there any evidence of a rheumatic infection.

The main symptoms are dyspnoea and swelling of the abdomen; the signs are similar to those observed in chronic congestive heart failure, with the important exceptions that the heart is not usually enlarged and evidence of valvular disease is absent. The jugular veins, and often the arm veins, are greatly engorged, the liver is enlarged but not tender, and ascites develops which is out of all proportion to any oedema that may be present in the extremities. The blood pressure and pulse pressure are below normal, and pulsus paradoxus is often found. Normal rhythm is usual, but in four of White's cases, and in one here described, there was auricular fibrillation.

X-ray Appearances

The heart shadow is normal in size and shape, or moderately enlarged. Calcification of the pericardium is frequently present, but its absence does not exclude the diagnosis; congestion of the pulmonary vessels and enlargement of the supracardiac shadow are also met with. On screen examination the normal heart pulsations are greatly reduced, and may even be absent, but these findings are more accurately shown by the modern method of kymography. An electrocardiogram may show low-voltage flattening or inversion of the T waves in Lead I or II, or in both leads. When inverted the T wave is coronary in Type. Low-voltage QRS waves are also commonly found.

Without surgical treatment the prognosis is hopeless, the disease usually terminating fatally within a few years. Resection of the pericardium by operation may, on the other hand, lead to restoration of normal health.

Case 1: Pick's Disease of Unknown Origin

A girl aged 9 attended the out-patient department at the West London Hospital in February, 1932, to 1933 suffering

from bronchitis. In 1936 she was admitted to the hospital under Dr. Sydney Owen. During the intervening three years she had had chicken-pox, mumps, measles, and bronchitis.

The girl was very debilitated and thin. The tonsils were much enlarged, and their crypts were full of carious debris. The lips were slightly cyanosed and the cervical veins engorged even in the upright position. The pulse was regular, rate 94, and the blood pressure 102/78. The heart was not clinically enlarged, and Broadbent's sign was absent. No murmurs were present, although a harsh systolic murmur had been noted before admission to hospital. A few rales were heard over the lung bases. The abdomen was distended and the umbilicus everted, but there were no signs of ascites. The liver was much enlarged, firm, and not tender. There was no oedema of the extremities. The urine contained no albumin or sugar. The cardiac impulse was poor over all chambers, but the mobility of the heart was unimpaired. There was considerable pulmonary vascular engorgement. Four examinations of the sputum were negative for tubercle bacilli, and the Mantoux reactions (1 in 10,000 and 1 in 1,000 dilutions) were negative. Blood Wassermann and Kahn reactions were negative. A blood count on September 14, 1936, gave: red cells, 4,400,000 per c.mm.; haemoglobin, 88 per cent.; white cells, 17,000 per c.mm.; colour index, 1.1. Anisocytosis was present. An x-ray examination made on August 25, 1936, by Dr. H. W. A. Post showed that the heart was not obviously enlarged. There was a prominence of the left border in the pulmonary artery region. A deposit of calcium was visible in the pericardium in the anterior and lateral views. The lung fields were clear.

During the five-weeks observation in hospital her condition improved somewhat, but slight pyrexia up to 99.6° continued until the time of her discharge. Three months later she was able to return to school; owing, however, to swelling of the abdomen and shortness of breath she soon had to be withdrawn. In January, 1938, the father wrote that she could only move about with difficulty in her room, and, apart from this, had to be wheeled around in a push-chair. Frequent cough added to her difficulties.

On January 24, 1938, at the age of 9½, she was readmitted to the West London Hospital. She was mentally backward and shy. Her face was puffy, and the veins in the neck were engorged when she was in the sitting position. Moderate orthopnoea was present. The pulse rate was 100, and pulsus paradoxus was absent. The apex beat was in the fifth space in the mid-clavicular line, and systolic retraction was visible in this region. The mitral first sound was accentuated, but no murmurs were present. The percussion note at both lung bases was impaired, owing to the high position of the diaphragm. The abdomen was grossly distended from ascites and the lower border of the liver was three inches below the costal margin. There was no oedema of the feet. On investigation the circulation rate (3 c.cm. 20 per cent. decholin solution), antecubital fossa to tongue, was 20 seconds (normal 9 to 16 seconds).

From a pre-operative x-ray examination made on January 11, 1938, it was observed that both domes of the diaphragm were elevated, causing the heart to lie more transversely; the deposits of calcium had increased and were greater on the anterior, left lateral, and inferior aspects of the heart. Congestion of the pulmonary vessels was visible. A kymogram (Plate, Fig. A) was taken the same day and the following appearances were noted: right border, pulsations almost absent; left border, aortic pulsations were not clearly defined. Abnormal movements were visible in the prominent middle arc region and slight movement in the upper ventricular area. The apex of the left ventricle, which was seen to be encased in a thick layer of calcium, was quite immobile.

OPERATION

On February 7, 1938, paracentesis abdominis yielded fourteen pints of straw-coloured fluid. Paracentesis was repeated two weeks later, and another ten pints were drained. On February 23, 1938, transternal pericardiectomy was performed by Mr. J. E. H. Roberts, assisted by Mr. A. Simpson-Smith.

Details of the surgical aspects of the case are to be published later by Mr. Roberts. The visceral pericardium was thickened and calcified, and was in some places firmly adherent to the myocardium; in other places it was more easily separated. It was incised and stripped off the anterior and left lateral aspects of the heart, in the process of which both pleural cavities were accidentally opened. Over the auricles the pericardium was more densely adherent, and owing to this and the patient's poor condition it was deemed advisable to terminate the operation as soon as possible.

POST-OPERATIVE HISTORY

Dr. R. G. Waller examined a section of the pericardium and found that it was greatly thickened (5 mm. in its thickest area) by an increase of collagenous and fibrous tissue; numerous calcareous plaques were present. There was no histological evidence of tuberculosis. Before the patient was returned to bed sufficient air was evacuated from both pleural cavities, by means of a pneumothorax apparatus, to leave slightly negative pressure. A drip blood transfusion was given, and during twenty days of a stormy post-operative period an oxygen tent was used. A week after the operation twelve ounces of ascitic fluid were aspirated from the abdomen, but since then there has been no further tendency for the fluid to reaccumulate and the abdomen has remained flat. Ten days after operation the circulation rate had slightly improved—19.5 seconds from antecubital fossa to tongue. Subsequently convalescence was uneventful, and when examined seven months after operation the patient had gained 8 lb. in weight, was leading a normal life, and was able to run round the ward without becoming short of breath; the distension of the cervical veins had disappeared.

Serial electrocardiograms (Figs. 1 and 2) were taken. Before operation (Fig. 1) the QRS and T waves were of low voltage in all leads. Three weeks after operation the T waves in Leads I and II became negative, but seven months later (Fig. 2) the T waves in the first two leads again became positive, and were of much higher voltage than in the pre-operative period. The QRS complexes in the limb leads were also much larger. Notching of the P waves in Lead II and right axis deviation resembled the findings in mitral stenosis.

FIG. 1.

FIG. 2.

FIG. 1.—Case I. Electrocardiogram (pre-operative) taken on January 11, 1938. Normal rhythm, rate 93 per minute. The QRS and the T waves in the limb leads, low voltage. The P waves in Lead II, large and bizarre. Notching of the T waves in the chest lead (IV R).

FIG. 2.—Case I. Electrocardiogram taken on September 16, 1938, seven months after pericardial resection. Slight right axis deviation has developed. Higher-voltage QRS and T waves in all leads. P waves in Lead II, notched.

X-ray examination on August 16, 1938, six months after operation, showed that the diaphragm was no longer elevated, but despite this the heart shadow was larger: the transverse diameter was 12 cm., which represented an increase of 1 cm. over the pre-operative measurement. Calcium deposits could now only be seen on the inferior aspect of the heart. The

prominence in the region of the pulmonary arc persisted, but a barium swallow examination, in the right oblique position, showed no enlargement of the left auricle.

A kymogram (Fig. 3) was taken on September 28, 1938, seven and a half months after operation. Pulsations in the right border were still minimal. The left border showed great increase in the pulsations, especially in the apex of the left ventricle, where they had been previously absent. The wave form, which did not conform to the normal ventricular pattern, is shown in the tracing; this may be due to unequal decompression.

FIG. 3.—Case I. Tracing of a kymogram taken seven and a half months after pericardial resection. Compared with Fig. A (Special Plate) the pulsations have reappeared at the cardiac apex, and are more obvious over the left border of the heart.

Case II : Pick's Disease ; Pulmonary Tuberculosis ; Auricular Fibrillation

A man aged 52 was admitted to the East Ham Memorial Hospital on March 23, 1937. There was a history of pneumonia and pleurisy six years previously, and since then he had complained of slight cough, increasing shortness of breath, and night sweats. For two weeks before admission his feet and abdomen had begun to swell.

On examination his nutrition was seen to be very poor; the eyes were rather prominent, but the thyroid was not enlarged; he was dyspnoeic and orthopnoeic. The veins of the neck, in the sitting position, were distended up to the angle of the jaw. The pulse was completely irregular in time and volume, and the rate was 100 per minute; the blood pressure was approximately 98/65. The apex beat could not be felt, but the left border of cardiac dullness was half an inch external to the mid-clavicular line, and the cardiac dullness did not shift with movement. The heart sounds were distant. There was flattening of the chest at both apices, and below the right clavicle the percussion note was impaired and crepitations were present; in addition rhonchi were heard over both lungs. The abdomen was tense and distended; the lower border of the liver could be felt two fingerbreadths below the costal margin, and there was dullness over the flanks. There was moderate oedema of the feet and ankles.

On admission the patient's temperature was 99°, but the next day it fell to normal, and remained so during the six weeks he was kept under observation. The urine contained a trace of albumin. The circulation rate was 25 seconds from the antecubital fossa to the tongue. An electrocardiogram (Fig. 4) showed auricular fibrillation with flattening of the T wave in Lead I, negative T waves in Leads II, III, and in the chest lead (IV R). Tubercle bacilli were present in the sputum. The Kahn reaction of the blood was negative.

X-ray Appearances.—Anterior and lateral views were taken of the chest and also a kymogram. In the anterior view the trachea and upper part of the mediastinum are displaced to the right. The heart is normal in size, but slightly irregular in outline, especially on its left border. An irregular deposit of calcium appears to be situated chiefly on the left side and inferior aspect of the heart. The upper part of the right lung

is opaque owing to thickening of the pleura. An area of translucency below the clavicle suggests the presence of cavitation. A deposit of calcium can be seen in the lung in this region. The middle and lower zones were comparatively clear. The right costo-phrenic angle is blocked by thickening of the pleura. Mottling is present in the left lung in the upper and middle zones. The left costo-phrenic angle is partly blocked by pleural thickening. In the lateral view a

FIG. 4.—Case II. Electrocardiogram taken on May 4, 1937. Auricular fibrillation, rate 94 per minute. QRS waves in Leads I and III, low voltage. T in Lead I, flat. T in Leads II and III, and in the chest lead (IV R), negative. Digitalis was not being administered at the time of examination.

marked deposit of calcium can be seen on the inferior border of the heart and also anteriorly where it extends in a continuous sheet downwards from the internal aspect of the manubrium sterni.

A kymogram was taken, but owing to the presence of dyspnoea it was necessary to make a record during quiet respiration. There were two respiratory phases during the exposure time of three seconds, the heart moving to the right during expiration and to the left during inspiration. The normal pulsations were absent except at the apex of the left ventricle, where small movements could be made out.

With rest, digitalis, and salyrgan injections moderate improvement occurred, but owing to the evidence of active tuberculosis in the lungs pericardial resection was not performed. The patient was subsequently discharged from hospital, and died in his home on July 12, 1937. No post-mortem examination was made.

Summary

Two cases of Pick's syndrome which were investigated by radiology, including kymography, are described. Case I, of unknown origin, was studied before and after operation. Case II, which was due to tuberculosis, was unsuitable for surgical treatment.

Kymography was found to be of value in showing the limitation of heart movement and as providing a record of the changes following surgical treatment.

REFERENCE

White, P. D. (1935). *Lancet*, 2, 539.

The Ministry of Labour has published a new edition of the pamphlet on *Dentistry*, which is No. 14 of the series compiled by arrangement with the Incorporated Associations of Headmasters and Headmistresses of Public Secondary Schools. The pamphlet sets out in detail the qualifications and training required and the prospects open to boys and girls of secondary school education wishing to make a career in dentistry. It may be purchased from H.M. Stationery Office or through any bookseller, price 3d.

E. H. HUDSON: SINGLE LARGE CYST OF LUNG SIMULATING HIGH-PRESSURE PNEUMOTHORAX

FIG. 1.—Radiograph showing the pulmonary remains at the apex and the base. August 13, 1936.

FIG. 2.—Radiograph showing the partially expanded lung and a pneumothorax. September 12, 1936.

FIG. 3.—Radiograph taken on September 24, 1936.

G. L. A. KONSTAM AND FRANKLIN G. WOOD: TWO CASES OF PICK'S DISEASE

FIG. A.—Kymogram of Case I before operation, showing the deposit of calcium on the lateral and inferior borders of the heart. The pulsations are absent at the apex of the left ventricle, and are minimal at the right border.

MILROY PAUL: ENTERIC INTUSSUSCEPTION DUE TO THE INVAGINATION OF MECKEL'S DIVERTICULUM

Inverted Meckel's diverticulum in lumen of resected gut.