foreign white cells; and the red cells of the transfused blood were rapidly phagocytosed. There was no other evidence of incompatibility between the patient's blood, which was Group A, and the donor's, which was Group O. There was no reaction following the transfusion and no haemoglobinuria. The death of the patient seven days later was easily accountable by his general condition.

The phenomenon of auto-erythrophagocytosis—ingestion by macrophages of the patient's own red cells—has been observed *in vitro* and has been described as occurring in the circulation in malaria and in trypanosomiasis by Connal (1912) and Macfie and Johnston (1913). In the literature consulted no reference was found to erythrophagocytosis either in agranulocytosis or as a result of blood transfusion. In this case the condition probably indicates either a degree of incompatibility between patient's and donor's blood insufficient to produce other reaction or, more likely, increased sensitivity of the patient to foreign red blood cells.

For the data and material described in Case II I am indebted to Air Commodore H. E. Whittingham.

REFERENCES

- Connal, A. (1912). *J. Path. Bact.*, **16**, 502.
 Cooke, W. E. (1927). *British Medical Journal*, **1**, 12.
 — (1934). *J. Lab. clin. Med.*, **19**, 453.
 Jackson, H., Parker, F., and Taylor F. H. L. (1932). *Amer. J. med. Sci.*, **184**, 297.
 Kennedy, W. P., and MacKay, I. (1936). *J. Physiol.*, **87**, 336.
 Macfie, J. W. S., and Johnston, J. E. L. (1913). *J. Lond. Sch. trop. Med.*, **2**, 212.
 Osgood, E. E., and Ashworth, C. M. (1937). *Atlas of Haematology*, Stacey, San Francisco.

THE ANATOMY OF THE BRONCHIAL TREE*

BY

J. HARDIE NEIL, D.S.O., F.R.A.C.S., F.A.C.S.

President, Pan-Pacific Surgical Association; Aural Surgeon, Auckland Hospital, New Zealand

W. GILMOUR, M.D.Glasg., D.P.H.Oxon

Pathologist, Auckland Hospital

AND

F. J. GWYNNE, M.B.Sydney

Radiologist to Mater Misericordiae Hospital, Auckland

(WITH SPECIAL PLATE)

Recent advances in radiology, bronchoscopy, and thoracic surgery have called for a description of the anatomy of the bronchi that will allow students and clinicians to visualize readily their distribution to the lung tissue. Current anatomical textbooks do not furnish the clinician or thoracic surgeon with the means for the precise localization of pulmonary lesions. The technically perfect film and the brilliant bronchoscopic field are of little avail if there is uncertainty in the conception of the anatomy of the bronchial tree.

The Work of Ewart and Others

Ewart (1889), in his classical work on the bronchi and pulmonary vessels, feared that excessive detail might cause the student to recoil from the study. His book contains a meticulous description of the bronchi down to the finest peripheral branches. He made continued reference to the work of Aeby (1880), who approached the subject from the point of view of comparative anatomy.

A comparison of the diagram of the bronchial tree of Aeby with that of Ewart shows that the former allows one to visualize the bronchial tree at a glance, while the latter with its multiplicity of smaller branches calls for an intensive study that confuses any who attempt to get a practical conception of the fundamentals. Ewart's description of the foliage and finer twigs tends to obscure the picture of the trunk and main branches, the recognition of which is all-important for a reasonably rapid conception of the general architecture of the lungs. Kramer and Glass (1932), apparently unaware of Ewart's work, used a terminology which allows one to picture readily the location of the segments involved by lung abscess. Nelson (1934), whose untimely death must be deemed a catastrophe to thoracic surgery, published an article dealing mainly with bronchiectasis, in which his lucid exposition of the anatomical data showed that he was conversant with the work of Ewart.

It is a tribute to Ewart and Aeby that, with crude appliances and lack of radiology and modern pathological technique, they arrived at a description which can hardly be improved upon apart from some detail. Ewart was quite definite as to the nature of the broncho-pulmonary segment as later described by Kramer and Glass. The latter state that each tertiary bronchus, with the lung tissue it serves, forms a distinct entity, the limits of which can be defined and the relations of which to the chest wall can be delineated. Ewart said that "within each lung large groups of lobules are kept in practical isolation from each other as regards their air supply. Each of these sublobar groups may be considered as forming separate respiratory districts." The recognition of this fact must cause all clinicians and radiologists to think and speak in terms of segments in the localization of lesions such as infections which apparently in the first instance affect a single broncho-pulmonary segment. Only in gross pathology do they pass over the border into adjacent segments and lobes (Figs. 1 and 2). Ewart certainly places students under a debt of gratitude by applying names to the bronchi that connote their territorial distribution and anatomical relations, thus avoiding the terminology of Aeby with its repetition of similar names qualified only by numerals.

Nomenclature: Some Suggestions

At this point a few observations on the terms used in describing the bronchial tree may be useful. The terms primary, secondary, and tertiary bronchi have been given different interpretations by different writers. It seems to us important to recognize as definite entities those branches which supply separate pulmonary segments that can be sharply defined by inflation and the relations of which to the chest wall can be delineated. By some these have been termed tertiary bronchi. We would suggest that they be termed segmental bronchi. This would avoid all confusion in regard to nomenclature.

The main bronchus extends from the bifurcation of the trachea down to its final subdivision into the axillary basal and posterior basal segmental branches. There appears to be no reason, either anatomical or clinical, for subdividing the main bronchus into one or more parts. We would therefore suggest that it be referred to as the main bronchus. Anatomical and clinical localization is readily afforded by the bifurcation and by the openings of the various vestibules and branches. As the segmental bronchi to the upper and middle lobes on the right side and the upper lobe on the left side (which is common to the upper and middle) do not arise directly from the main bronchus it is necessary to have a name for this intervening part, and we suggest that these areas be

* Read at the meeting of the Auckland Division of the British Medical Association, November, 1938.

referred to as the vestibules. We therefore think that the following names might be used to identify the various parts of the bronchial tree: (1) main bronchi; (2) vestibules; (3) segmental bronchi. There is no practical utility in giving names to the further subdivisions to the bronchial tree. Any attempt to do so would introduce a complicated system of nomenclature, which we have set out to avoid.

We now show the tables of Ewart and of Kramer and Glass naming the segmentary bronchi, and offer our suggestions for a revised nomenclature.

EWART	KRAMER and GLASS	SUGGESTED NOMENCLATURE
<i>Upper Lobe :</i>	<i>Upper Lobe :</i>	<i>Upper Lobe :</i>
Apical— <i>a</i>	Apical— <i>a</i>	Subapical— <i>b</i>
Posterior apical— <i>b</i>	Paravertebral— <i>b</i>	Apical— <i>a</i>
Pectoral— <i>c</i>	Anterior— <i>c</i>	Axillary— <i>c</i>
Axillary— <i>d</i>	Axillary— <i>d</i>	Anterior— <i>d</i>
<i>Middle Lobe :</i>	<i>Middle Lobe :</i>	<i>Middle Lobe :</i>
Pectoral— <i>e</i>	Anterior— <i>e</i>	Anterior— <i>e</i>
Axillary— <i>f</i>	Axillary— <i>f</i>	Axillary— <i>f</i>
<i>Lower Lobe :</i>	<i>Lower Lobe :</i>	<i>Lower Lobe :</i>
Posterior horizontal— <i>g</i>	Apical— <i>g</i>	Apical— <i>g</i>
Retrocardiac— <i>h</i>	Cardiac or mesial— <i>h</i>	Cardiac— <i>h</i>
Antero-basal— <i>i</i>	Antero-lateral— <i>i</i>	Subapical— <i>j</i>
Lesser posterior horizontal— <i>j</i>	Postero-lateral— <i>k</i>	Anterior basal— <i>i</i>
Axillary basal— <i>k</i>	Paravertebral— <i>l</i>	Axillary basal— <i>k</i>
Posterior basal— <i>l</i>		Posterior basal— <i>l</i>

The letters in italics denote the segmental bronchi in the order given by Ewart, and Kramer and Glass.

In this list we have placed the bronchi in the sequence in which they leave the bronchial tree. This arrangement is helpful to bronchoscopists. Reference is made to the variable position of the subapical bronchus (or bronchi) in the lower lobe later in the text (Figs. 3 and 4).

Upper Lobe

In the upper lobe, Aeby and Ewart gave full representation to the lateral distribution of the posterior apical of Ewart, which we term subapical. Kramer and Glass, to whom we personally are indebted for elucidation of the bronchial tree, were apparently not aware that the surface distribution of the segment extended to the junction of the greater and lesser fissures of the lung on the right side, where we have invariably found it distinct from, and not a branch of, the apical as described by Ewart. On the left side it comes off early on the dorsal aspect of a bronchus that continues on as the apical. As this segment is recognized by radiologists as a site of predilection of tuberculosis, its importance might well be stressed by a more complete territorial designation. To accomplish this would involve calling it the mediastinal-paravertebral-axillary. This is obviously unsuitable. The term subapical is suggested, as the segment corresponds in distribution with the one we have similarly named in the lower lobe.

Middle Lobe

In the middle lobe the choice of terms lies between anterior and pectoral. One would like to pay deference to Ewart's classical and monumental work and call it pectoral, but anterior is suggested on account of its universal application.

Lower Lobe

In the lower lobe "apical" is obviously a convenient term, although Nelson used the term "dorsal" and again pointed out that it is at times a separate lobe. "Cardiac" is a term used by students of comparative and human anatomy, although the segment has no exclusive relation

to the heart. The segment occurs only on the right side. In the left it is absent, the corresponding area of the lung being served by the anterior basal. In the koala, or Australian native bear, this segment is incorporated in the right lung as in the human, but in all the other animals that we have investigated it is a separate lobe.

The term "base" is used by anatomists when referring to the diaphragmatic surface of the lung. Ewart refers to the base of the lung as "the broad lower extremity of the lung including its inferior third," although he recognizes that part of the middle lobe in the ventral or anterior aspect of the lung would come within these limits. Practically all the lung territory served by the anterior basal (anterior division), the axillary basal, and the posterior basal segmental bronchi is contained in this so-called "base." It is a convenient term, practical and ingrained in the literature and clinical description. It is consequently worthy of retention, and substitution is difficult.

The terms anterior basal (although this has an anterior and a lateral division, each of which at times has a separate bronchial opening), axillary basal, and posterior basal describe the location of the segments with exactitude and are acceptable in that they resemble Ewart's original designations. Our findings must necessarily be in general agreement with those detailed by Ewart and more conveniently represented by Kramer and Glass, and Nelson, but we find in the subapical segmental bronchus or bronchi of the lower lobe a probable source of confusion to anatomists and also to bronchoscopists.

Between the origin of the apical bronchus of the lower lobe and the origin of the axillary basal from the main bronchus are found one or more openings. Before our recent access to Ewart's book we thought that we were the first to describe them, as they were not figured by Kramer and Glass, and Nelson. However, Aeby depicts one on each side of the tree in the position we have mentioned, and Ewart refers to it as the lesser posterior horizontal, a single bronchus on the right side, taking origin between the anterior basal and axillary basal. On the left side he refers to it as a branch of the axillary basal.

In our specimens we have found from one to three such subapical bronchi on the right side, and one on the left, quite apart from the other segmental bronchial openings mentioned. In a relatively simple type of lung, such as that of a pig (Plate, Fig. A), distal or caudal to the apical bronchus of the lower lobe may be seen a series of branches of descending magnitude on the dorsal, lateral, ventral, and, in lesser numbers, on the mesial aspects. Homologues of some of these may be found in the monkey, cat, sheep, dog, opossum, rabbit, kangaroo, wombat, koala, and duiker deer. Each mammal develops a characteristic architecture of its bronchial tree. Some bronchi individualize themselves in accordance with the pattern of the tree of the species, whilst others remain vestigial or disappear.

With a knowledge of the positions of the segmental bronchi in man, one can readily recognize the homologues in mammals. Distal to the apical bronchus of the lower lobe we found in all our mammalian specimens homologues of the subapical bronchus or bronchi. In the human we have observed it arising from the dorsal, mesial, or lateral aspect of the right main bronchus. We show an illustration (Plate, Fig. B) of the right bronchial stem in a child with a subapical bronchus in each of these three positions. On the left side (Plate, Fig. C) we have found it situated dorsally, mesially, laterally, and twice ventrally. It may be absent on the left side, in which case we have found a branch going to its customary distri-

RIGHT LUNG, ANTERIOR VIEW

- UPPER LOBE
1 SUBAPICAL
2 APICAL
3 AXILLARY
4 ANTERIOR
- MIDDLE LOBE
5 ANTERIOR
6 AXILLARY
- LOWER LOBE
7 APICAL
8 CARDIAC
9 SUBAPICAL
10 ANTERIOR BASAL
11 AXILLARY BASAL
12 POSTERIOR BASAL

RIGHT LUNG, POSTERIOR VIEW

- UPPER LOBE
1 SUBAPICAL
2 APICAL
3 AXILLARY
4 ANTERIOR
- MIDDLE LOBE
5 ANTERIOR
6 AXILLARY
- LOWER LOBE
7 APICAL
8 CARDIAC
9 SUBAPICAL
10 ANTERIOR BASAL
11 AXILLARY BASAL
12 POSTERIOR BASAL

FIGS. 1 and 2.—Diagrammatic drawing of the broncho-pulmonary segments and the beginning of the segmental bronchi.

RIGHT LUNG, LATERAL VIEW

RIGHT LUNG, MEDIASTINAL VIEW

FIGS. 3 and 4.—Broncho-pulmonary segments. (If these diagrams are enlarged by a pantograph on to sheets of cellophane so that the transverse diameter at the level of the seventh dorsal vertebra is about seven and a half inches, superimposition on a standard lateral film will allow the broncho-pulmonary segments to be delineated.)

bution arising from the axillary basal (described by Ewart as being its only origin) or from the posterior basal. As stated above, it is additional to all other constant bronchi in the lungs of man and of the animals we have examined. The reason for its occasional absence from the common site in the left lung of man and its irregular appearance within the limits mentioned is not at present clear. It is possibly becoming vestigial in man, as its variations would suggest. We have named it the subapical, and in metal casts and by single bronchial fillings with lipiodol and by inflation have shown its distribution to be to the mesial, posterior, and lateral areas of the lower lobe between the apical and posterior basal segments.

Commentary

With the realization of its varied location and the recognition that the other segmental bronchi are constants, the bronchial tree can be described with the surety of ordinary anatomical description.

This dissertation cannot be suggested as detracting from the work of Ewart, who must have devoted many years to his investigations. Alone, and with crude instruments and materials, he described fully the bronchial tree even to its minute branches. Increasing experience of later investigators has confirmed the general accuracy of this master of bronchial anatomy.

BIBLIOGRAPHY

- Aeby, C. T. (1880). *Der Bronchialbaum der Säugethiere und des Menschen*, Leipzig. (Quoted by Ewart.)
 Ewart, William (1889). *The Bronchi and Pulmonary Vessels*, J. and A. Churchill, London.
 Kramer, Rudolph, and Glass, Amiel (1932). *Ann. Otol.*, **41**, 1210.
 Neil, J. Hardie, Gilmour, W., and Main, W. W. (1936). *Proceedings of Second Congress, Pan-Pacific Surgical Association*.
 — Gwynne, F. J., and Gilmour, W. (1937). *Med. J. Austral.*, **2**, 165.
 — — — Main, W. W., and Fairclough, W. A. (1937). *Ann. Otol.*, **46**, 338.
 — — — — — (1938). *Austral. New Zeal. J. Surg.*, **8**, 118.
 Nelson, H. P. (1934). *British Medical Journal*, **2**, 251.

The eighth annual report of the National Birth Control Association (president, Lord Horder), with which is now incorporated the Birth Control International Information Centre, the Birth Control Investigation Committee, and the Society for the Provision of Birth Control Clinics, points out that the amalgamation last spring has left the N.B.C.A. as the only central birth control organization apart from the Society for Constructive Birth Control. The chairman and secretary gave evidence before the Interdepartmental Committee on Abortion, a memorandum having been previously sent in. The medical subcommittee, under the chairmanship of Mr. A. C. Palmer, completed another year of active work, and among leaflets issued was one entitled "Services to Private Practitioners by the Medical Subcommittee of the N.B.C.A." A Scottish Advisory Committee was formed in 1938 and a number of new branches were set up in England and Scotland; some of them have already opened clinics. Early last spring a move was made to new and larger headquarters at 69, Eccleston Square, London, S.W.1. A conference of all branches, held in London before the annual meeting, was so successful that this will now be an annual event. Out of the 409 maternity and child welfare authorities in England and Wales 246 have now taken action of one sort or another under the Ministry of Health memoranda. Of these 246, seventy-nine now have post-natal gynaecological or birth control clinics; 113 refer cases to voluntary or municipal clinics, to hospitals, or to private practitioners; and thirty-eight make grants or lend premises, or do both, to voluntary clinics. Some take action in more than one of these ways.

TWO CASES OF PICK'S DISEASE STUDIED WITH THE AID OF KYMOGRAPHY

BY

G. L. S. KONSTAM, M.D., M.R.C.P.

*Physician to the West London Hospital, with Charge of the
Cardiographic Department*

AND

FRANKLIN G. WOOD, M.B., D.M.R.E.

*Radiologist to the London Chest Hospital and the
German Hospital, Dalston*

(WITH SPECIAL PLATE)

The following two cases of Pick's disease (chronic constrictive pericarditis) were considered worthy of publication, not only because of their comparative rarity, but because opportunity was afforded of studying them by kymography. The first case is of particular interest, as kymograms were taken before and after operation.

Our knowledge of this syndrome and its treatment by pericardial resection has been considerably extended by Paul D. White (1935), whose St. Cyres Lecture summarized his study of fifteen personal cases. The salient feature met with is fibrous or callous thickening of the pericardium, which by gripping the heart in an unyielding envelope prevents normal diastolic filling and leads to what has been termed "inflow stasis." The syndrome occurs in both sexes, and its onset is more usual in early than in middle life. The aetiology is often obscure, but there may be a past history of pneumonia, tuberculosis, polyserositis, pleurisy, and acute pericarditis. It is noteworthy that in none of White's series was there any evidence of a rheumatic infection.

The main symptoms are dyspnoea and swelling of the abdomen; the signs are similar to those observed in chronic congestive heart failure, with the important exceptions that the heart is not usually enlarged and evidence of valvular disease is absent. The jugular veins, and often the arm veins, are greatly engorged, the liver is enlarged but not tender, and ascites develops which is out of all proportion to any oedema that may be present in the extremities. The blood pressure and pulse pressure are below normal, and pulsus paradoxus is often found. Normal rhythm is usual, but in four of White's cases, and in one here described, there was auricular fibrillation.

X-ray Appearances

The heart shadow is normal in size and shape, or moderately enlarged. Calcification of the pericardium is frequently present, but its absence does not exclude the diagnosis; congestion of the pulmonary vessels and enlargement of the supracardiac shadow are also met with. On screen examination the normal heart pulsations are greatly reduced, and may even be absent, but these findings are more accurately shown by the modern method of kymography. An electrocardiogram may show low-voltage flattening or inversion of the T waves in Lead I or II, or in both leads. When inverted the T wave is coronary in Type. Low-voltage QRS waves are also commonly found.

Without surgical treatment the prognosis is hopeless, the disease usually terminating fatally within a few years. Resection of the pericardium by operation may, on the other hand, lead to restoration of normal health.

Case 1: Pick's Disease of Unknown Origin

A girl aged 9 attended the out-patient department at the West London Hospital in February, 1932, to 1933 suffering

FIG. A.—Cross-section of left bronchial stem of a pig, at the level just above the homologue of the subapical bronchus.

FIG. B.—A pathological specimen from a child aged 4 who died of bronchopneumonia, lung abscess, and empyema following measles. There are shown the orifices of three subapical segmental bronchi which are the variants of the bronchial tree.

FIG. C.—Showing the constants of the left bronchial tree and a large subapical bronchial orifice.

C. C. HOLMAN: PROGRESSIVE STREPTOCOCCAL
ULCERATION

CASE II.—Condition on November 1, 1938.