SURGICAL PROCEDURES IN GENERAL PRACTICE

This is one of a series of articles contributed by invitation

PRURITUS ANI

BY

A. LAWRENCE ABEL, M.S., F.R.C.S.

By the term "pruritus ani" is literally meant "an itching at the anus." *Gould's Medical Dictionary* defines pruritus as "itching, a peculiar uncomfortable sensation due to irritation of a peripheral sensory nerve and leading to scratching." As itching cannot occur without some causative factor pruritus must be regarded as a symptom rather than as a disease in itself. Nevertheless it is true that definite changes in the anal and peri-anal skin occur in the vast majority of those who complain of irritation, and it is reasonable to apply the term "pruritus ani" only to cases in which these skin changes are present. If the underlying cause or causes—for they are often multiple—are correctly diagnosed and adequately dealt with suitable local treatment to the affected skin is likely to meet with complete success. Conversely, if the factors which have produced the irritation are inadequately dealt with no amount of local treatment is likely to effect a permanent and complete cure.

For practical purposes the simplest classification of the causes of pruritus ani would appear to be as follows: (1) excessive moisture; (2) functional (that is, hysterical or psychological); (3) infective.

Excessive Moisture at the Anus

This condition may arise from within the bowel or from the skin itself. In the former case seepage of mucus from the mucosa of the rectum and anal canal is the commonest cause, but pus, blood, or liquid paraffin may also seep past the sphincters. In the latter case the condition is due to lack of cleanliness, excessive sweating, local irritation by clothing, or a specific disease of the skin.

CAUSES WITHIN THE BOWEL

Any abnormality of the rectum or anal canal tends to give rise to a hyperactivity of the mucous membrane and an excessive secretion of mucus, larger or smaller quantities of which, leaking past the sphincters, cause sogginess of the anal and peri-anal skin. Even minute quantities may produce the typical changes of pruritus ani. The skin typically becomes reddened, the rugae enlarge from oedema, and superficial erosions often occur. As the condition becomes more chronic thickening and induration arise from cutaneous and subcutaneous fibrosis. These changes in turn are followed by fissuring and cracking of the epithelium, accompanied by pain, and infection by intestinal and other organisms, which still further increase the local dermatitis. The commonest diagnosable and treatable conditions in the rectum are as follows:

Rectal Constipation.—Careful examination will often reveal an imperfectly emptied rectum, even after what appears to the patient to be a satisfactory daily evacuation of the bowel. Similarly, in patients who have two or more evacuations of the bowel, careful examination will often reveal that the rectum has only been partly emptied at the first action and is not completely emptied at the second or even the third. The presence of a faecal

mass resting like a bung on the pile-bearing area of the rectum will in a large proportion of cases give rise to enough local congestion and hyperaemia to produce a slight excess of mucus, which may seep through the anal sphincters and reach and irritate the external skin. Another effect of this type of constipation is to cause flatus to be voided frequently, and this will further allow the passage of mucus and tends to slight soiling around the anus.

Pectenosis.—Passive congestion, with varying degrees of sphincteric spasm, may eventually give rise to a deposition of fibrous tissue beneath the mucous membrane of the anal canal in its middle portion, which is known to anatomists as the pecten. This band of fibrous tissue, named by W. Ernest Miles the "pecten band," is another common abnormality in cases of pruritus ani, and the condition is known as pectenosis.

Internal haemorrhoids in early, intermediate, or late stages of development are common accompanying factors. A fissure, prolapse of the rectal mucosa, hypertrophied anal papillae, inflammation of the Morgagnian crypts, and blind internal fistulae may, severally or in multiple, be the underlying causes of the condition. Occasionally the excess of mucus comes from a simple or malignant tumour of the rectum or anal canal. Many of these factors are brought into play or have their effects magnified by too rich a diet and by alcohol. This is especially true in those classes of society which can afford them. Rare cases are found to have an achlorhydria, and Swift believes that in these there is a diminished absorption of vitamin A from the dietary. This lack of vitamin A results in deficient nutrition of epithelium. Similarly, a lack of absorption of vitamin B may give rise to altered sensation of nerve endings. Difficult cases should accordingly bring into mind the possibility of avitaminosis. I have only once known tobacco to be the most important determining factor, and in this case the unfortunate individual was a medical man.

MOISTURE OF THE SKIN

The commonest cause under this heading is lack of cleanliness. The relatively more dirty habit of using paper, which is common throughout the "civilized" races, instead of washing after defaecation is responsible for a large number of the milder grades of pruritus ani, especially in more hairy individuals. A thorough wash night and morning in all adults is obviously the counsel of perfection. Excess of sweating, especially in the more arduous occupations, or in times of stress and worry in some individuals, is often sufficient to produce the typical skin changes.

Thick woollen underclothing may be responsible for or may aggravate pruritus. Those of us who have had the privilege of seeing a large number of our fellow-countrymen let down their nether garments, and who have taken the opportunity to observe the same, are aware that the British housewife loves to do her most thorough darning in the area which is the seat of pruritus ani. These patients often improve considerably if their underwear is made of pure silk or at any rate of artificial silk, or, failing that, of linen mesh. Occasionally the peri-anal affection is associated with a generalized derma-

titis, a gouty diathesis, or with glycosuria. It frequently accompanies pruritus vulvae, and then usually has its origin in the field of gynaecology. Skin tags around the anus increase the likelihood of lack of cleanliness and excess of sweating, and often have to be removed before much headway can be made with local treatment.

Psychological Causes

Although it is true that there is almost always a peculiar nervous temperament, with an inclination to easy worry, in a very large majority of those who suffer from pruritus ani, it would appear that many cases of the more serious types of pruritus are entirely due to an abnormal mental make-up. In these cases associated hysterical manifestations, such as loss of corneal reflex, anaesthesia of the palate, patchy anaesthesia of the face and legs, grossly exaggerated knee-jerks, are often present. Worry is a factor both before and after the development of the condition. For obvious reasons this type of case is the most difficult to treat. The condition may be compared to the clinical entity of dermatitis of the vulva with cervical erosion in highly strung neurotic women.

Infective Causes

The most frequent single infective factor, especially in children, is the common threadworm. One or more of these may sometimes be seen wriggling at the anus, or a proctoscope may reveal their presence in the rectum. The threadworm tends to come into the rectum and anal canal towards evening, having migrated there from higher up. Doubtful cases should therefore always be examined in the latter part of the day. It is my practice at the Gordon Hospital to subject all patients, where indicated, to a high colon wash-out, specifically to exclude the presence of *Enterobius vermicularis*. The next most common parasite is the *Pediculus pubis*. Careful inspection will reveal the presence of the parasite and its ova not only around the anus but also in the pubic region. Scabies may also affect this region and give the typical appearance, and may extend to other parts of the body.

Very rarely the cause of a local infection is a fungus or yeast. In these cases the affected areas of skin have a circumscribed and well-defined border, and microscopical examination of a suspected case will reveal the presence of either *Monilia albicans* or epidermophyton or trichophyton, and clinch the diagnosis.

Symptomatology

The most prominent symptom is itching. This may be anything from a mild irritation lasting for a few seconds after the patient gets warm in bed, or after the bowels have moved, up to an intolerable irritation, which may be more or less continuous, accompanied by burning and smarting and with soreness due to the excoriations produced by scratching. The patient may be able to control the itching by pinching or squeezing the affected part, but often he wakes at night to find himself in the act of scratching, and his sleep may be seriously disturbed, to the point of affecting his health.

A careful case history must be taken of all patients, and examination must be made in a good light; it should consist of external inspection and palpation, and digital and endoscopic exploration of the anal canal and rectum. Careful examination of the abdomen and pelvis should also be made. The urine must always be examined for sugar, and the patient's constitutional state and mental make-up taken carefully into account. Never rely upon general causes for pruritus ani, except when the most

minute investigation has failed to reveal any local source of irritation.

The changes seen in a typical case have been detailed above; the skin may be macerated and moist, or shiny and dry, with denuded epithelium; there may be associated scratch marks with linear pigmentation and decapitated follicles. Pyogenic infection may have supervened, with pustules and even boils. In severe cases these are concomitant with the pruritus and are the result of scratching and superadded infection.

The treatment of cases of pruritus ani will be described under the following headings: hygienic; dietetic; medicinal; local treatment—(a) directed to the cause (operative and non-operative), (b) directed to the skin (by surface applications, by subcutaneous injection, and by operation).

Hygiene and Diet

As already stated, silk or artificial silk underclothing should be substituted for wool or flannel. Careful cleanliness is most important. Daily, or preferably twice-daily, washing is advised for all adults. Hairy individuals occasionally need peri-anal shaving until the skin becomes normal.

Depending on the severity of the condition, alcohol, coffee, tea, and sugar should be limited or even forbidden. Meat should be restricted, and shell-fish, tinned food, pickles, spices, and all highly seasoned foodstuffs should be avoided.

Medicinal Treatment

As already indicated, careful attention must be paid to the state of the bowels. The ideal at which to aim is a regular, smooth, hundred-per-cent. daily action, which should take place shortly after the patient has broken his fast. Thereafter the rectum should be completely empty for the remainder of the twenty-four hours. The drug that appears to have the best effect upon the last few inches of the large intestine is senna. The most satisfactory method of administering this is as follows:

R.	Pulv. fol. sennae ...	...	...	...	} āā 3 grains
	Pulv. rad. jalapae ...	...	...	...	
	Pulv. sulph. sublim. ...	...	...	...	
	Theriacae nigrae ...	...	...	ad 1 drachm	

Fiat confectio. Mitte 4 ounces.

Sig.—One drachm nocte.

The patient should be instructed to start by taking a small eggspoonful of this electuary regularly at night-time, varying the quantity with the effect produced. After two or three days the patient will find that he gets a regular, soft, and complete evacuation of the bowel each morning; and its use should be continued until considerable progress with the remainder of the treatment has been made. It will be found that after two or three weeks the nightly dose can be stopped for one night and still the bowels will work the following morning. The next night the patient should take his dose. He may then miss two nights, still with two daily evacuations. In other words, this preparation is not what is commonly known as "habit-forming," but it actually does get the bowel into a good habit, which is continued after the medicament is stopped. In rare cases the complete senna leaf is found to cause griping, and an alternative way of giving senna is as an infusion. Eight or ten leaves are soaked in cold water for not more than three hours, and the resulting "tea" taken about 10 p.m. If cold water is used and not hot, and not more than three hours are allowed for soaking, the griping element of senna is not obtained, and a satisfactory action of the bowels occurs.

The following evening a smaller or larger number of leaves are infused, until the appropriate number to suit that particular individual is found. If liquid paraffin is used it must not be given in doses large enough to cause seepage, as this in itself is a potent cause of irritation, which outweighs its benefit as a lubricant. Any aperient that produces numerous loose watery stools is to be avoided. When the itching disturbs the night's sleep, allonal, luminal, bromide, valerian, phenacetin, and veganin are valuable as sedatives. Achlorhydria or avitaminosis should receive suitable treatment.

Local Measures

A number of the underlying causes of pruritus ani can best be dealt with by a radical operation. Cases of pectenosis should be treated by pectenotomy—that is, division of the pecten band. Internal haemorrhoids in the third degree or stage of their development and the still more advanced intero-external (or van Buren) piles require the modern ligature operation; some cases of chronic fissure and all fistulae, and of course any neoplastic disease, must undergo radical surgery; and skin tags causing irregularities of the surface should be removed.

Injection of Sclerosing Fluid

The early and intermediate stages of internal haemorrhoids may satisfactorily be dealt with by the injection of a sclerosing fluid. The one most commonly in use is 5 per cent. phenol in almond oil. This is injected beneath the mucous membrane high up in the pile. The routine treatment at the Gordon Hospital is to inject a large quantity, up to but not exceeding 10 c.cm., into one pile only at the first sitting. Approximately one week is allowed to elapse before the next injection, and each of the three primary piles is treated in turn. When all three have been dealt with, an interval of three to four weeks is allowed to elapse and the condition reviewed to ascertain if further injections are necessary. The larger variety of piles will require two, or occasionally three, applications. The second type of injection fluid which is sometimes required for the lower portion of the pile is from 5 to 7 minims only of a 20 per cent. solution of phenol in equal parts of glycerin and water. This is given deeply into the centre of the pile mass itself.

The effect of any sclerosing injection may be augmented by the use of an astringent ointment, the most efficacious of which is made up as follows:

R_x Unguentum hydrarg. subchlor. B.P. ... ½ ounce
Sterile white vaselin ... ad 2 ounces
Fiat ung. Mitte 2 ounces.

The patient is instructed to place a little of this ointment on his finger and gently to milk it into the rectum before and after each action of the bowels, and again before going to bed, throughout the course of his injection treatment. Immediately the injections and the use of the ointment are started the quantity of mucus produced by the pile-bearing area is diminished and the irritation begins to subside.

In cases of anal fissure, injections may again be employed with considerable benefit. In this case an injection containing an anaesthetic with a prolonged effect is given with strict aseptic precautions into the substance of the external sphincter muscle. B.A.B.A. with percaine, or proctocaine, is most commonly used, and from 5 to 10 c.cm. may be given in a fan-shaped manner on one or several occasions. Its effect is augmented by the use of an astringent anaesthetic and antiseptic ointment, as,

for example, the aforementioned ointment containing 10 grains of percaine hydrochloride to each 2 ounces.

In those cases where threadworms are a cause of the condition treatment should consist of a rectal injection of 4 ounces of a salt-and-water solution, using two teaspoonfuls of salt to 4 ounces of water. The injection should be given in the evening with an ordinary bulb or Higginson enema syringe. The cure may be accelerated by the use of either santonin 2 grains with 1 grain of calomel each night for three nights, or one tablet of butolan three times a day for a week. Half these doses should be given to children. Great personal cleanliness is important, especially the washing of the hands after defaecation in order that reinfection may not continually occur.

Scabies quickly responds to the use of sulphur ointment, and should disappear if the underclothing is changed and cleaned night and morning for three days. Pediculosis is cured by shaving and the use of a mercurial ointment. Fungus infections are best treated with frequent swabbings of the affected parts with 1 per cent. solution of methyl or gentian-violet in 25 per cent. spirit, followed by the application of a good dusting powder.

1. Surface Applications to Skin

In cases where an acute dermatitis is present Miles finds the following ointment of value:

R_x Pulv. zinc. ox. ... 2 drachms
Lin. camph. ... 3 drachms
Anaesthesin ... 2 drachms
Lanolin ... 1 ounce
Vaselin ... ad 3 ounces

Fiat ung. Mitte 3 ounces.

Sig.—To be smeared on the affected part two or three times by day and on going to rest.

For local application by the patient when the dermatitis is less acute, Gabriel has found the following lotion very efficacious:

R_x Phenol ... 1 drachm
Zinc oxide ... 2 drachms
Pulv. calamin. prep. ... 1 drachm
Glycerin ... 2 drachms
Spirit. rect. ... 2 drachms
Aqua rosae ... ½ ounce
Milk of magnesia ... ad 4 ounces

Sig.—To be dabbed on the anal region morning and evening; if necessary a compress moistened with the lotion may be applied during the night.

I have found the following treatment of great help in many cases, especially where superficial fissures and local soreness exist. The affected skin is painted with a lotion consisting of 20 per cent. phenol in equal parts of glycerin and water. After 15 or 20 seconds the excess of this lotion is removed by gently dabbing with cotton-wool, and the following ointment then applied:

R_x Pulv. zinc. ox. ... 2 drachms
Lin. camph. ... 3 drachms
Percaine hydrochloride ... 10 grains
Sterile white vaselin ... ad 2 ounces
Fiat ung. Mitte 2 ounces.

Sig.—To be applied to the affected part two or three times a day and on going to bed.

The 20 per cent. phenol in glycerin and water solution should be applied by the medical attendant only, and may be repeated in from three to seven days, depending on the severity of the case.

Other types of ointment that have been found efficacious have been percainal and nestosyl: the latter may be combined with a nestosyl suppository passed into the anal canal before going to bed. In those cases where oedema of the skin around the anus occurs, Miles finds that a

lotion of lead acetate—prepared by mixing one drachm of liquor plumbi subacetati with two tablespoonfuls of fresh milk—gives great relief. Where an unhealthy surface epithelium exists superficial layers of the dermis may be destroyed by using a saturated solution of silver nitrate applied to the surface with a camel-hair brush and after ten minutes neutralized by a normal solution of sodium chloride. This may be applied twice a week, and will stimulate the superficial excoriations to heal. At a further stage a dusting powder containing zinc and starch, or zinc starch and boric acid, may be used by day and the lotion or ointment by night.

2. Subcutaneous Injection

It occasionally happens in the most severe grades of pruritus ani, and especially in the so-called functional type of case, that local applications to the surface have little or no beneficial effect, and recourse has then often to be had to the subcutaneous injection of oil-soluble anaesthetics or sclerosing solutions. In these severe cases local anaesthesia of the irritating area can be produced, with much benefit to the patient's state of mind, a relief from scratching, and a healing of the local excoriations and dermatitis. The mere prick with a needle appears often to improve the patient's psychology. A number of oil-soluble anaesthetics are available. The most popular are Gabriel's percaïne in oil, Gorsch's anucaïne, B.A.B.A. of Martindale, and Morgan's proctocaine. B.A.B.A. with the addition of 1 per cent. percaïne also gives good results. With strict aseptic precautions a large injection of the solution, varying from 10 to 30 c.cm., is given by means of a long needle, through healthy skin, in a fan-like manner deeply into the peri-anal region. A finger in the anal canal makes sure that the latter is never punctured. Gabriel recommends, and I believe wisely, that the treatment should be carried out in a hospital or nursing home, where careful attention can be paid to the patient's sleep, the regulation of the bowels, and local treatment by surface applications.

Haynes advises the use of from 10 to 20 c.cm. of a sclerosing solution of 1 in 2,000 hydrochloric acid. Through a single posterior puncture a four-inch needle is passed, and a sterile solution of 1 in 1,000 percaïne is injected into the subcutaneous tissue, using from 10 to 15 c.cm. The syringe is then loaded with the hydrochloric acid solution, and as the needle is withdrawn this solution is injected so that it is evenly distributed in the same plane as the percaïne. Other workers have used alcohol. Stone gives two or three minims of absolute alcohol through multiple small punctures into the subcutaneous tissue. Buey injects 40 per cent. ethyl alcohol through a long (four-inch) needle from a single posterior puncture, using from 20 to 40 c.cm. The risk of sloughing is considerable in both these methods. For the severe, intractable type of case, in which no local pathology can be found to account for the condition, large doses of the oil-soluble anaesthetics appear to me to give the best results.

Provided every other aspect of the patient's treatment is being or has been carefully carried out, a small dose of x rays (half an erythema dose) sometimes greatly benefits the typical intractable case.

3. Operation

Surgical procedures designed to divide the peripheral nerves and undercut the anal and peri-anal skin are only mentioned in order to condemn them; the results are so unsatisfactory that they are seldom if ever worth undertaking.

SURGERY OF THE GALL-BLADDER

PROFESSOR GREY TURNER'S THIRD LECTURE

The third and last of Professor Grey Turner's Lettsomian Lectures on the surgery of the gall-bladder and common bile duct was delivered before the Medical Society of London on March 6. In speaking of the preparation of the patient for operation Professor Turner said that it was the custom to think mostly in terms of the jaundiced patient, but any operation on the gall-bladder was potentially a serious undertaking. The late Lord Moynihan said: "Surgery has been made safe for the patient; let us make the patient safe for surgery." While he could not entirely subscribe to the first part of that aphorism, everyone would agree as to the wisdom of the second. Before any operation on the gall-bladder the patient should be in hospital and receive a certain preparation, the amount and type depending on circumstances. Where any signs of obesity or of trouble from intestinal conditions were evident it was a good thing to get the patient accustomed to the passage of a stomach tube before the operation; this might be valuable in after-treatment, though it was a trying ordeal for the unaccustomed patient.

Preparation of the Patient

It was just thirty-seven years ago in that very room that Mayo Robson laid great stress on the extreme importance of the preparation of the jaundiced patient (his address was reprinted in the *British Medical Journal* of January 18, 1902). Mayo Robson gave calcium chloride by the mouth and rectum. The lecturer saw it so used, but was never quite satisfied that given in that way it was really of great value, but it did point to the necessity for doing something to overcome the risks associated with haemorrhage. It behoved them in every case to have that complication in view. A great many of these jaundiced patients had suffered much pain and had had morphine, and it was important to keep them under observation for some time for the effect of drugs of that kind to be overcome. It should be seen that they had sufficient fluid (not forgetting the natural rather than the unnatural channels), and it was well to pay attention to the condition of the bowel. He believed that the intravenous exhibition of calcium chloride was certainly of value. The use of blood transfusion before or after operation was a recourse which should always be borne in mind. Whether it should be used as a routine was a moot point, but when there was evidence of a haemorrhagic tendency it should never be omitted. It might be wiser in such cases to operate in two stages. In his experience the use of ascorbic acid (vitamin C) had been of some value in particular cases, and he had great hopes of the help to be obtained from vitamin K, which seemed to have striking anti-haemorrhagic properties.

Technique of Operation

Coming to the technique of the operation Professor Turner spoke first of the anaesthetic. He did not believe that it was always necessary to use spinal anaesthesia to get the best results. Local anaesthesia was perhaps contra-indicated in deep jaundice because the actual trauma of the injection provided avenues from which haemorrhages might extend and become serious. The serious haemorrhage that sometimes took away the resistance of these patients was not always in the peritoneal cavity; he had evacuated from the abdominal wall on more than one occasion large masses of blood clot.

Whatever was to be done in the surgery of the upper abdomen, it was important not to be hampered by the want of proper exposure. All that was requisite might perhaps be done with one or other of two incisions—the vertical incision through the rectus muscle associated with the name of Mayo Robson or the oblique incision below the costal margin, which offered certain advantages. If the vertical incision was used he would plead for its upward extension. Surgeons must have been hampered when the vertical