obtainable in London; and to the Bursars of certain of the Colleges at Oxford and Cambridge for their courteous verification of data relating to certain of the oarsmen concerned.

REFERENCES

- Cooper, E. L., O'Sullivan, J., and Hughes, E. (1937). *Med. J. Austral.*, 1, 577.
 Drinkwater, G. C., and Sanders, T. R. B. (1929). *University Boat Race Official Centenary History, 1829-1929*, Cassell, London.
 Dublin, L. I. (1928). *Harper's Monthly Magazine*, 157, 229.
 — (1930). *New York Times*, July 20, Special Features Section.
 — (1932). *Statistical Bulletin, Metropolitan Life Insurance Co.*, August.
 Fairbairn, S. (1931). *Fairbairn of Jesus*, p. 69, London.
 Greenway, J. C., and Hiscock, I. V. (1926). *Yale Alumni Weekly*, June 11.
 Lehmann, R. C. (1931). *The Complete Oarsman*, 4th ed., London.
 Lewis, Sir T. (1937). *Diseases of the Heart*, 2nd ed., p. 154, London.
 Meylan, G. L. (1903-4). *Harvard Graduates Magazine*, 12, 362, 367, 543.
 Morgan, J. E. (1873a). *University Oars*, p. 300, London.
 — (1873b). *Ibid.*, p. xiii.
 — (1873c). *Ibid.*, p. 20.

CHRONIC URTICARIA *

BY

NORMAN BURGESS, M.A., M.D., M.R.C.P.

Physician in Charge of the Skin Department, Bristol General Hospital; Clinical Lecturer in Diseases of the Skin, University of Bristol

The subject of chronic urticaria is one of great interest, first because its complex nature, and frequently its long duration, set a difficult problem, and, secondly, because perhaps more than any other skin condition, it brings us in close touch with general medicine. I have purposely avoided the acute cases, whether of external or internal origin, as these rarely cause any difficulty, and I am restricting this discussion to those cases of urticaria of internal origin which are already of long standing when we first see them. With regard to the final mechanism responsible for wheal formation, it is now generally believed to be due to the release of relatively large amounts of H-substance as a result of cell damage; this may be caused in several ways, of which antigen-antibody reaction is one. So far as possible I propose to consider the subject from the clinical point of view, and to base my remarks on cases which have been under my care, but it would perhaps be wise, before proceeding to this clinical study, to discuss some of the theories that have been put forward to account for the production of chronic urticaria.

Theories of Causation

It is thought that the toxin responsible for the condition may be absorbed from foci of infection in the teeth, nose, throat, lungs, gall-bladder, or urinary tract. In other cases the patient suffers from urticaria when exposed to cold, and this has been shown to be due to the presence of a dermolysin in the blood which unites with the cells at a low temperature and lyses them on rewarming, setting free H-substance. Another cause of urticaria, probably more common than one might imagine, is emotional upset. In this connexion an interesting theory has been put forward by Stokes and Pillsbury (1930), who believe that as a result of nervous strain there is a diminished secretion of hydrochloric acid in the stomach, leading to an increase in the Welch bacilli in the intestine and so to the production of a histamine-like substance, the absorption of which can give rise to the

triple response. As a further result of hypochlorhydria there is diminished absorption of calcium from the food; again, diminished pancreatic and intestinal secretion leads to imperfect digestion of proteins with absorption of abnormal cleavage-products of protein from the alimentary tract, and because of this the patient may become sensitized to food and bacterial proteins. Another view is that the release of histamine as a result of antigen-antibody reaction is accompanied by emotional upset, the frequent repetition of this leading to a conditioned reflex. Thus the emotion can then itself release histamine even in the absence of the allergen.

It is also believed that the urticarial mechanism can be set in action by the products of damaged epidermal cells, and Lewis showed that a needle-prick in the epidermis can give rise to a vascular response. It is probable, however, that in perhaps the majority of cases the toxin is absorbed from the alimentary tract. In the first place the condition may be caused by allergy due to specific food sensitiveness. It has been shown by A. and M. Walzer (1927, 1928) that the protein may be absorbed unchanged into the blood stream and, reaching sensitized tissue, produce an antigen-antibody reaction. This is especially likely to occur if there is hypochlorhydria, or if the proteopexic power of the liver fails. Secondly, abnormal split-products of protein derived from food, due to deficient digestion, or products of bacterial action on proteins, may be absorbed and, escaping destruction in the liver, produce the symptoms. This will take place if the detoxicating power of the liver is at fault.

Barber and Oriel (1928), as a result of extensive biochemical investigations of the blood and urine during and after allergic attacks, came to the conclusion that there is evidence of hepatic disturbance in these cases, and that a failure of the proteopexic and antitoxic power of the liver is to some extent responsible for the condition. Oriel (1931) also found that it was possible to precipitate a proteose-like substance by adding excess of alcohol to an ether extract of the urine. He found that the amount of this proteose was greatly increased during an attack of urticaria or angioneurotic oedema. He claimed that in certain allergic diseases, including urticaria, the proteose contains the specific hapten responsible for the condition, and that when it is injected intradermally it causes a reaction similar to that produced by the allergen to which the patient is sensitized. He also showed that urinary proteose can act as an antigen, and that sensitivity can be transferred by the injection of the serum of a sensitized patient. This was carried out by the Prausnitz-Kustner technique. He believes that proteose is sometimes the primary allergen, but that in many cases it acts as a secondary allergen in patients sensitized to an exogenous substance.

Another factor which is of importance in chronic urticaria is heredity, for it is well known that in certain families there is a predisposition to urticaria, asthma, hay-fever, eczema, Besnier's prurigo, and migraine. This inherited state has been described by Coca (1926) as "atopy." I do not think that it would be profitable, in the present state of our knowledge, to discuss the views advanced by Coca to explain this hereditary tendency. An interesting theory to account for familial predisposition to allergic conditions is that put forward by Langdon-Brown (1932), who suggests that this tendency is due to absence of the ferment histaminase, which destroys histamine as soon as it is released. This view is supported by the work of Barber and Oriel, who found an increase in the amino-acids in the blood and urine in

* Read at a meeting of the St. John's Hospital Dermatological Society, November 23, 1938.

patients suffering from allergic diseases. I have already briefly referred to this work, and shall consider it in greater detail when discussing the treatment of chronic urticaria.

Present Series of Cases

I will now proceed to a clinical study of cases which have been under my care during the past few years. In fifty-eight consecutive cases seen in private and hospital practice there has been a very pronounced sex incidence, only twelve cases occurring in males, against forty-six in females. If, as Stokes suggests, emotional factors play an important part in the production of this condition, the larger number of cases in women may perhaps be accounted for. Secondly, there is quite a marked age incidence, the greatest number of cases occurring in the fourth decade. I have set this out graphically in the accompanying chart. Forty-nine out of the fifty-eight cases

occur between 20 and 60 years, and there is a sharp rise in the curve between 30 and 40 years. This is probably due in part to the incidence of focal infection being greater during these periods. Also dyspepsia is more common in middle life, while emotional and nervous strain are more likely to be present at this time. The other possible aetiological factors have been examined in thirty-three consecutive private cases, in which the methods of treatment have also been studied.

The following were the aetiological factors in this group. An allergic family history was obtained in nine cases. Focal infection in teeth (fifteen) or in the nose and throat (eleven) was present in twenty-six cases, food sensitization was present in six cases, and dyspepsia in nine. Two patients were worse when exposed to cold, one case followed an attack of influenza, while six cases were secondary to eczema and were presumably caused by the products of damaged epidermal cells. In three cases a psychological factor was discovered; in one case the condition appeared to be associated with excessive doses of thyroid, while in two cases no cause was found. In many cases more than one factor was present. It will be seen that thirty of the cases showed abnormalities associated with the alimentary tract—that is, teeth, food sensitization, and dyspepsia.

In a disease of such complexity as chronic urticaria, in which several factors may be present in any given case, it is impossible in most instances to clear up the condition by a single therapeutic measure, and I feel that it is the failure of the practitioner to realize this which is to some extent responsible for the long duration of many of these cases. The routine I have adopted may be considered under three main headings: (a) treatment based on aetiological factors discovered by investigation, (b) treatment based on theoretical considerations, (c) desensitization.

Treatment Based on Aetiological Factors

In the first place the patient is examined for evidence of focal infection, and when present this is treated, if possible. The dramatic effect that can follow the elimination of infective foci is shown in the following cases.

Case 1.—A single woman aged 27 had suffered from urticaria daily for three months. The attacks always began between 5 and 6 p.m. Her teeth had recently been reported as healthy by a dental surgeon. An attempt to desensitize her by autohaemotherapy was unsuccessful. The teeth were then radiographed, and apical infection of two of them was discovered. Their removal was followed immediately by complete recovery.

Case 2.—A married woman aged 53 had a three-weeks history of daily attacks. Apical infection of a lower bicuspid was discovered by x-ray examination. Three days after the removal of this tooth the condition began to clear up and recovery was complete.

The patient is asked to keep a diet chart, and to record each day in parallel columns the food and drink taken, and the state of the eruption. By omitting various articles of diet in turn for a few days it is possible to discover specific food sensitiveness when present. In cases where imperfect digestion of proteins as a result of hypochlorhydria is suspected, the administration of hydrochloric acid with meals is often effective. It would of course be best in such cases to give fractional test meals, but in actual practice, especially in private cases, this may be difficult to arrange, and one is apt to rely on the therapeutic test. It is also sometimes possible to discover psychological difficulties which may be causal factors in the production of the condition. It may be a great surprise to the patient to find that her domestic difficulties and her skin eruption are in any way related, and the recognition of the association may lead to prompt recovery even if the domestic troubles cannot be remedied. This is well shown in Case 3.

Case 3.—A married woman aged 36 had suffered from urticaria for a year. She had much domestic unhappiness, which she felt compelled to conceal and repress. There was no possible satisfactory solution of her difficulties. The relation of this to her eruption was pointed out to her, an association which she had previously failed to realize. She was also ordered glucose and whole-blood injections. Her doctor reported that he had given her three injections, but that the condition had completely cleared up after the first one. In this case the rapid recovery must be attributed to psychotherapy rather than to any physical treatment.

In cases secondary to eczema, in addition to desensitization, to which I shall refer later, the original eczematous condition must of course be treated.

Case 4.—A married woman aged 57 had suffered from generalized seborrhoeic dermatitis for three weeks. Urticarial attacks had begun shortly after the onset of the dermatitis. She stated that she had suffered from eczema and nettle rash some years previously. She was treated with local applications and whole-blood injections. After two injections she was quite free from skin disease, with the exception of dandruff.

Treatment Based on Theoretical Considerations

The second part of the treatment is based on the observations of Barber and Oriel, to which I have already briefly referred, on the changes in the blood and urine during and after allergic attacks. I am inclined to think that this work has to some extent unfortunately been overlooked as a result of the subsequent controversy as

to the specificity of the proteose which Oriel isolated from the urine, and which was said to contain the specific hapten responsible for the symptoms. In their original paper Barber and Oriel showed that just before and during an allergic attack there is a rise in the amino-acids and non-protein nitrogen and a fall in the total chlorides in the blood, while the urine shows a rise in free acidity, ammonia excretion, excretion of amino-acids and non-protein nitrogen, together with an increase in the specific gravity as a result of diminished excretion of water. The urine also shows retention of chlorides and deposition of urates on cooling. When the attack is over these changes are reversed: there is a marked diuresis and the urine becomes alkaline. They also found positive van den Bergh reactions of the biphasic type in a large proportion of their cases. They believe that these changes are indicative of a disturbance of hepatic function, and that the increased ammonia excretion and the retention of chlorides are protective mechanisms. On these findings they based the following treatment:

1. A lacto-vegetarian diet. This is advised because in many cases the antigens responsible for allergic conditions are absorbed from the alimentary tract, being specific food proteins, intermediate products of protein digestion, or toxic substances formed by bacterial action on protein.

2. Ammonia in the form of sal volatile is given because the increase in ammonia excretion is regarded as a protective mechanism.

3. Glucose is prescribed because glycogen is essential for the antitoxic and proteopexic functions of the liver.

In my cases I have not altered the patient's normal diet unless the diet chart has indicated specific food sensitiveness, when that article of diet has been eliminated. Most of my patients have been given glucose, while some have had ammonia. I have given the glucose in the form of Glucose D; this also contains calcium, which is probably of value, as some cases of urticaria show diminished blood calcium.

Desensitization

Finally, in most of the cases an attempt has been made to desensitize the patient. This has been tried in various ways. It will be remembered that Vallery-Radot and Pagniez (1916) showed that attacks of urticaria due to food can sometimes be prevented by giving the patient, about an hour before his meal, a small amount of the food to which he is sensitive. More recently Urbach and Fasel (1931) have adopted and extended the method by giving specific peptones, which they call "propeptans." In many of my cases I have used a preparation called anaclasia; this contains various peptones derived from common articles of diet, together with magnesium and sodium hyposulphites, which have also been used as desensitizing agents. Another method of desensitization that I have used extensively is intradermal injections of proteose prepared from the urine passed during or immediately after an attack of urticaria. I usually give 0.1 c.cm. of a 1 in 1,000,000 solution of proteose every five to seven days.

A further method of desensitization is autohaemotherapy. Five to ten c.cm. of blood are withdrawn from a vein and immediately reinjected intramuscularly. This method has been employed for the past twenty-five years, although its mode of action is still uncertain. In 1931-2 I carried out some experiments in order to try to throw some light on its mechanism (Burgess, 1932, 1933). My results showed that its effects do not depend on the production of leucocytosis, but that cases which were

improved by the treatment had given positive reactions to their own proteoses before and negative reactions after treatment. It is therefore possible that the blood contains antigens. In other cases peptone injections were given.

Results of Treatment

The results obtained were as follows. Of the thirty-three cases treated twenty-two recovered completely, five showed considerable improvement—that is, whereas previously they had suffered from severe daily attacks they now have occasional slight attacks—two did not improve, while I have been unable to follow up four cases. Therefore, of the twenty-nine cases of which I have full records twenty-seven either completely recovered or improved considerably. In the accompanying table I have analysed

Table showing Results of Treatment

	No. of Cases	Recovered or Improved	No Improvement	No Record
Group a:				
Focal infection ..	9	9	—	—
Psychotherapy ..	3	3	—	—
HCl	8	8	—	—
Totals ..	20	20	—	—
Group b:				
Diet	6	6	—	—
Glucose	27	21	2	4
Ammonia	8	5	1	2
Totals ..	41	32	3	6
Group c:				
Anaclasia	18	14	2	2
Proteose	11	8	2	1
Autohaemotherapy	13	10	—	3
Peptone	5	5	—	—
Totals ..	47	37	4	6

the results obtained from the various methods of treatment. As one would expect, the treatment which I have included under Group *a*—that is, treatment based on aetiological factors discovered by investigation—shows the best results, in all cases being followed by complete recovery or considerable improvement. The results obtained from the treatment based on theoretical considerations (Group *b*) appear to support the theories concerned, but are not quite so good as in the first group. Desensitization (Group *c*) also proved satisfactory, although one cannot claim that the methods which theoretically should be specific give superior results to the non-specific methods.

Conclusion

It would thus seem that there is evidence to support the theories put forward to explain the aetiology of chronic urticaria. It is clear that urticaria is a symptom rather than a disease, an end-result of many different pathological processes, and every case must be judged on its merits. The routine by which I have endeavoured to systematize, so far as possible, the investigation and treatment of these cases may be summarized as: (1) Removal of foci of infection when present. (2) Elimination of foods to which the patient is sensitive. (3) Investigation of psychological difficulties. (4) Treatment of dyspepsia when present. (5) The administration of glucose, calcium, and ammonia. (6) Some method of desensitization. If the objection be made that this method of treatment is of the "shot-gun" variety, it must be remem-

bered that in a condition of such complexity good results are only likely to be obtained by a judicious selection of various therapeutic measures in the same case.

REFERENCES

- Barber, H. W., and Oriel, G. H. (1928). *Lancet*, 2, 1009, 1064.
 Burgess, N. (1932). *Brit. J. Derm. Syph.*, 44, 124.
 — (1933). *Ibid.*, 45, 333.
 Coca, A. F. (1926). *Arch. Path. Lab. Med.*, 1, 96.
 Langdon-Brown, W. (1932). *Practitioner*, 129, 641.
 Oriel, G. H. (1931). *Proc. roy. Soc. Med.*, 24, 1171.
 Stokes, J. H., and Pillsbury, D. M. (1930). *Arch. Derm. Syph.*, Chicago, 22, 962.
 Urbach, E., and Fasel, P. (1931). *Arch. Derm. Syph.*, Vienna, 164, 133.
 Vallery-Radot, P., and Pagniez, P. (1916). *Presse méd.*, 24, 529.
 Walzer, A. and M. (1927). *Amer. J. med. Sci.*, 173, 279.
 — (1928). *Arch. Derm. Syph.*, Chicago, 17, 659

INDUCTION OF LABOUR

THE COMBINED METHOD OF PREMEDICATION
WITH CASTOR OIL AND QUININE, AND
ARTIFICIAL RUPTURE OF
MEMBRANES

BY

EILEEN C. WISE, M.R.C.S.

Visiting Obstetrician, British Hospital for Mothers and Babies,
Woolwich

Artificial rupture of the membranes as a means of inducing labour has been employed for so many years that its value as an obstetric manœuvre might well be expected by now to be precisely established. Goodwin (1937) reports its use 150 years ago, and states that Denman in 1793 was one of its earliest advocates. Yet it has not gained general approval, largely on account of the traditional association of "dry labour" with delay and difficulty. It seems possible that confusion has arisen from a failure to distinguish sufficiently clearly between the course of labour when the only abnormal feature is the early loss of liquor amnii, and the course when its escape is associated with other factors which really play the major and responsible part in producing difficulty. If the early escape of fluid could be fairly exonerated from the drawbacks for which it is commonly blamed, and if, further, its artificial production could be established as a reliable means of inducing labour, a valuable point would be reached for practical guidance. There are many situations in obstetrics where reliable control over the start and completion of labour can be calculated to lessen risk to mother and child or both. The difference in effort required to expel a foetus of, say, 6½ lb. and one of 8 lb. may be of vital service to a woman with morbus cordis. The same difference may save the foetus where contraction of the maternal pelvis exists. Many other instances occur to the mind, of which mention may perhaps be made of the great advantage of being able to prevent gestation from continuing beyond the normal period of forty weeks.

Many opinions could be quoted both in favour of and against this method of inducing labour. A good sample in British and American literature is contained in the writings of Drew Smythe (1931), Drew Smythe and Thompson (1937), FitzGibbon (1931), Tennent (1938), Guttmacher and Douglas (1931), Jackson (1934), Wilson (1934), and Goodwin (1937). Drew Smythe (1931) recommends induction of premature labour by artificial rupture of the membranes for all cases in which the foetus presents

by the vertex. He includes in his list of indications pyelitis of pregnancy, cardiac disease, and disproportion. This last condition he considers to be one above all others in which induction of labour is indicated. Eden and Holland (1937) are in favour of the method, describing it as both safe and reliable.

Indications for Induction in the Present Series

This method has been employed 681 times during the last six years at the British Hospital for Mothers and Babies, and with the experience gained, and as certain points in technique have been improved, a high appreciation of its value has grown. The cases chosen for analysis number 287, and they comprise all those in which the method has been used over a period of two years. It may be regarded as the routine policy of election at this hospital in the obstetric conditions presently to be described. The indications for induction in this series of cases were: disproportion, 100 cases; at term to prevent post-maturity, seventy-one cases; toxæmia of pregnancy, forty cases; chronic bronchitis and various states of ill-health classified as maternal debility, twenty-six cases; post-maturity, twenty-five cases; morbus cordis, seventeen cases; hydramnios and twins, five cases; ante-partum hæmorrhage, three cases. Total cases 287.

Method of Induction

Once the decision to induce labour was reached the method employed was carried out in the following stages. (1) A course of early continuous premedication with sulphate of quinine. (2) Preliminary admission to hospital. (3) The use of castor oil and an increase in quinine medication. (4) Artificial rupture of the membranes followed by an immediate and deliberate expulsion of liquor amnii.

Early Quinine Medication.—All primigravidae and those multigravidae with a history of defective uterine action in previous labours received 1½ grains of quinine sulphate three times daily during the last three weeks for which the pregnancy was allowed to continue.

Preliminary Admission to Hospital.—The routine for primigravidae selected for induction was admission to hospital five days before the anticipated date of artificial rupture of the membranes. For multigravidae this period was reduced to three days.

The Use of Castor Oil and Increased Quinine Medication.—At 6 a.m., castor oil 1 oz.; at 10 a.m., 2 p.m., and 6 p.m., quinine sulphate powders 10 grains; and at 5 p.m., a simple enema and a hot bath were administered to primigravidae on the day after admission (second day) and repeated on the fourth day; to multigravidae they were given on the second day.

Artificial Rupture of the Membranes.—This was performed on the fifth day in primigravidae, and on the third day in multigravidae. The operation was carried out in the labour theatre with all the usual antiseptic and aseptic precautions. The patient being in the lithotomy position, the operator passed the first two fingers of the right hand into the vagina, and then a finger through the os until the foetal head was reached. In some cases in which the cervix was very posteriorly placed it was necessary to hook it forward with the examining finger. A long stylet or vulsellum forceps was passed along the hand into the vagina and guided through the os until it was in contact with the foetal head. The membranes were ruptured and the liquor allowed to drain off. This was