others the blood-laevulose curve is not only raised but is also prolonged.

One case of carcinoma is of special interest as giving some indication of the extent of liver damage which must occur before the laevulose tolerance begins to show deficiency. The blood-laevulose curve, estimated five days before death, showed that the rise above the apparent initial level was, at half an hour after the dose of laevulose, 14 mg.; at one hour, 16 mg.; and at two hours, 10 mg. per 100 c.cm. This was a border-line result—only just outside the normal range. At necropsy the liver was found to be extensively disorganized by carcinoma, infarction, and necrosis, and it was estimated that about half the liver tissue had been destroyed. Now in such a case the condition resembles the result of partial hepatectomy. Little, if any, blood will pass from the portal veins to the hepatic veins through grossly damaged tissue. Almost all the blood which passes through the liver will traverse tissue of good functional capacity. It seems that in such circumstances a deficient laevulose tolerance indicates that at least half the liver tissue is out of action. In diffuse lesions, such as acute hepatitis, the condition is quite different, since all the circulatory paths through the liver will traverse damaged tissue, and this probably accounts for the fact that in acute hepatitis the laevulose tolerance is usually definitely deficient during the attack.

The diagnostic value of the test is limited by the existence of a large functional reserve in the liver. A normal result does not exclude liver disease. But, like renal function tests, the laevulose-tolerance test may give information as to the extent of the damage when a lesion has been recognized on clinical evidence. In some cases the test has diagnostic value; I have met with instances in which deficient laevulose tolerance has been demonstrated when the existence of liver disease could be neither proved nor excluded on the clinical evidence. The test may sometimes assist diagnosis—for example, in cases of unexplained haematemesis, unexplained anaemia, and in detecting hepatic cirrhosis in Banti's disease at a stage when the clinical evidence is inconclusive.

Summary

As a result of earlier work it has been shown that the laevulose-tolerance test becomes more reliable if the results are based on estimations of blood laevulose rather than total blood sugar.

My modification of the diphenylamine method for the estimation of blood laevulose has resulted in greater speed and accuracy of analysis.

Normal results for blood-laevulose curves obtained by this method are presented.

Some additional results in cases of liver disease are described.

I am indebted to the honorary medical and surgical staff of the Royal Victoria Infirmary for the opportunity to examine cases and for the use of clinical notes; to my colleagues on the pathological staff for histological and post-mortem data; and to the volunteers who offered themselves as subjects for the study of normal results.

REFERENCES

- Herbert, F. K. (1938). *Biochem. J.*, **32**, 815.
— and Davison, G. (1938). *Quart. J. Med.*, **31**, 355.
Jordan, R. C., and Pryde, J. (1938). *Biochem. J.*, **32**, 279.
Scott, L. D. (1935). *Brit. J. exp. Path.*, **16**, 489.
Steinitz, H. (1937). *Acta med. scand.*, **93**, 98.
Stewart, C. P., Scarborough, H., and Davidson, J. N. (1937). *Edinb. med. J.*, **44**, 105.
— (1938). *Quart. J. Med.*, **31**, 229.

ASTHMA AS A SEQUEL OF NASAL INJURY

BY

W. A. BOURNE, M.D., M.R.C.P.

Assistant Physician, Royal Sussex County Hospital

The association of asthma with disease of or pressure on areas of the nasal mucosa has been recognized for many years, and has often been made the basis of anti-asthmatic treatment. The importance of nasal pathology is, however, now much debated, and an authority has stated that "physicians nowadays are beginning to realize that it is only a minor cause in some cases and a major secondary effect in a majority of adults" (Bray, 1931). The importance of trauma in this respect has not been stressed, and as it may at times appear to play a part—not merely minor or secondary, but actually a sole part—in the production of asthma, the following two cases may be of interest.

Case I

The patient, a messenger girl aged 17, had been exceptionally healthy, without chest trouble of any kind, urticaria, migraine, or other allergic manifestations. The family history was similarly free. On June 27, 1932, she was flung from her bicycle, and her head struck the ground violently. She was admitted to Exmouth Hospital, where on examination she was found unconscious, with stertorous respiration, pupils dilated and non-reactant, and limbs flaccid, with occasional spasms. There was a tri-radiate wound, the largest limb of which extended upwards and to the left from the nasion for a distance of three inches, exposing the frontal bone. There was extensive bruising of the face, with haematomata of both orbits. There were also abrasions of hands, knees, and shins. The head wound was cleaned and dressed without suture. For five days the patient's condition appeared grave. Symptoms of cerebral concussion and irritation then disappeared rapidly, the wound closed, and in three weeks she was discharged at her own request. As the facial swelling subsided it was seen that there had been a fracture of the bones of the nose, with lateral displacement of moderate degree. This was not treated. She continued to attend as an out-patient, and appeared to have made a good recovery. There were no symptoms of chest trouble.

On August 7, 1932, she suddenly developed spasmodic dyspnoea of expiratory type, asthmatic in character, very severe, and relieved by adrenaline. Attacks of this kind continued in spite of treatment on usual medical lines, with local nebulae. Protein-sensitivity tests gave no result. In November she was referred to a hospital nose and throat department, where it was reported that there was deflection of the septum and scarring, definitely due to the accident, but that in view of the patient's chest condition operation was not advised. The asthmatic condition continued for a further period of sixteen months, with some improvement in the summer of 1933, but with prolonged attacks of great severity. The patient became kyphotic, dyspnoeic, and cyanosed, instead of stalwart and upstanding, with the clear colour that is often seen in the red-haired. She was quite unable to work, in spite of courses of vaccine and peptone treatment and breathing exercises.

In March, 1934, operation for resection of the septum was performed. The immediate effect of this was to some extent disappointing. Nasal obstruction was relieved, but the asthma continued, slowly diminishing in severity until May, 1935, since when the patient has had no attacks. Treatment of any kind ceased in February, 1935.

Case II

A retired lieutenant-commander, aged 40 and unmarried, had a history free from any chest trouble or allergic manifestation. The family history was similarly free. On August

18, 1935, when in perfect health, he was involved in a motor accident, sustaining cerebral concussion, fractured right fibula, lacerated wounds of the forehead and left orbital regions, and bruising of the facial tissues, hands, and left knee. He was admitted to Hove Hospital, where he was unconscious for twenty-four hours. Three attempts to control the fragments of the fibula were made, and finally a bone peg was inserted and he was discharged with the leg in plaster.

On October 4, 1935, he developed acute nocturnal dyspnoea, expiratory in type, of great severity. On examination the lungs were full of expiratory sounds. There was a forked scar over the side of the bridge of the nose, the two limbs running up the forehead and over the right eyebrow. The nose internally showed obstruction of the left naris. Adrenaline relieved the attack, but though further treatment with a nasal spray of ephedrine and chlorotone, and the usual drugs internally, kept the more violent paroxysms under control, the patient was in constant discomfort. With the former case in mind, he was referred to Mr. Harold Downer, who reported: "The nasal septum is markedly deflected to the left, and there is opacity of the left antrum."

Submucous resection was performed on November 25, 1935, by Mr. Downer. By November 27 the asthma was much improved, and since January, 1936, has not recurred.

Comments

It is submitted that nasal injury resulting in deformity was the precipitating factor in the development of asthma in these patients, who had never previously shown any asthmatic tendency, and have shown none in the considerable time which has elapsed since full recovery from operation. On the other hand, the great majority of cases of nasal injury have no sequel of this kind. A number of reasons may be suggested for the appearance of asthma in these two individuals. Doubtless unknown predisposing factors were present, as emphasized by Maxwell (1936). In addition, however, the above accidents produced severe trauma. The violence with which the girl struck the ground, and the man the wind-screen, was probably more than is involved in most nasal accidents. There was also an element of psychical trauma. Shock at the time, the possibility of police intervention, litigation in the civil courts, and anxiety regarding means of livelihood were added. Further—and it is suggested that this is most important—it happened that other and more urgent injuries than those to the nose were present, and owing to the pressure of the treatment of these the condition of the nose and the nasal passages could not be determined in the early stages and was not investigated before discharge from hospital. For these reasons deflection of the septum and scarring of the wound in the mucosa probably combined to produce a trigger area which, in patients whose nervous balance was disturbed, resulted in the sudden onset of acute asthmatic attacks. These appeared after intervals of six to eight weeks.

In fractures of the nose unaccompanied by other injury treatment is much more early given, and even if external reposition is not perfect, the nares are likely to be left patent and conditions for asthma production are prevented. When the injuries in these two cases had produced asthma, the presence of asthma was no contraindication to operative treatment. Delay in operation was followed by a poor immediate result and a lengthy interval before the disappearance of the attacks. Immediate operation was followed by rapid relief and disappearance.

The operation was that of simple resection, and it should be emphasized that no suggestion is made that nasal treatment for asthma of other origin should take this form. Nor is it regarded as clear why such operation

should relieve mechanical pressure, since the asthmagenic area of Dixon and Brodie does not appear to coincide with that concerned in submucous resection. However, though it is realized that two cases are quite insufficient for far-reaching conclusions, the following of some practical importance seem justifiable.

Conclusions

Nasal injury may be a precipitating factor in the production of asthma.

In cases of nasal injury accompanied by more severe injury elsewhere, the condition of the nose should be borne in mind, ascertained before discharge, preferably by specialist opinion, and remedied if in any way doubtful.

Once asthma is established, operation to remedy nasal deformity is indicated at the earliest possible moment.

The presence of asthma is no contraindication to operative treatment, and the results may be expected to be good in proportion to the promptitude of operation, provided that the history and the condition of the patient show no other cause for the development of the asthma.

I am much indebted to Mr. Harold Downer for his criticism and advice.

REFERENCES

- Bray, G. W. (1931). *Recent Advances in Allergy*, p. 110, London.
Maxwell, J. (1936). *British Medical Journal*, 1, 874.

MANAGEMENT OF A HAY-FEVER IONIZATION CLINIC

A COMPARATIVE STATISTICAL SURVEY OF RESULTS IN 1937 AND 1938

BY

PHILIP FRANKLIN, F.R.C.S.

Honorary Laryngologist, Margaret Street Hospital for
Diseases of the Chest

The Margaret Street Hay Fever Ionization Clinic, attached to the Margaret Street Hospital for Diseases of the Chest, was opened in the spring of 1937, and the results of the treatment for 1937 were reported in the *Journal of April 30, 1938* (p. 948). The figures for 1938 show a great improvement over those of 1937. This improvement may either be due to the general mildness of hay fever during the season of 1938 or be the effect of our modified technique. In the first place, in 1938 we to a large extent gave up prophylactic ionization, and most of the patients were treated either immediately before the expected onset of the hay fever or actually at the beginning of the attack. Secondly, last year we were using an ionization current of 6 to 8 milliamperes, whereas in 1937 the current was only 3 to 5 milliamperes. Subsequent experience will show whether the improved results of 1938 are really due to the modified technique.

Organization

The organization of the clinic has been further simplified, and can be briefly described as follows. The appointment book is arranged in such a way as to show at a glance the ionization sets (of which there are eighteen) still available at any particular time. The patient is received by an attendant, who takes his particulars and at the same time sprays the nose with a 2 per cent. solution of larocain. The patient is then ready for the nose-packing, which is carried out by medical assistants. The packing material consists of thin squares of cotton-wool reinforced by a single layer of gauze and soaked in a 1 per cent. solution of zinc sulphate. The cotton-wool