work-place in increasing or depressing the rate of morbidity. It by no means follows that morbidity and mortality rates rise and fall together. It may and does happen that an establishment with a low mortality rate has a very high sickness-absence rate. But a high sickness rate often means psychologically unsatisfactory conditions. Morbidity, if it could be fairly measured, might well be a more valuable industrial pointer than mortality. Here is a task for the future.

At least I hope to have convinced you that even in mortality data alone there is material which not only the professional statistician but the social worker, the administrator, and indeed every educated citizen, should study and understand. All this light on contemporary social conditions should not be imprisoned between the covers of a Blue-book, but should illumine our understandings and our hearts, teaching us to work for a better England, like Chadwick long ago.

THE NATURE OF CHRONIC MASTITIS

BY

H. J. B. ATKINS, D.M., M.Ch., F.R.C.S.

Assistant Surgeon to Guy's Hospital; Co-Director of the Mastitis Clinic, Guy's Hospital

The resistance of the body to infectious disease involves the operation of three distinct mechanisms. The first of these depends on an inherited quality of cell structure or metabolism, and is called a natural immunity. The second is a serological reaction leading to the production of antibodies, and is termed acquired immunity. The third is a local tissue reaction whereby leucocytes and fibroblasts, summoned to the site of invasion, erect a barrier of granulation tissue against the further inroads of the disease. This local reaction accounts for a series of clinical conditions known collectively as the chronic granulomata.

If we attempt to relate neoplastic to infectious disease, we find certain broad features of resemblance. Infections are caused by the introduction into the body of pathogenic organisms, neoplasms by the operation of some extrinsic factor. Individuals, families, and even races exhibit a natural immunity to certain infectious diseases, and there appears similarly to be a natural immunity to neoplastic disease in certain strains or species of animals, depending equally on inherent qualities of cell structure or metabolism. A type of acquired immunity in the serological sense is met with in relation to implanted tumours, but has not been demonstrated with spontaneous tumours or in relation to the extrinsic factor itself. It is, however, probable that some immunizing mechanism, closely allied to that obtaining in bacterial invasion, is provoked by neoplastic disease. It is thought that such a mechanism may be responsible for the rarity of the occurrence of two primary growths in one individual. It remains to inquire whether there is any process involved in neoplastic disease comparable to the formation of granulation tissue in infectious disorders, and, even more remotely, whether there is a series of clinical conditions analogous to the chronic granulomata.

Is Chronic Mastitis a Precancerous Condition?

Evidence relating to this problem is obtained from a study of the natural history of the disorder known as chronic mastitis. There has long been an apparently irreconcilable difference of opinion between the histologist and the clinician as to the precancerous nature of this disease, a difference of opinion based on evidence so

conflicting as to demand the closest analysis. For many years the histologist has maintained that the epithelial proliferation of the mammary ducts to which, with cyst formation, he has given the name of "chronic mastitis" leads to cancer (Cheate and Cutler, 1931). He says this because he can take a series of microscopical sections from different breasts and, arranging them in order, display an unbroken sequence of epithelial change from the normal at one end of the scale, through the epithelial patterns characterized by him as chronic mastitis, to carcinoma at the other. This evidence is suggestive but not conclusive. By reversing the sequence it could equally well be maintained that the normal breast develops from carcinoma, a postulate which we know from outside evidence to be absurd. The assumption, however, that the order chosen arbitrarily by the histologist represents a natural sequence receives strong confirmation from a collateral source.

The experimental pathologist has succeeded in producing cancer of the breast in animals by the injection of oestrin; and Lacassagne (1932, 1933, 1934, 1936) has shown that the preliminary changes in the breast epithelium which eventually lead to cancer are precisely similar to those designated by the histologist as chronic mastitis. On this evidence it must be concluded that the epithelial changes observed under the microscope in sections taken from the breasts of cases of chronic mastitis may be precancerous. On the other hand, an overwhelming number of clinicians of experience believe that the disease which they term chronic mastitis has no especial liability to become cancer. This belief is founded on impressions derived from long association with many cases of chronic mastitis. Impressions are apt to be fallacious, and the only scientific evidence on this aspect of the problem comes from Bloodgood (1932), who followed up a series of cases for many years and concluded that the incidence of breast cancer in patients suffering from chronic mastitis was no greater than in the general population. In the mastitis clinic at Guy's Hospital, where 170 cases of mastitis, many of them of long standing, are being investigated and followed up, no case has developed carcinoma within the two and a half years' life of the clinic.

In a previous paper (Atkins, 1938) I have pointed out that the only way to ascertain whether a disease A predisposes to a disease B is to take a sufficient number of cases of A, follow them up for some years, and observe whether disease B occurs in a greater proportion of these than in the general population. This investigation has been undertaken in relation to chronic mastitis and cancer, and has not revealed a predisposition of the one for the other. Although statistical evidence to be conclusive must be based on a larger number of cases than have yet been investigated, it is certain that carcinoma of the breast follows chronic mastitis in a strikingly small proportion of cases, having regard to the precancerous nature of the epithelial changes which accompany the disease.

Significance of the Fibrous-tissue Proliferation

These precancerous tendencies must therefore be arrested, and it is in this connexion, I think, that the histologist has concentrated too exclusively on the epithelial changes of chronic mastitis, and not given sufficient weight to the significance of the fibrous-tissue proliferation. In the biopsy material from the mastitis clinic a fibroblastic reaction has been an outstanding feature of the microscopical picture. In comparing a series of slides from this clinic with a series taken from the breasts of women who had died from diseases unconnected with the breast, I found that it was as easy to distinguish the mastitis clinic

sections from the post-mortem room sections by observing the fibrous tissue as by concentrating on the epithelial pattern. The epithelial changes designated chronic mastitis by the histologist and assumed to be precancerous were present in a small proportion of the control sections, but it was quite exceptional to find an excess of highly nucleated fibrous tissue in sections taken from patients not complaining of the symptoms of chronic mastitis. The clinician when he refers to chronic mastitis means a painful nodular breast, and an outstanding feature of the painful nodular breast is fibrosis with or without cysts. If now this fibrosis accompanies epithelial changes, which threaten cancer but rarely fulfil their threat, it is possible that the fibrous tissue may be produced in response to that threat and may play a part in preventing it from coming to fruition. Consequently it would be just those patients who are able to develop a fibrous-tissue reaction to epithelial overgrowth—that is, those who complain of the symptoms of chronic mastitis—who would be unlikely to develop cancer. The conception that the fibrosis of chronic mastitis is an expression of a local defence mechanism resolves the problem of the precancerous nature of this disease in conformity with both the histological and the clinical evidence. Moreover, it falls into line with the established procedure of Nature in dealing with infectious disease and provides the counterpart to the granulation tissue which is such a prominent feature of that process.

That the problem of the protective function of fibrous tissue is an exceedingly complicated one is appreciated from a consideration of the relative malignancy of the different types of carcinoma of the breast. The malignancy of a breast cancer cannot be assessed from the amount of fibrous tissue present, as would be the case if fibrosis were the only mechanism of defence against the malignant cell. Where so many factors operate, these may exert a predominating influence on the clinical picture. In the absence, for instance, of a certain degree of natural or acquired immunity no amount of fibrosis could of itself control the spread of the disease. One factor among many might only be expected to colour the whole picture when it was operating in its most intense form. Such an intensity of fibrosis is met with in the atrophic scirrhus: here the combination of fibrosis with the most anaplastic type of epithelial cell leads to a control of epithelial growth which may delay the spread of the disease for many years. Thus in those cancers where the morphology of the epithelial cell would lead us to expect a rapid growth but which lie dormant for years, and in chronic mastitis where precancerous changes suggest a malignant tendency which is not fulfilled, the only accompanying phenomenon observable is fibrosis. On that account the theory of the protective function of fibrous tissue against epithelial new growth is put forward as a working hypothesis. This appeal to the old tissue-tension theory of Waldeyer and Thiersch does not suffer from the objections to that theory in its original form. It is not suggested that an upset of the equilibrium between fibrous and epithelial tissue is the whole story of the development of cancer, but it is proposed that fibrosis is engendered by a tendency to epithelial overgrowth and that the fibrosis is one part of the complicated mechanism of defence against neoplastic disease comparable to granulation tissue in infectious disease.

Pathology of the "Fibromatoses"

Lastly, if this be so, it is pertinent to inquire whether there is a series of clinical conditions comparable to the chronic granulomata and expressive of a local tissue

reaction to the threat of malignant invasion elsewhere in the body. We have discussed the relation of chronic mastitis to this problem, but there is a class of clinical phenomena, which I will call the fibromatoses, which might repay investigation in this connexion. I refer to leucoplakia, Riedel's thyroiditis, fibrotic prostate, linitis plastica, and fibromatosis of the colon. An entirely satisfactory pathology has not as yet been assigned to these conditions, and in order to stimulate further investigation the suggestion is put forward that the fibromatoses, including chronic mastitis, may be the homologues in neoplastic disease of the infective granulomata.

Summary

The natural history of chronic mastitis is considered, and an analysis made of the evidence as to whether the condition is precancerous.

It is submitted that the fibrosis occurring in chronic mastitis is part of the mechanism of defence against neoplastic disease, comparable to granulation tissue in infectious disease.

The suggestion is made that the "fibromatoses" might repay investigation in connexion with the appearance of cancer in other parts of the body.

REFERENCES

- Atkins, H. J. B. (1938). *Lancet*, **1**, 707.
 Bloodgood, J. C. (1932). *Amer. J. Cancer*, **16**, 103.
 Cheate, Sir Lenthal, and Cutler, Max (1931). *Tumours of the Breast*, London.
 Lacassagne, A. (1932). *C. r. Acad. Sci., Paris*, **195**, 630.
 — (1933). *C. r. Soc. Biol., Paris*, **114**, 427.
 — (1934). *Ibid.*, **115**, 937.
 — (1936). Five Lectures, International Cancer Research Foundation, Philadelphia.

BLOOD LAEVULOSE IN THE LAEVULOSE TOLERANCE TEST

WITH SPECIAL REFERENCE TO NORMAL STANDARDS

BY

FREDA K. HERBERT, M.A., M.B., B.S.

(From the Pathological Department of the Medical School, King's College, and the Royal Victoria Infirmary, Newcastle-upon-Tyne)

When laevulose reaches the normal liver by the portal veins it is rapidly utilized, part being oxidized and part stored as glycogen. The rate of utilization is almost as rapid as the rate of absorption from the intestine, so that very little laevulose reaches the general circulation after oral administration of the sugar. If the glycogenic function of the liver is impaired by disease the rate of utilization fails to keep pace with the rate of absorption, and an abnormally large amount of laevulose passes through the liver into the systemic blood stream. These facts provide the basis for the laevulose-tolerance test of hepatic function. In the past the method of performing the test has been to follow the changes in total blood sugar after a dose of laevulose, and it has been assumed that any increase above the fasting level is due to laevulose.

Recent Investigations

In recent years methods have been developed for the analysis of blood laevulose, and various authors have made fresh investigations of the laevulose-tolerance test in which the blood laevulose has been directly estimated