WAR WOUNDS AND AIR RAID CASUALTIES

The following is one of a series of lectures on the Treatment of War Wounds and Air Raid Casualties recently delivered at the British Postgraduate Medical School

BLOOD TRANSFUSION

BY

JANET M. VAUGHAN, D.M., F.R.C.P.

I propose to discuss the problem of blood transfusion particularly from the point of view of a war-time emergency service, though much of what I have to say is also applicable to a transfusion service in times of peace.

Sources of Blood for Storage

Blood for purposes of transfusion may be freshly removed from a living donor and immediately given to the patient, or blood that has been stored may be used. Recognized sources of blood for storage are the placenta, the cadaver, and adult donors. It is obvious that under war conditions fresh blood cannot be used as a routine. The disorganization of means of communication caused by aerial bombardment makes it impossible to call upon donors at those times when blood is most needed. Medical and surgical personnel will be required to attend to patients during severe air raids or at periods of active fighting. They must administer blood and not spend time withdrawing blood. Donors obviously cannot be sent up to the front-line dressing stations. It will be absolutely essential, therefore, to have adequate supplies of stored blood.

Even in ideal circumstances the yield of blood from the placenta is small, on an average 80 c.cm. Though certain workers advocate its use at least in times of peace, opinion on the whole suggests that the labour involved in its collection is not repaid by the small amounts of blood which result (Goodall *et al.*, 1938; Howkins and Brewer, 1939; Page *et al.*, 1939; Halbrecht, 1939). Cadaver blood has not been used in this country. In Russia it is claimed to be satisfactory (Yudin, 1937). On theoretical grounds, in time of war it should prove an excellent source, since in persons dying a sudden death spontaneous fibrinolysis occurs, and it is not therefore necessary to use an anticoagulant. In practice, however, the collection of blood from war cadavers has, according to Saxton (1939) and Duran (1939), proved unsatisfactory.

Stored blood from adult donors has, however, been an almost universal success. Recent experience in this country confirms the work of Russian (Filatov, 1937; Bagdassarov, 1937) and Spanish workers (Duran, 1939) on this point (Elliott, Macfarlane, and Vaughan, 1939; Boland, Craig, and Jacobs, 1939). Biddle (1939) has for the last year used nothing but stored blood at the Ipswich Hospital. He claims that in giving 150 transfusions he has had fewer reactions than during the period when he used fresh blood. He had slight reactions in about 5 per cent. of his cases. Experience in the department of pathology of the British Postgraduate Medical School has been similar. In receiving fifty consecutive transfusions with stored blood four patients had a slight rise of temperature and four had mild rigors. Saxton has reported a higher reaction rate in Spain (1939), but it appears probable that complicating factors, to be discussed later, may have been present.

Such figures as are available suggest that the rise in red cells and haemoglobin following the use of stored blood is as satisfactory as that achieved by fresh blood, while improvement in the clinical condition of the patient

is equally good provided that certain criteria of satisfactory preservation, to be discussed later, are adhered to (Elliott, Macfarlane, and Vaughan, 1939).

Blood Transfusion Institutes

Experience both in Russia and in Spain has shown that the best results are obtained if blood is withdrawn, stored, and distributed from special centres equipped for the purpose. It is obvious that the existence of such centres does not prohibit small and isolated units from arranging for their own supplies in times of emergency. It is, however, easier to ensure adequate sterility of apparatus and methods of collection with trained personnel working in suitably equipped centres.

Donors

It proved possible in Spain, in spite of war-time conditions, to organize a service of donors who came up to give blood at regular intervals. Blood was withdrawn apparently without ill effects as often as once every three weeks in many cases. On theoretical grounds it would be wise to have a longer interval—if possible at least three months, as is advised at present by the Red Cross Voluntary organization. Such a period will, however, clearly prove impossible in time of war: 500 to 600 c.cm. of blood is a reasonable amount to take from each donor. It has been suggested that it is wiser to take blood from a donor who has had no food for several hours. We kept a careful note of the time and character of the last meal in all the donors of blood for storage, and could not detect any relation between reactions and the recently fed donor.

In Barcelona, Duran (1939) stored only blood from Group IV Moss (O international), since under war conditions adequate grouping at the front proved impossible. In our limited experience here we found that it was wise to keep blood of all groups. This policy resulted in no wastage. If donors are to be bled as frequently as once a month they should be given large doses of iron, and if possible extra rations, at the time of withdrawal.

Apparatus

Various types of apparatus have been used for the withdrawal, storage, and administration of blood. No one method is at the moment completely satisfactory. On theoretical grounds the ideal technique for war-time should fulfil the following conditions:

1. The whole apparatus must be autoclaved in one piece ready for use without further manipulation if sterility is to be secured.
2. Blood must be collected without exposure to air for the same reason.
3. All glass-ware must be strong.
4. The container should be completely filled to reduce movement of blood to a minimum. Even slight shaking of blood after storage for longer than twenty-four hours is found to increase haemolysis, so that any technique that reduces agitation to a minimum is to be advocated.
5. No glass tubing should be left in contact with the blood after collection, as it is found to favour clot formation.
6. The technique of both collection and administration should be as simple as possible.

The method described by Duran (1939) fulfils the first four of the above conditions. The special storage

vessels required, however, even if somewhat simplified, are not easy to obtain on a large scale. Further, the method of collection and closing of the storage vessels appears too complicated for emergency conditions where extremely large quantities of blood may be needed.

The technique described by Elliott and others (1939) fulfils the first five conditions. The vessel is, however, completely filled, and in order to give blood by a pressure method a two-way tap and syringe is required. This can well be manipulated by a single individual, but it might be thought that it makes the apparatus too complicated for emergency use where skilled personnel will not necessarily be available.

Some modification of Boland, Craig, and Jacobs's (1939) method, where blood is stored under pressure, so facilitating administration, will probably prove the most satisfactory technique available should emergency immediately arise. Both collection and administration are easy and no complicated apparatus is required. The vessel is not, however, completely filled with blood, and if shaken in transit haemolysis is likely to occur. This risk is lessened if blood is transported immediately after collection, since red-cell fragility is not greatly increased in the first twenty-four hours of storage. Immediately after blood is collected air can be pumped into the container to raise the pressure, using a valve-tap somewhat on the principle of a bicycle valve, or pressure can be raised by the use of a Higginson syringe just before blood is administered. In giving blood to patients suffering from severe shock or haemorrhage whose veins are likely to be collapsed, some form of pressure is a great advantage. In many cases it saves the necessity of cutting down on the vein and inserting a cannula.

Method of Collection

In collecting blood for storage it is essential to adopt a strictly aseptic technique. We found three instances of contamination with *Staphylococcus aureus* in our first twenty samples of blood withdrawn with reasonable precaution. After that more rigid rules were laid down regarding asepsis, and no further contamination occurred. The operator wore a mask and scrubbed up as he would in the theatre. The skin over the antecubital fossa of the donor was cleansed with soap and water and then with alcohol. The whole apparatus, including the needle and preservative required for collection, was autoclaved together, so that it was not handled in any way, and it was so arranged that the blood was never exposed to the air.

Blood that flows freely into the preserving fluid keeps in more satisfactory condition than blood that flows slowly. It is therefore important to place the donor in a comfortable position with his arm near the edge of the arm rest, so that no kinking of rubber tubing can occur. If care is taken over the comfort and position of the donor it was found that 500 c.cm. of blood could be easily collected in from four to seven minutes. Before cleaning the donor's skin we placed an ordinary blood-pressure cuff round the upper arm and maintained a steady pressure of 80 mm. Hg. Boland and others (1939) have described the collection of blood into a flask containing a partial vacuum. Other workers have advocated the use of a Higginson syringe to maintain a negative pressure in the collecting vessel. Before withdrawing the needle from the vein, and after disconnecting it from the collecting flask, a few drops of blood should be taken into a tube containing 1 per cent. sodium citrate in normal saline for purposes of grouping and into a second tube for a Wassermann test. For convenience these two small tubes may be slipped under a rubber band attached to the collecting vessel before it is autoclaved.

Preservatives

Four preservatives have been used: (i) sodium citrate in varying concentration; (ii) physiological citrate; (iii) glucose citrate; and (iv) a mixed salt solution known as I.H.T. (see

table). We at first added 10 c.cm. of 3.8 per cent. sodium citrate to every 90 c.cm. of blood, giving a concentration of 0.38 per cent. sodium citrate, approximately similar to that used by Saxton (1937). Bagdassarov has used a higher concentration of 0.54 per cent. Repeated reference is made in the literature to the use of physiological citrate solution, though I have been able to trace only one figure given for the exact

Table showing the Preservatives used in Storing Blood

Solution	Authority	Constituents	% Concentration	Volume of Solution	Volume of Blood	Final % Concentration
(I) Citrate	Saxton ..	Sodium citrate	3.8	10	100	0.34
	Vaughan ..	" "	3.8	10	90	0.38
	Duran ..	" "	4.0	10	100	0.36
	Bagdassarov	" "	6.0	10	100	0.54
(II) Physiological citrate	Page <i>et al.</i>	{ Sodium citrate	1.25	10	10	0.625
		{ Sodium chloride	0.85			0.425
(III) Glucose citrate	Domanig	{ Sodium citrate	1.0	10	10	0.5
		{ Glucose	2.0			1.0
	Duran	{ Sodium chlorate	0.2	10	100	0.1
		{ Sodium citrate	4.0			0.36
(IV) I.H.T.	Moscow	{ Glucose	1.0	10	10	0.09
		{ Sodium chloride	0.7			0.35
		{ Potassium chloride	0.5			0.25
		{ Magnesium sulphate	0.004			0.01

composition of the solution. Page, Seager, and Ward (1939) used 1 per cent. of sodium citrate in 80 c.cm. of normal saline to which was added 80 c.cm. of blood. Glucose citrate is also mentioned, and the following formula is given by Domanig (1937): sodium citrate, 1 gramme; glucose, 2 grammes; sodium chlorate, 0.2 gramme; water, 100 c.cm. This is mixed with blood in equal quantities. Duran is reported by Ellis (1937-8) to use a final strength of 0.36 per cent. sodium citrate and 0.09 per cent. glucose (see table). The majority of workers have employed the citrate. We found that, however carefully blood was mixed with the citrate solution, clots always tended to form, which, though easily removed by filtration, made the administration of the blood rather more complicated than necessary. After twenty transfusions had been given with citrated blood we therefore used the anticoagulant first devised by the Institute of Haematology and Blood Transfusion at Moscow, commonly known as I.H.T. (Bagdassarov, 1937). The formula is: sodium chloride, 7 grammes; sodium citrate, 5 grammes; potassium chloride, 0.2 gramme; magnesium sulphate, 0.04 gramme; distilled water, up to 1,000 c.cm. It is mixed volume for volume with blood. Why this particular proportion of salts was chosen I have been unable to discover. Results with its use proved satisfactory, and no further trouble with clot formation was experienced. All observers agree that considerable dilution of blood for storage is an advantage; haemolysis occurs less readily, and, since stored blood anyhow tends to be sticky, dilution facilitates administration. As much as 2,600 c.cm. of mixed blood and I.H.T. has been given to a patient with no ill effects.

Grouping

This is not the place for a discussion of the technique of blood-group determination. It is necessary, however, to emphasize that accurate determination of blood groups is essential to the success of any transfusion service. Severe reactions are almost invariably due to the administration of incompatible blood. Such a mistake usually results from the use of weak test sera. All serum should have a minimum titre of 1 in 100 at least if it is to be kept for any length of time, and preferably higher. It is not always easy to find individuals with such a high titre. One of the most important tasks of any large transfusion service should be to guarantee a supply of such serum.

No serum, however good, should be kept for more than six months. It should not be stored in small capillary tubes for longer than a few weeks. Serum is best kept in bulk in 5-cm. ampoules, and a small supply should be put up in capillary tubes for immediate use. It is probable that many of the severe reactions following the use of stored blood in Spain were due to wrong grouping dependent upon unsatisfactory supplies of test sera, rather than to any inherent fault in stored blood. If possible cross-matching between recipients' serum and stored corpuscles should be carried out, but in times of emergency this may be a counsel of perfection.

Storage: Temperature and Duration

We employed a temperature of 4° C. for storage of blood. Satisfactory results have been reported with storage at a temperature as low as 1° C. and as high as 7° C. (Bagdassarov, 1937; Boland *et al.*, 1939). Haemolysis occurs if blood is frozen.

All observers agree that blood begins to deteriorate after the fifteenth day of storage, though it can be given without ill effects after storage for thirty-four days. We purposely kept some samples of blood so as to test the results of its administration, and found that even when it was thirty-three days old the patient seemed none the worse for its administration. That after the fifteenth day less satisfactory results are obtained is probably dependent upon the fact that the red cells become increasingly fragile about this time and haemolysis increases rapidly. The tendency to haemolysis becomes evident in citrated blood rather earlier than it does in blood preserved in I.H.T. Elaborate studies of the effect of storage, both upon red-cell characters and the biochemical composition, have been made by Russian workers and are summarized by Filatov (1937). The sugar content decreases from the second day of storage, while the lactic acid content increases, together with the inorganic phosphorus. Haemoglobin becomes more stable. The albumin and globulin contents are constant for at least fourteen days. The reaction becomes increasingly acid. Complement is completely destroyed in fifteen days, and after seven days is reduced to 50 per cent. The red cells remain unaltered for fifteen days; after that time they begin to show great variation in size and shape, and there is a rapid increase in cell fragility with increasing haemolysis. The reticulum is lost after five days, presumably owing to ripening of the young red cells. There is a gradual decrease in agglutinability. The sedimentation rate is gradually slowed. The platelets are said to have completely disappeared by the fifteenth day, changes beginning on the first day of storage. There is considerable divergence of opinion about the effect of storage upon white cells. Filatov (1937) believes that degeneration begins between the third and fifth days. Irger (quoted by Filatov, 1937) goes so far as to maintain that 57 per cent. of the white cells are intact on the twenty-fifth day. In practice it is found that if the plasma immediately above the sedimented red cells is amber rather than red in colour no excessive haemolysis has occurred and the blood is suitable for use.

Sterility

The literature contains few precise statements concerning the sterility of blood withdrawn for storage. In our own series of fifty samples blood was withdrawn by means of a capillary pipette from the main lot, for purposes of culture, the day after collection. Four tubes containing heart broth were inoculated, two with 2 c.cm. and two with 1 c.cm., one from each set being incubated aerobically

at 37° C. and the other two anaerobically. Cultures were plated out daily for fourteen days. Three cases of contamination with *Staphylococcus aureus* occurred in the first twenty of the fifty samples collected. The remaining thirty were sterile. It is possible that this high rate of contamination was in part due to technical factors which, as already discussed, were later eliminated. The contaminant usually grew in about half the tubes and did not develop until the sixth day of culture. Dr. T. C. Stamp, to whom we are indebted for arranging for these tests, believes that cultures should be kept under observation for at least one week before reporting a negative. Aratyunjan (quoted by Filatov, 1937) reports 8 per cent. of contaminations in a series of sixty-five samples; Shorina, 3.9 per cent. of infections (quoted by Filatov, 1937). Saxton reported a much higher rate of contamination in Spain. It is likely that under the stress of war-time conditions sterilization of apparatus was far from satisfactory. Any handling of the collecting apparatus, such as putting on a needle sterilized by boiling rather than using one attached to the rubber tubing before autoclaving, we found increased the contamination rate. Unless contamination was gross we were unable, on naked-eye examination, to detect any difference between sterile and infected blood. Biddle, who tested a few only of his 160 lots of blood, found them all sterile. Boland *et al.* (1939), out of forty samples of blood cultured, found two to be contaminated with *Staph. albus*.

Available evidence suggests that if the blood is withdrawn with a careful aseptic technique the risk of contamination is slight. In times of peace culture should be a routine. Under emergency conditions it would be impossible to employ this additional safeguard. Duran (1939), in his latest apparatus, stores the blood in contact with oxygen. He considers that anaerobes obviously cannot grow in these circumstances, while aerobes, if present, consume the oxygen, so that the blood, which should be bright red, becomes dark in colour. In this way, in his opinion, contamination is readily detected without the necessity for culture. We found one positive Wassermann reaction in a group of fifty donors; Biddle (1939) found none.

Method of Administration

Stored blood can be administered by ordinary gravity, by some form of pressure apparatus, or as a continuous drip. It must always be filtered. In Spain, we believe filtration was effected at the base soon after withdrawal of the blood (Duran *et al.*, 1937). We either poured the blood through sterile muslin when it was given by the continuous-drip method, or passed it through a round glass bulb containing glass-wool before it reached the patient's circulation when gravity or pressure apparatus was used. Boland *et al.* (1939) used about 40 c.cm. of glass shot (diameter 2-3 mm.), added to each bottle before autoclaving. It is essential to mix and warm the blood before administration. This can be done by standing the container in water at 40° C. for about fifteen to twenty minutes and rotating it gently at intervals. It is also an advantage to have some form of closed container for keeping the blood warm during administration. A metal water-bath kept in a cork-lined plywood case serves this purpose well. Special care is needed in cross-matching, since there is some evidence of a definite decrease in the agglutinin content of preserved red cells. Duran (1939) pooled all blood taken from Group IV donors and sent out mixed lots for therapeutic use. He found this method satisfactory, and it obviated all need for cross-matching.

Transport

If the best results are to be obtained with stored blood it is essential that it should be kept at a low temperature until immediately before use. Blood that has once been warmed even for a short time tends to deteriorate more rapidly than blood kept at a steady low temperature. Special refrigerator vans were used successfully in Spain for the transport of stored blood up to front-line hospitals. In designing such transport vans it is important to remember that stored blood should not be shaken more than is absolutely necessary. Transport vehicles should therefore be fitted to carry containers in a steady upright position. For the transport of small amounts of blood in times of peace the metal water-bath and cork-lined box used for keeping the blood warm during administration may equally well be used for keeping it cold.

Indications for Transfusion of Blood

The relative merits of gum saline, glucose saline, and blood transfusion are always a subject of debate. The chief value of transfusion is its immediate effect in increasing blood volume, plasma proteins, haemoglobin concentration, blood platelets, and other factors concerned in blood coagulation. In some instances it may act as a stimulus to blood regeneration (Vaughan, 1936). In time of war, transfusion is required primarily for patients who have had recent severe blood loss. The indications for transfusion in such cases must be largely clinical, since owing to the concentration of blood that occurs the red-cell count and haemoglobin estimation may be normal or even polycythaemic. Restlessness, pallor, dyspnoea, a systolic blood pressure of 70 mm. Hg or less, and a rising pulse rate are indications that blood transfusions are needed. Transfusions of fresh blood are also of definite value in patients with prolonged sepsis and anaemia, due, for instance, to a gunshot wound that fails to heal. Experience in Spain suggests that possibly stored blood is not so effective in such cases, but carefully controlled observations on this point are still lacking.

Experience has shown that stored blood is life-saving both in time of war (Saxton, 1937) and in time of peace (Elliott *et al.*, 1939). It is clear, however, that much still remains to be learnt about the most satisfactory methods of preservation, about the changes that occur during storage, and about its value in conditions other than sudden haemorrhage.

I must express my gratitude to Dr. Duran Jorda for the information he has given me about his experience with stored blood in the Spanish war.

REFERENCES

- Bagdassarov, A. (1937). *Sang.*, 11, 466.
 Biddle, E. M. C. (1939). *Lancet*, 1, 405.
 Biddle, E., and Langley, G. F. (1939). *British Medical Journal*, 1, 555.
 Boland, C. R., Craig, N. S., and Jacobs, A. L. (1939). *Ibid.*, 388.
 Domanig, E. (1937). *Wien. klin. Wschr.*, 50, 1067.
 Duran, Jorda F. (1939). Personal communication.
 — (1939). *Lancet*, 1, 773.
 — *et al.* (1937). *Rev. Sanid. Guerra*, 1, 329.
 Elliott, G. A., Macfarlane, R. G., and Vaughan, J. M. (1939). *Lancet*, 1, 384.
 Ellis, R. W. B. (1937-8). *Proc. roy. Soc. Med.*, 31, 684.
 Filatov, A. N. (1937). *Vestn. klin. Khir.*, 51, 194.
 Goodall, J. R., Anderson, F. O., Altimas, G. T., and MacPhail, F. L. (1938). *Surg. Gynec. Obstet.*, 66, 176.
 Halbrecht, J. (1939). *Lancet*, 1, 202.
 Howkins, J., and Brewer, H. F. (1939). *Ibid.*, 132.
 Page, A. P. M., Seager, K. G., and Ward, E. M. (1939). *Ibid.*, 200.
 Saxton, R. S. (1937). *Ibid.*, 2, 606.
 — (1939). *Ibid.*, 1, 405.
 Vaughan, J. M. (1936). *The Anaemias*, 2nd ed., London.
 Yudin, S. S. (1937). *Lancet*, 2, 361.

PREVENTION OF BLINDNESS

GENERAL ASSEMBLY OF INTERNATIONAL ASSOCIATION

A general assembly of the International Association for the Prevention of Blindness, which has its headquarters in Paris, was held in London on April 19 during the period of the annual congress of the Ophthalmological Society of the United Kingdom, the proceedings of which were reported in the *Journal* last week (p. 887). Dr. P. BAILLIART, the president, gave a brief address of welcome, and presented the gold medal of the association to Dr. PARK LEWIS, who, in accepting it, recalled the history of this movement. In 1909 at the International Ophthalmological Congress at Naples a proposal was made for an organization of this kind, but not until 1929, at the congress held in Holland under the presidency of Professor Van der Hoeve, did the association come into being, along with the International Organization against Trachoma.

Prevention of Ophthalmia Neonatorum

The meeting then received a report by Dr. A. H. H. SINCLAIR (Edinburgh) on ophthalmia neonatorum in Great Britain, particularly in Scotland, where, he said, the new rules framed by the Central Midwives Board for that country provided for the application of the Credé method of prophylaxis, midwives being required to take precautionary measures, including the instillation of 1 per cent. silver nitrate drops. In view of the fact that blindness or serious impairment of vision still occurred it was considered that these drops should be used as a routine measure for prophylaxis; no substitute had an efficiency equal to that of silver nitrate. The arguments against the use of silver nitrate by midwives were no longer valid, as their training had now reached a standard sufficiently high to make the method safe in their hands. He also discussed the treatment when the disease had become established, and expressed the hope that M & B 693 for gonococcal ophthalmia, and prontosil for pneumococcal and streptococcal cases, would fulfil their early promise, and that venereal disease in the mother, which at present constituted the greatest menace to the eyesight of the child, might be brought under control.

Professor VON SZILY (Germany) declared that the prophylactic method of Credé should be applied thoroughly and conscientiously in every case, as prescribed in the regulations for midwives in the Reich since 1920. The compulsory notification of all cases of gonococcal ophthalmia of the newborn should be strictly observed. This method offered the best assurance that all suspected cases of blennorrhagia received special treatment at an early date. Dr. R. P. WILSON (Egypt) touched on the interesting fact that ophthalmia neonatorum is rare in Egypt. It had been for centuries the tradition and superstition of the native midwives to put either a mixture of onion juice and salt or the juice of a lemon into the eyes of newborn children. This strange and crude practice might be an important factor in the prevention of ophthalmia neonatorum in Egypt and surrounding countries where it was also followed, seeing that lemon juice had a high percentage of free citric acid, and onion juice contained allyl sulphide, which also had powerful germicidal properties.

Professor F. TERRIEN (France) said that on the basis of replies from almost all the French maternity clinics it appeared that the instillation of silver nitrate, generally 1 per cent., immediately after birth and before the cord had been ligatured, was carried out in almost every case. The desirability of this measure was laid down by the Académie de Médecine in 1937. Dr. J. D. M. CARDELL recalled the fact that when in 1881 Credé introduced his method it was in the form of 2 per cent. silver nitrate drops instilled into the eyes of the newborn child, thus at once achieving a dramatic reduction in the incidence of ophthalmia neonatorum, the incidence in the Leipzig lying-