A COMPARATIVE STUDY OF THE TUBERCULIN "PATCH TEST" AND THE INTRACUTANEOUS MANTOUX TEST IN CHILDHOOD

BY

DONALD COURT, M.B., M.R.C.P., D.C.H.

*Medical Registrar to the Children's Department, Westminster
Hospital*

(WITH SPECIAL PLATE)

The object of this brief communication is to present certain observations on a recent advance in the method of detecting tuberculous infection. From the discovery of tuberculin in 1890 to the present time a growing number of tests have appeared having as their common factor the allergic response of infected individuals to the products of bacillary growth. The more significant of these included the original subcutaneous reaction of Koch, the "Stichreaktion" or "needle-track" reaction of Epstein and Escherich, the cutaneous test of von Pirquet, the conjunctival approach of Calmette, the percutaneous or inunction test of Moro, and finally the intracutaneous test of Mantoux and Moussu. This march of events was naturally towards a greater diagnostic accuracy. From the von Pirquet reaction, with a potential error according to Hamburger of 40 per cent., we reach the present position where, by using the intracutaneous or Mantoux test, the error in negative reactors is but little over 2 per cent. The position is even more emphatically stated by Opie, who claims that the intracutaneous test is the only means by which sensitivity to tuberculin can be measured so accurately that two successive tests can be compared.

The increase in accuracy having been achieved, the next issue was naturally a simplification in technique. The only previous departure from the "inoculation" method, in its several varieties, were the conjunctival approach of Calmette and the inunction test of Moro. In the latter the skin was rubbed with ether to remove fat, and a concentrated old tuberculin was then applied in ointment. This method, though more sensitive than the von Pirquet reaction, still had an error of 15 to 20 per cent. (Lovett). Nevertheless in recent years there has been a return to the percutaneous approach. The modification referred to in this communication is the Vollmer (Lederle) patch test.

Technique of the Vollmer Patch Test

The patch consists essentially of a small strip of adhesive plaster 4 inches by 1 inch, on which are set three squares of filter paper (Plate, Fig. 1). The two outer squares have been saturated with undiluted old tuberculin, the centre with glycerin broth, to act as a control. Two groups of children were investigated in the wards and out-patient department of the Hospital for Sick Children, Great Ormond Street. Two tests were carried out on each child.

1. *The Mantoux Test.*—Into the skin of the right forearm 0.1 c.cm. of 1:1,000 O.T. (old tuberculin) was injected intradermally, and the resulting area of erythema and induration read at the end of forty-eight hours. Unless this area possessed one diameter of 10 mm. or more the test was considered negative.

2. *The Vollmer (Lederle) Patch Test.*—An area of skin over the sternum, or in younger children in the interscapular region, is cleansed with ether or acetone. The chosen site is allowed to dry, and the patch, after removal of the crinoline, is applied, and left in place for forty-eight

hours. Washing of the adjacent parts during this period should be avoided. The strip is then removed, and the result read after a further twenty-four hours. The reason for this delay is twofold. First, the reaction is more pronounced at this time, and a result that would appear equivocal on first removing the patch is found to be definitely positive. Secondly, any erythema due to irritation by the plaster itself will have subsided. It may be noted that the latter occurred in only two of the cases tested, and never gave rise to difficulty in interpretation. A positive reaction may appear as a sharply circumscribed, indurated, and reddened square set with small vesicles (Plate, Fig. 2). In other cases only a few discrete pin-head papules were present. Usually both squares reacted, but occasionally only one square did so. The central square remains unchanged. It must be stressed that any reaction, however slight, within the limits of the test squares is significant. The degree of the reaction would appear to vary with the type of skin, and bears no obvious relation to the extent or activity of the tuberculous infection present.

The Present Investigation

Two groups of children were investigated by the intradermal and percutaneous methods, as outlined above. The first were taken in a routine fashion without regard to age, sex, or clinical diagnosis. The ages ranged from 11 months to 12 years. The results (110 cases) were as follows:

Mantoux-positive ..	31—28%	Patch-positive ..	31—28%
Mantoux-negative ..	79—72%	Patch-negative ..	79—72%

In this group, therefore, there was no discrepancy between the two methods. In only one case did the application of a second patch prove positive when the first had given a negative result. "Plaster reactions" occurred in two cases, but gave no difficulty in interpretation.

The second group consisted of 100 children in whom a positive reaction to 1:1,000 O.T. intradermally was established. Comparison with the Vollmer patch test here gave the following results in 100 cases: Mantoux-positive, 100; patch-positive, 98; patch-negative, 2. In this group two children gave a negative response at first, but were positive with a second patch. The reason for this is not clear, but it is assumed that the absorptive power of the skin depends upon its texture and on the efficiency of fat removal during the preliminary cleansing with ether or its equivalent. A defect in the patch is much less likely than the preceding factors, which are obviously capable of some variation in different individuals. In the first child in which discrepancy occurred between the two tests the percutaneous method gave a negative result on three separate occasions. Generalized tuberculosis was present, however, and the Mantoux reaction itself was only weakly positive, indicating the depression of tuberculin sensitivity which is known to occur under such conditions. In the case of the second child no adequate explanation was forthcoming.

These results agree in the main with those from other similar investigations. In Monrad's (1936) series of 1,720 children, 1,362 were Mantoux- and patch-negative, 341 were Mantoux- and patch-positive, and seventeen were Mantoux-positive and patch-negative. There was agreement in 99 per cent. of the whole group, but in only 95 per cent. of the Mantoux-positive series. Wolff and Hurwitz (1937) tested some 936 children under 16. They found that in active cases there was no discrepancy between the two methods, but a difference of 1.8 per cent. in the series as a whole. In a group of 536 cases Dudley Hart (1938) found that 100 were Mantoux-positive and ninety-seven patch-positive.

Vollmer and Goldberger (1937), the pioneers in this work, established the superiority of the patch test over that of von Pirquet. In a recent communication—September, 1938—using a method similar to that described above, their results were as follows: Of 169 children with active tuberculosis only one failed to react positively to the patch, though Mantoux-positive—a discrepancy of less than 1 per cent. In 118 routine cases the Mantoux test with 1:1,000 O.T. did not reveal a single case of infection that was not already discovered by the patch test. These last results compare closely with those obtained above.

Advantages of the Patch Test

1. *Simplicity in Technique.*—It cannot be denied that the Mantoux reaction is reliable only if the injection is a true intradermal one, with exact amounts of material. The patch test, on the other hand, demands a minimum of manipulative skill. This, though obvious, is none the less important in routine work, notably in general practice. At the same time it obviates yet another of the "pricks" to which children are subjected in these days of increasing investigation. There seems to be no tendency on the part of the child to interfere with the patch once it is applied.

2. *Durability of Material.*—For solutions of old tuberculin to be reliable they must be renewed at least every two months. The patch test material, if kept in a cold dry place, will remain reliable for a year.

3. *The Absence of Undesirable Complications.*—General and focal reactions never occur, and necrosing local reactions are unknown.

The Disadvantages

1. The patch test is slightly less reliable than the intracutaneous method. Nevertheless from the present series, and those reviewed in the literature, it is clear that at least 95 per cent., and at most 98 per cent., of cases infected by the tubercle bacillus can be exposed by this method. This leaves 2 to 5 per cent. in which further investigation will be required. In such cases the patch test should be supplemented by the Mantoux, using a 1 in 100 dilution. If the discrepancy still remains, the latter, because of its greater range of sensitivity, should be taken as the reliable indicator.

2. Severe local skin reactions have been reported. I have no personal experience of this whatever. In the main, the patch test would appear to produce a less intense reaction than the corresponding Mantoux. Furthermore, there is little or no associated pruritus, and the erythema fades within two to three weeks at most.

Conclusions

The Vollmer patch test approaches in reliability the intracutaneous method of Mantoux.

In a very high proportion of cases it will reveal the presence of tuberculous infection. For the remaining minority a combination of the patch test and the Mantoux, using stronger concentrations of old tuberculin, or its equivalent, will prove satisfactory.

My thanks are due to Dr. Donald Paterson for his help and permission to use his patients. The "patch tests" were very kindly supplied by Chas. F. Thackray of Leeds.

BIBLIOGRAPHY

- Hart, F. Dudley (1938). *Lancet*, 2, 609.
Hart, P. D'Arcy (1932). *Med. Res. Cncl. Spec. Rep. Ser. No. 164*.
Monrad, S. (1936). *Ugeskr. Laeg.*, 98, 885.
Vollmer, H., and Goldberger, E. W. (1937). *Amer. J. Dis. Child.*, 54, 1019.
— (1938). *Ibid.*, 56, 584.
Wolff, E., and Hurwitz, S. (1937). *J. Amer. med. Ass.*, 109, 2042.

THE DIAGNOSTIC VALUE OF THE INTRADERMAL TUBERCULIN TEST IN CHILDREN

BY

D. B. BRADSHAW, M.B., D.P.H.

Immunological Officer, Hospital for Sick Children, Great Ormond Street

The graded intradermal tuberculin test was introduced by Mantoux in 1910, and has for some years been accepted, at least in this country, as the most reliable tuberculin test. Although the information afforded by the result of the test in any given case is usually clear, it is difficult from the figures available to answer the question of how useful the test is as a general diagnostic measure, and it is to provide this information that the following analysis has been made. The figures heretofore published relate to special groups, such as children from tuberculous homes (Schlesinger and Hart, 1930), the clinically tuberculous (Hart, 1932), the clinically non-tuberculous (Hart, 1932), and children attending a special (chest) hospital (Dow and Lloyd, 1931); nor are figures compiled in other areas or relating to normals directly relevant. The fundamental facts underlying the use of the Mantoux test for diagnostic purposes are:

1. The test is positive in almost 100 per cent. of proved cases of tuberculosis (moribund cases excluded) (Hart, 1932).

2. A positive reaction is proof of past or present tuberculous infection, but it can never be direct evidence that active tuberculosis is present or that any given lesion or symptom is tuberculous in origin.

3. It follows from (1) and (2) that the value of the Mantoux test lies mainly in the exclusion of tuberculous infection which is implied by a negative reaction. Its limit of usefulness is therefore determined by the percentage of positive reactors among the general population. Where this is low, as in infants and young children, the test has found its chief application; where high, as in the adult population, the test can seldom yield useful information.

In its diagnostic application the test is used in patients presenting symptoms or signs that are probably or possibly tuberculous in origin. In the patently tuberculous and non-tuberculous the test is not required for diagnosis. The value of the Mantoux test therefore depends upon the proportion of negative reactors in the group of "probably or possibly tuberculous." This proportion is limited by the factors set forth above, but will vary from one community to another and even within the same community according to age groups, social standing, etc. In addition the increasing use of safe milk, segregation of open cases, and improvement in social conditions should lead to a progressive drop in the number of those who have become tuberculin-sensitive. The test should therefore be more useful with the passage of time. In the light of what has been said there can be no absolute or final assessment of the value of the Mantoux test for diagnosis, and the view is widely held that it is only worth while performing the test in the younger children.

The Present Investigation

At the Hospital for Sick Children the test is extensively employed, and we have for some time had the impression that it is of greater value than is usually accepted. The present analysis has amply borne out that impression. The figures under review relate to Mantoux tests carried out in the out-patient department between 1934 and mid-1938. Except where otherwise stated the tests were done

A. TUDOR EDWARDS: MODERN PRINCIPLES OF TREATMENT IN BRONCHIECTASIS

FIG. 6.—Right middle and lower lobes after lobectomy, showing advanced bronchiectasis secondary to a chronic middle-lobe abscess.

FIG. 7.—Large single infected congenital cyst after lobectomy of the right upper lobe in the case referred to in Fig. 5.

FIG. 8.—Left lung after pneumonectomy in the case referred to in Fig. 4.

DONALD COURT: TUBERCULIN "PATCH TEST" AND
INTRACUTANEOUS MANTOUX TEST IN CHILDHOOD

FIG. 1.—Three squares of filter paper on adhesive plaster; the outer squares are saturated with undiluted old tuberculin, the centre square (control) with glycerin broth.

FIG. 2.—A positive reaction twenty-four hours after removal of the patch.