are affected, the main blood supply being unaffected. It is of interest that in this type of case the infection does not readily tend to spread from one group of muscles to another.

Treatment

Prophylaxis by means of anti-gas-gangrene serum or the use of this serum after the infection has been established is in my opinion of little or no value, though it is obviously difficult to substantiate this statement fully in the case of prophylaxis. The giving of anti-gas-gangrene serum at the same time as anti-tetanic serum is advisable therefore, and though it may do no good it can do no harm. The best form of prophylaxis is the early and careful débridement of the wound, all dead tissue, especially muscle of doubtful vitality, being removed and the wound being left open for some days if there is obviously risk of severe infection. In established cases the line of treatment depends on the nature and extent of the infection: where there is massive gangrene in a limb with damage to the main artery early amputation is essential. Such amputations should be well above the limits of the infected area; if flaps can be fashioned they should be ample, and it is advisable that they should not be sutured for some days. Exposure of the stump to the air with the minimum amount of dressings is also of extreme value. The immediate amelioration in the patient's condition after amputation for gas gangrene is most remarkable.

In group gangrene, where a muscle or a group of muscles are attacked, resection of the affected area will often cut the infection short and save both limb and life. I will give two examples of such cases to illustrate the method:

Case 1.—This patient was wounded at 1 a.m., the right leg being shattered in the lower third. He was admitted at 3.30 a.m., and a primary amputation was performed by equal lateral flaps in the upper third of the leg at 4 a.m. The flaps were lightly sutured and provision was made for continuous eusol irrigation. All went well for twenty-six hours; the pulse then ran up to 120 (it had previously been 80 to 90), the tongue became furred, and the patient was drowsy and obviously very ill. On examination of the limb the front of the leg showed mottling of the skin, with swelling and a tympanitic note on percussion. The patient was at once anaesthetized and the wound was opened up, when it was found that the entire anterior tibial group of muscles in the stump were brick-red in colour and non-contractile. They were removed *en masse*, and the wound was left open and treated by continuous eusol irrigation. After twenty-four hours the patient's condition improved steadily, and he was evacuated to the base nine days later. At the first operation it was noted that the anterior tibial artery was occluded but that the anterior tibial muscles were apparently healthy and contractile: the arterial occlusion clearly predisposed to the development of the gangrene at a later date. In this case it is doubtful whether the patient would have survived an amputation through the thigh. Local excision of the infected muscle group gave him a useful stump with a normal knee-joint.

Case 2.—This patient was wounded at 4 p.m. by a rifle grenade fragment and admitted three and a quarter hours later as a walking case. When he arrived he was clearly very ill: his pulse was 120, of poor quality; the tongue was furred and the face pinched. There was an irregular entry wound the size of a threepenny-piece on the posterior aspect of the right arm just above the internal condyle of the humerus; there was no exit wound. The whole arm appeared swollen and was crepitant to the touch; there was a tympanitic note on percussion. He was operated on at once, when it was found after excision of the wound of entry that the projectile was lying to the inner side of the vessels in the

lower third of the arm. A second incision was made over the site of the projectile, and subcutaneous tissue was found to be oedematous and the biceps muscle showed signs of gangrene. The whole biceps was therefore exposed by a long incision and the inner half was found to be changed in colour and non-contractile: the outer half of the muscle appeared to be normal. As the infection seemed to be localized to the inner half of the muscle this portion was excised in its entirety. The wound was left open for seven days, and was then closed by secondary suture. The ultimate result was extremely satisfactory.

BIBLIOGRAPHY

- Cole, L. (1936). *British Medical Journal*, 1, 1191.
 — (1938). *Proc. roy. Soc. Med.*, 31, 1205.
 Frankau, C. H. S., Drummond, H., and Neligan, G. E. (1917). *British Medical Journal*, 1, 729.
 Wallace, C. (1917). *Ibid.*, 1, 725.

THE ORGANIZATION OF OUT-PATIENTS AT A TEACHING HOSPITAL

BY

WALTER HOWARTH, M.A., M.B., F.R.C.S.

Surgeon to the Ear, Nose, and Throat Department,
St. Thomas's Hospital

It has for a long time been a reproach that at the large London hospitals, in which very considerable numbers of out-patients are dealt with each day, the patients may have to wait, and indeed often do wait, many hours before their case is finally attended to and they are free to go home. A considerable period of waiting may also be entailed when patients come up regularly for treatment, and it is sometimes the case that a patient may wait several hours before being seen. If the patient is at work and time is valuable, a visit to the hospital may well mean the loss of half a day, and if this has to be repeated at frequent intervals it may give rise to some dissatisfaction on the part of the employer as well.

From time to time attempts are made to expedite the examination of patients, either by having a continuous "sorting" system or by the appointment of senior officers qualified to deal with the cases as they arise and so obviate the necessity for the patient to wait to be seen by the visiting surgeon or physician at a later hour. Improvements have undoubtedly been effected by these and other means, but the lot of the patient who comes for regular treatment (and these form the bulk of the hospital out-patient attendances) has not improved very much. In my department at St. Thomas's Hospital a scheme has been worked out and has been put to a practical test with such satisfactory results that it seems worth while to draw attention to it in the hope that others may think it worthy of a trial. The details of the scheme have been largely worked out by the lady almoner to my department, Miss Turner, and by my colleague, Mr. Bateman, and the greatest credit is due to them.

The System at St. Thomas's

In a teaching hospital, not only have the patients to be examined, diagnosed, and treated, but some also have to be reserved for teaching purposes and examined by the students. A certain number of those who attend for regular treatment have also to be examined with the students, so that the results of treatment may be followed up and demonstrated. This necessity for teaching—of vital importance, I think, from the student's point of view—naturally complicates and often delays the actual treatment of the patients, so that any plan that ensures a more even distribution of time is an advantage for everyone concerned. A careful examination of the time-sheet will show the method that has been adopted, and a mere glance at the accompanying typical daily time-sheet that has

been filled in by the lady almoner—upon whom the bulk of the work falls—will give an idea of the orderly arrangement that prevails. (The handwritten details, which are the bureau numbers of the patients, are printed in italics.) A few words of explanation are, however, necessary.

St. Thomas's Hospital						
Ear, Nose, and Throat Department						
Surgeon's Name: MR. HOWARTH.				Date		
VISITING SURGEON	CH. ASSIST.	H.S.	C.A.1.	C.A.2		
1.30			Examine New Cases	Examine New Cases	Examine New Cases	1.30
2.0	Sorting new Cases with H.S. and C.A.s	Old Cases 1. 16,321 2. 27,832 3. 22,560 4.	New Cases with Visiting Surgeon	New Cases with Visiting Surgeon	New Cases with Visiting Surgeon	2.0
2.15		5. 15,519 6. 45,395 7. 4,943 8. 41,284				2.15
2.30	Teaching	Referred Cases 1. 38,844 2. 47,560 3. 4.	Old Cases 5,988 19,608	Old Cases 1. 35,222 2. 17,345 3. 28,814 4. 21,806	Old Cases 1. 47,493 2. 1,037 3. 39,828 4.	2.30
2.45		5. M. Box 6. R. Day 7. 8. 4,640	47,335 24,473	5. 3,828 6. 49,037 7. 8. 48,695	5. 34,454 6. 45,176 7. 8. D. Jones	2.45
3.0		9. 51,805 10. 11. 12.	40,326 45,188	9. 17,881 10. 48,051 11. 18,387 12.	9. 11,600 10. 37,824 11. 22,164 12.	3.0
3.15		Old Cases 9. 46,605 10. 38,983 11. 14,676 12.	42,073 3,618	13. 37,343 14. 48,613 15. 45,125 16.	13. 48,257 14. 247 15. 4,445 16. 47,252	3.15
3.30	No Booking Discussion	13. 38,993 14. 47,809 15. 16.	46,549	17. 49,807 18. 31,576 19. 43,223 20.	17. 44,877 18. 49,037 19. 33,488 20.	3.30
3.45	Ward Cases and Nurses and Stu- dents, &c.	17. 7,181 18. 46,549 19. 48,964 20.	48,160	21. 24,938 22. 52,204 23. 24.	21. 51,438 22. 42,795 23. 24.	3.45
4.0	Special Cases 2,053 29,647 35,164 1,268			25. 46,145 26. 49,467 27. 28.	25. 24,584 26. 51,578 27. 28.	4.0
4.15	47,017 51,973		52,240	43,242 11,680	43,184	4.15

The staff of the department on an out-patient day consists of the visiting surgeon, the chief assistant, the house-surgeon, and two (or more) clinical assistants. The session begins at 1.30 p.m., and by that time the new patients sent up by doctors have been to the central bureau and received their bureau number (for follow-up and filing purposes) and their "letters," on which details of their case will be written. These cases are examined by the house-surgeon and the clinical assistants in the first place. When the visiting surgeon arrives he goes through these cases with the house-surgeon and the clinical assistants for the next half-hour, checking their observations and selecting cases that he thinks may be useful for teaching. Cases that are not required for teaching are referred to the chief assistant. The visiting surgeon then teaches the students on the selected cases for the next hour, and after that there may be a short discussion on some unusual problem that has arisen. Cases referred from the wards are then seen, as well as any nurses or students who may require examination and treatment. Lastly, special cases that the surgeon particularly wishes to keep in touch with are seen. These cases will have

been given a card by the lady almoner with the time, say 4.15 p.m. or later, written on it. The ward cases, nurses, and students would not come during the teaching hour, but would arrive at a definite time after this.

While the visiting surgeon is engaged in the way that has just been outlined, the chief assistant would be dealing with the remainder of the new cases and then, later in the afternoon, with old cases which were in the habit of coming up to receive his personal treatment. These would arrive in batches of four at times corresponding to the times written on their cards. The chief assistant decides whom the patient should see at his next attendance and fills in the name (surgeon, house-surgeon, etc.) on the last line of the card, the appropriate time being added later by the lady almoner, who arranges the rota of attendances for subsequent out-patient days. Meanwhile the house-surgeon and the clinical assistants are busy seeing other old cases, who also arrive in batches of four at times stated on their cards, so that their period of waiting is reduced to a minimum. The lady almoner will have arranged an early or a late appointment to suit the convenience of the patient.

The time of the house-surgeon is not fully booked up each quarter of an hour with old patients, as in this particular department his activities (myringotomies, puncture of antra, cauterization of noses) cannot be definitely circumscribed. The clinical assistants are fully occupied with the treatment of the old cases as they arrive at their stated times.

When the patients leave the department they go to the lady almoner's room and she arranges the times of their subsequent visit, noting from their card which officer they are to see. It will thus be observed that practically all the organization of attendances falls upon the lady almoner, while the successful working of the scheme depends on the punctuality and regularity of attendance of the visiting surgeon and other officers concerned.

THE NEXT STAGE IN EVACUATION

The Minister of Health has sent to all local authorities concerned in the Government's Evacuation Scheme in England and Wales a communication (Ministry of Health Circular 1800, H.M. Stationery Office, 1d.) asking them to work out in detail their plans for the evacuation and reception of school children and others who are to receive priority under the scheme. Enclosed with this communication is a memorandum (Memo. Evacuation No. 4, 3d.) indicating the general lines of the action which would be necessary in an emergency, the problems likely to arise, and the manner in which they can best be dealt with.

Practically all the returns showing the results of the evacuation survey have been received and examined. The receiving and evacuating authorities can now, in contact with one another and with the evacuation authorities, work out their plans in greater detail in the light of information on the numbers in the priority classes and the accommodation available.

Course of Action in an Emergency

The scheme is to provide for the transference from the crowded areas of the great cities of some three million persons, including school children, young children with their mothers, expectant mothers, blind persons, and cripples. Having decided which portion of their area and which schools in the area are to be included in the evacuation plans, and having obtained the approval of the Minister to their proposals, evacuating authorities must ascertain the numbers of persons in the priority classes who would desire to be evacuated in the event of an emergency.

The success of plans for the evacuation of school children will depend largely on the co-operation of teachers, and autho-