positive (Figs. 3 and 4). In sanatorium for ten months, bed rest for six months. Now no sputum; considerable resolution.

Case IV.—Female aged 16. Adolescent phthisis. Large cavity in right upper zone with fluid level; sputum positive (Fig. 5). Artificial pneumothorax attempted, contraselective collapse obtained. A.P. discontinued. Bed rest in sanatorium for eight months. Several skiagrams during this period showed cavity becoming progressively smaller. Now completely healed (Fig. 6); no sputum, excellent general condition.

A SPUTUM-NEGATIVE CASE

Case V.—Female aged 45, first seen July, 1935. Cough, loss of weight (1 stone in two months), and lassitude for the last six months. Clinical and x-ray investigations showed evidence of infiltration of the upper zones of both lungs. Originally no sputum, but subsequently sputum was examined by all available methods and found negative for tubercle bacilli. On routine sanatorium treatment for twenty-four weeks and regular supervision subsequently. In May, 1937, no abnormal physical signs. X-ray examination showed complete resolution of the lung lesion.

Is Case V to be regarded as a chronic infiltration of lung tissue casting shadows identical with those of pulmonary tuberculosis and not apparently to be connected with the tubercle bacillus? The chronicity of this case shows that it is not quite analogous with those described by Scadding; but recent work reveals how much we have to learn about simple inflammatory processes in the lung, and that caution is necessary before results are attributed to a particular type of treatment.

Summary

A discriminating view on the treatment of pulmonary tuberculosis has been suggested, bearing in mind the two aspects of the disease—namely, the constitutional aspect and the local manifestation in the lung.

The tendency of certain fresh tuberculous pneumonic and bronchopneumonic lesions, with or without cavities, to heal spontaneously has been outlined.

The desirability of treating such cases by strict and subsequently modified bed rest with clinical and radiological control has been emphasized.

The prolonged time factor for the successful treatment of such cases by general and local rest and the importance of the complete co-operation of the patient are outlined.

The immediate institution of collapse therapy in such types of localized lesions is not justified, but may demand serious consideration if with serial x-ray control the lesions remain stationary or retrogress.

On the other hand, collapse therapy in cases of the productive type of disease should not be unduly delayed, and in certain cavity cases may prove to be a life-saving device.

BIBLIOGRAPHY

- Amberson, J. B. (1936). *Amer. Rev. Tuberc.*, **33**, 269.
 Ameuille, P. (1926). *Progr. méd.*, **41**, 1933.
 Bacmeister, A. (1927). *Beitr. klin. Tuberk.*, **67**, 157.
 Brooks, W. D. W. (1938). *Brit. J. Tuberc.*, **32**, 14.
 Coryllos, P. N. (1933). *Amer. Rev. Tuberc.*, **28**, 1.
 — and Ornstein, G. G. (1938). *J. thorac. Surg.*, **8**, 10.
 Ellman, Philip (1932). *Chest Disease in General Practice*, H. K. Lewis, London.
 Fales, L. H., and Beaudet, E. A. (1931). *Ann. Rev. Tuberc.*, **23**, 690.
 Hebert, G. T. (1939). *Tubercle*, **20**, 145.
 Jaquerod, M. (1930). *Cure de Repos*, Masson et Cie, Paris.
 — (1936). *Rev. Tuberc.*, 5 s., **2**, 1102.
 — and Ellman, Philip (1927). *Brit. J. Radiol.*, **32**, 343.
 McMahon, B. T., and Kerper, E. H. (1933). *Amer. J. med. Sci.*, **186**, 170.
 Paterson, Marcus S. (1911). *Auto-inoculation in Pulmonary Tuberculosis*, London.
 Scadding, J. G. (1937). *British Medical Journal*, **2**, 956.

THE CAUSAL ORGANISM OF SOUTH AFRICAN "SANDWORM" ERUPTION

A PRELIMINARY NOTE

BY

N. L. MURRAY, B.Sc., M.B., B.Ch., D.P.H.

Union Health Department, Durban

(WITH SPECIAL PLATE)

The "creeping eruption," or "sandworm" as it is known in South Africa, is a very common condition, particularly in Natal and Zululand. Up to the present time its aetiology has never been satisfactorily established. The purpose of this article is to illustrate a hitherto undescribed causal organism and to describe the procedure of isolating it.

The affliction is characterized by a thin inflammatory raised line or track about one-eighth of an inch wide, which increases in length as the parasite burrows its way in a tortuous manner through the superficial layers of the skin and produces a severe irritation (Plate, Fig. 1). The feet of children who play bare-footed are most commonly affected, but any part of the skin may be attacked, including the genital organs. The buttocks, too, are a favourite site. It affects natives as well as Europeans, and no age is exempt. The cases occur most often in the hot wet months of the year. Lay opinion that infection is incurred only on the seaside beach sand above high-water mark is not proved. In fact, evidence shows that persons who have not been near the beach at all have contracted it, and in one case seen recently it was associated with gardening. This man had been digging and transplanting cabbages at his home in Durban, where he developed a severe infection of both hands within a few days. The distribution in South Africa is not fully worked out yet, but cases seen inland have almost invariably been attributed to a holiday on the Natal coast.

The infection is spread by scratching the parts to relieve the pruritus, and sepsis usually follows. The sepsis may be severe, and it has led to permanent scarring; one subject managed to transfer the infection from one hand to the other and then to the face, which became very painful and swollen, particularly about the chin and lips. During recent weeks eight cases have come to my notice, and I had the good fortune to have two adults who were willing to submit to the tedious process of looking for the causal organism.

Procedure of Search for the Organism

A dissecting microscope, some cedarwood oil, and a sharp eye-spud were used. The cedarwood oil clears the skin somewhat and permits an even incident light to be focused on the track. The capillary loops of the true skin are visible, and appear dilated below the burrow, due to reaction to the irritation of the parasite and any eggs or debris produced by it. It is also possible that a secretion by the parasite in the form of a lysin assists the fairly rapid tunnelling through the epidermis and contributes to the severe itching complained of.

The raised reactionary track, which is palpable and obvious to the naked eye, is much wider than the true burrow through the cells. The obvious track is about 3 mm. wide, but the true track is only about 0.33 mm. (roughly the thickness of a fine dental needle) until it becomes septic. It is superficial in its entire extent and

may show small pockets at intervals where egg-laying probably occurs. When the track through the skin is located the cedarwood oil is wiped off with xylol and the parts cleaned with ether or spirit. Using the eye-spud, the track is then laid open to its terminus, taking care not to draw blood, which obscures visibility. The drops of clear fluid in the burrow are transferred to a slide and covered with a cover-slip for examination under an ordinary microscope. A drop of distilled water on a clean slide prepared with Mayer's egg albumen facilitates the procedure. In this way I was successful in obtaining what appear to be four stages in the development of a

Cross-section of "sandworm" track in human skin. Photomicrograph of Dr. F. G. Cawston's case (1928). The track lies entirely in the superficial layers of the epidermis and contains numerous lymphocytes. The stratum corneum has been separated by artefact. Serial sections of skin in two of my own cases revealed the track in an identical situation.

parasite hitherto undescribed as a cause of "creeping eruption."

The investigation was begun with the idea of looking for either a fly larva (*Gastrophilus* var. or *Hypoderma*) or a larval dog-hookworm (*Ankylostoma braziliense*), both of which are quoted in the textbooks as a cause. A cursory examination under the microscope made it obvious that from the size of the true track the causal organism was likely to be much smaller than expected. No sign of the above-mentioned larvae was evinced in any of about thirty tracks examined. A number of slides were prepared from droplets taken from various portions of tracks in the skin of a man infected with "sandworm" (Plate, Fig. 1), and examination of these revealed a mite about 300 μ in length and two developmental stages (Figs. 3 and 4). Tracks in a second case—that of "sandworm" of the hand (Fig. 6)—after a patient search produced a similar mite (female) (Fig. 5) and a number of eggs (Fig. 2). The minuteness of the parasite and its transparency when unstained make it very difficult to be seen in the skin. To realize the size, it should be noted that the mite could be dropped into a pinhole with room

to spare, and that its egg is little larger than a polymorphonuclear leucocyte.

Commentary

The attention of interested readers may be drawn to a note by Theobald which states: "It has been asserted by Haller that the Cape ailment (Port Natal sickness) is caused by mites, but this statement has been contested" (Fantham *et al.*, 1916). No mention is made of a "creeping eruption." The nearest resemblance, in the available literature, to the mite discussed in the present investigation is *Tetranychus molestissimus*, which is found in the Argentine and Uruguay on the under-surface of the leaves of *Xanthium macrocarpum*. It attacks mammals and men, producing severe itching, accompanied by fever in the latter (Fantham *et al.*, 1916).

The photographs of the various stages discovered must suffice for this article. Fuller details of the classification, life history, and occurrence are being prepared, and in a subsequent article the treatment will also be included.

Dr. J. Mackay, district surgeon, Durban, kindly referred suitable cases to me and provided the stimulus to investigate this common ailment. Mr. O. F. Gibbs, senior technical assistant, Government Laboratory, Durban, gave valuable advice in the photomicrography. I acknowledge my thanks for their assistance. I am also indebted to Dr. F. W. P. Cluver, deputy chief health officer, for permission to publish this report.

REFERENCES

- Cawston, F. G. (1928). *J. trop. Med. Hyg.*, 31, 201.
Fantham, Stephens, and Theobald (1916). *Animal Parasites of Man*, p. 488.

OBSERVATIONS ON THE PERFORMANCE OF BLOOD-GROUP TESTS

BY

G. L. TAYLOR,* M.D., Ph.D.

AND

ELIZABETH W. IKIN,* B.Sc.

(From the Galton Laboratory, University College, London)

Compared with other parts of the world blood-group investigations in this country have been few. In a recent series of papers Taylor and Prior (1938a, 1938b, 1939) have given some account of what has been done, have described in detail their investigations of family and unrelated material in England, and have shown by statistical analyses that their results agree with current genetic theory.

Below is a description of the technique used in the Galton Laboratory for the diagnosis of the blood-group factors A₁, A₂, B, O, M, and N, followed by observations on technical points mostly concerned with the M-N factors, the detection of which, authorities are agreed, is difficult and full of pitfalls for the inexperienced. At the present time there may be interest in such a communication coming from a unit which for more than three years has been devoted, almost exclusively, to work on blood-group factors. It may appear that the difficulties are overstressed and that the technique is elaborate; the workers in this unit are of the opinion that anything less well-controlled and less careful would not be good enough.

Blood-group tests on 1,073 unrelated persons, a sample we believe to be representative of the population in England, are reported and analysed; of these 1,073 tests, 422 have already been reported by Taylor and Prior (*loc. cit.*). Our technique undergoes slight modification from time to time, as in the case of the strength of red-cell

* Supported by the Rockefeller Foundation.

N. L. MURRAY: CAUSAL ORGANISM OF SOUTH AFRICAN "SANDWORM" ERUPTION

FIG. 1.—Case I. Adult European male affected with "sandworm." Ten days' duration. Note the tortuous track in skin over crest of ilium. Each of the other patches represents a track which has been scratched to relieve the intense irritation.

FIG. 2.—The egg of the causal organism, about $12\ \mu$ in size, lying among debris from the track.

FIG. 3.—Developmental stage of the organism—nearing adult size.

FIG. 4.—Adult mite; stained specimen, $300\ \mu$ in length, extracted from track shown in Fig. 1. The breadth has been somewhat increased by pressure of the cover-slip; two pairs of hind legs partly obscured. Note hairs and terminal claws on legs.

FIG. 5.—Case II. Adult female mite (unstained). Obtained from the hand.

FIG. 6.—Case II. "Sandworm" of the hand. Both hands affected, and very septic. Note raised track doubling on itself at tip of third finger.