of either hand fully; dorsiflexion of ankles relatively weak; gait much less waddling in nature; unable to climb stairs or rise up from floor yet; deep reflexes in upper limbs still much diminished, absent in lower limbs; abdominals absent. Weight 8 st. 3½ lb. Discharged for a month's holiday in the country.

November 19.—Feels stronger; can now ascend stairs, but has still some difficulty in getting up from the floor; no obvious muscular wasting; deep and superficial reflexes *in statu quo*. Weight 8 st. 8½ lb. December 3: Residual weakness now very slight; reflexes *in statu quo*. Discharged home.

TREATMENT

In addition to the usual measures, vitamin B₁ was given daily—4 mg. intramuscularly for ten weeks and 3 mg. orally thereafter. Dilute HCl was also given until the result of the second test meal was learned.

Discussion

Apart from a certain degree of mental disturbance the patient was in good health when taken ill. The discovery of achlorhydria was most intriguing, all the more so when a second gastric analysis, carried out when recovery was well on the way, showed free HCl to be present in amounts even greater than the average. The association of polyneuritis and achlorhydria has been stressed lately. Minot, Strauss, and Cobb (1933) investigated a series of fifty-seven cases of alcoholic polyneuritis. Only seven showed a normal free HCl content, fifteen secreted distinctly diminished amounts, and twenty-one secreted no free acid at all. Douthwaite (1936) recorded seven cases of "gastro-genous polyneuritis." Each was associated with a gastric disturbance involving a reduction or complete absence of free HCl. The cause of the achlorhydria in the present case is obscure. No other gastric symptom was elicited. One might theorize doubtfully upon the possibility of psychic disturbance having affected gastric secretion through the medium of the diencephalo-pituitary apparatus. Evidence is accumulating that some forms of polyneuritis, and even of localized neuritis, are in reality due to a deficient intake or absorption of vitamin B₁. In America alcoholic neuritis has been clearly shown to be produced by such a lack (Strauss, 1935, and later workers). The coexistence of achlorhydria in many such cases is surely more than a coincidence. Nevertheless one not infrequently encounters cases of polyneuritis in which the parenteral administration of the vitamin appears to have little effect. Recovery was not hastened very appreciably in the present case. There may be a deficient formation or absorption of some other neurotropic factor.

I am indebted to Dr. W. R. Reynell, senior physician to the West End Hospital, for permission to publish this case.

REFERENCES

- Douthwaite, A. H. (1936). *British Medical Journal*, 2, 535.
Minot, G. R., Strauss, M. B., and Cobb, S. (1933). *New Engl. J. Med.*, 208, 1244.
Strauss, M. B. (1935). *Amer. J. med. Sci.*, 189, 378.

The People's League of Health after an investigation extending over many months has now organized a scheme of clinical research with a view to determining the influence of diet on expectant and nursing mothers in relation to maternal mortality and morbidity. Systematic observation and records under expert direction are being collected at the ten London hospitals which are co-operating with the League in this research, and the results will be studied and published in due course. The League has just concluded the first year of its research, and it will probably be another six or nine months before the work is completed and the report published.

MYOMECTOMY IN THE PREGNANT UTERUS

BY

RUFUS C. THOMAS, F.R.C.S.Ed., M.R.C.O.G.

*Assistant Medical Officer of Health for Obstetrics and
Obstetric Consultant, County Borough of Croydon;
Obstetrician and Gynaecologist, Mayday
Hospital, Croydon*

DeLee, in his *Principles and Practice of Obstetrics* (p. 576), says: "Myomectomy has been frequently done during pregnancy, but it is a very bloody operation, and is followed by abortion in 17 per cent. of cases (Winter)." Speaking of the treatment of fibroids during pregnancy he says: "In the absence of unbearable pain, of severe haemorrhage, of very rapid growth, of great distress from over-distension of the belly, and of signs of necrosis of the tumour, or torsion, or acute abdomen, it is justifiable to wait."

By no means all fibroid tumours give rise to symptoms during pregnancy. All such cases, however, cause the obstetrician a certain amount of anxiety. It must be remembered that tumours which are easily palpable on the anterior or lateral uterine wall may be accompanied by others on the posterior wall, which are therefore inaccessible by the ordinary methods of examination. There is always the possibility that a fibroid low down in the posterior wall may cause obstruction of labour. For this reason in all cases where a fibroid is palpable I make an examination under a general anaesthetic at about the thirty-seventh week in order to be sure that such a condition does not exist. If such a growth is found the mode of delivery can be planned accordingly. Even after apparently safe delivery the anxiety is not over, as post-partum haemorrhage is not uncommon from inefficient retraction or where the placental site has lain over the situation occupied by the growth. The placenta may be adherent to a submucous fibroid, and may necessitate the "digging out of the two at once," with the danger of perforation of the uterus. Irregular haemorrhages which prolong the lying-in period are quite common, and the occurrence of pyrexia and offensive lochia following pressure necrosis of the tumour is well known and often gives rise to considerable anxiety. If symptoms appear some months before the expected date of delivery the possibility of a prolonged period of rest in bed, whether at home or in hospital, is one which the patient does not relish. If symptoms are acute the question of operative intervention must be considered.

The three cases reported below resemble each other in that pain was the chief symptom, and operation was undertaken on this account. The actual condition found at laparotomy was, however, different in each case. The following are the records.

Case I

A primipara aged 30 was admitted to hospital in great pain; this she referred to an area on the left side of the uterus, which was enlarged to about twenty-two weeks' pregnancy. A small mass, very tender, could be felt on the left uterine wall below the level of the umbilicus. She had been vomiting for several days, her tongue was furred, and she looked ill. On opening the abdomen a fibroid tumour the size of a large egg was found, having a broad base. An elliptical incision was made at the edges of the tumour, which was easily shelled out. Bleeding was temporarily controlled by hot packs, the cavity obliterated by mattress sutures, and the uterine incision closed by a continuous suture of No. 0 catgut. The hand was then

passed behind the uterus and both ovaries were found to be the seat of tumours about 4 inches in diameter, the two tumours lying side by side in the bottom of the pouch of Douglas. They were removed without difficulty, being easily delivered out of the abdomen with little disturbance of the uterus. The abdomen was closed in three layers, closure being reinforced with three figure-of-8 tension sutures of silk-worm-gut. Morphine was given during the next few days, but the patient showed no sign of aborting, and was discharged in three weeks. She had a spontaneous delivery at term, the child weighing 7½ lb.

The right ovarian tumour proved to be a dermoid, with the usual contents of hair, fatty fluid, and several teeth. The left tumour was a single cyst containing clear fluid. Both together would have been sufficient to cause obstruction during labour.

Case II

A primipara aged 29 had been attending the ante-natal clinic from the second month of pregnancy. I first saw her at the end of the fourth month, when she was in considerable pain and had a slight vaginal loss. There was a very tender swelling in the right iliac fossa, which she stated had increased considerably in size during the past few days. She was admitted to Mayday Hospital and operated upon the same day. A fibroid tumour the size of a tennis ball was found in the right side of the uterus, its upper limit being just below the round ligament. It was enucleated after making elliptical incisions above the edges. The cavity was obliterated by mattress sutures and the uterine incision closed with continuous catgut. Closure of the abdomen was again reinforced with three figure-of-8 silk-worm-gut tension sutures. There was again no attempt at abortion, and she was discharged twenty days later. She gave no anxiety during the remainder of her pregnancy and had a spontaneous delivery at term, the child weighing 7 lb. 13 oz. The uterus involuted normally, but she had a small thrombosis in a vein of the right leg. She was discharged at her own request on the sixteenth day, and when seen at the post-natal clinic a month later was found to be normal in every way.

Case III

A primipara aged 31, married nine years, was admitted to Mayday Hospital with a history of three and a half months' amenorrhoea and pain in the right side for two days, which had become acute on the day of admission. There was no vomiting or vaginal discharge. On examination a tender, slightly mobile swelling was found in the right iliac fossa, apparently separate from the uterus. She was kept under observation for the next twelve days, with opiates to relieve pain. There was, however, no improvement, and the swelling remained acutely tender. Laparotomy was therefore carried out, and on exposing the uterus a fibroid of about the same size as that found in Case II was seen. It had a short but broad pedicle, which had undergone partial torsion, there being about one and a half twists. An incision was made into the uterine wall enclosing the pedicle and the tumour resected. Some troublesome bleeding from the uterine walls occurred, and several vessels were separately tied. The uterine muscle was then brought together by mattress sutures, the peritoneal coat being closed with continuous No. 0 catgut. The abdomen was closed as in the previous cases.

The patient had severe pelvic pain for the whole of the second night and third day after operation, and was given morphine and corpus luteum extract. After this her convalescence was normal. She was readmitted to Mayday Hospital two weeks before term, and was delivered a week later, the child weighing 6 lb. 10 oz. On discharge the uterus was noted as slightly bulky, but there was no loss, and when she was seen at the post-natal clinic a month later involution was complete.

All three uterine tumours on being cut proved to be myomata, and each had small areas of haemorrhage in the centre.

Points of Interest

In all three cases the chief complaint was acute abdominal pain.

Case I, in addition to the uterine tumour, had bilateral ovarian tumours, which might have caused obstruction of labour. In one sense it was fortunate that the patient developed acute symptoms from the uterine tumour, as this occurrence led to the discovery and removal of the ovarian tumours, which might otherwise have been missed, giving trouble only after labour had started. In view of the extensive operative measures it is remarkable that this uterus remained quiescent.

Case II had a vaginal loss when first seen, and would probably have aborted if myomectomy had not been carried out.

Case III undoubtedly threatened to abort after operation. This is not surprising when it is remembered that torsion of the pedicle had begun, which must have rendered the uterus irritable, and that there was superadded the excision of the tumour. This patient was also the least advanced in pregnancy.

All three patients were operated upon in the second trimester. This is the period of pregnancy when the uterus is often sluggish, as shown by the occasional difficulty encountered in inducing therapeutic abortion. I have several times had to insert laminaria tents, after dilatation, rupture of the membranes, and cervical packing had failed to stimulate uterine action. In two cases I have had to empty the uterus by vaginal hysterotomy.

All three cases went on to term and had spontaneous deliveries of live babies, involution being satisfactory in each case.

A MANŒUVRE TO FACILITATE ABDOMINAL PALPATION

BY

CHARLES H. OSBORN, F.R.C.S., F.R.A.C.S.

*Surgeon to In-patients, St. Vincent's Hospital,
Melbourne, Australia*

Palpation holds pride of place among the routine methods of physical examination of the contents of the abdomen, owing to the fact that this cavity, unlike others in the body, is enclosed more by muscles than by bones, but it is obvious that the amount of information to be gleaned from the use of this method must necessarily depend upon the degree of muscular relaxation which can first be obtained. It is true that it is impossible to overcome that involuntary or reflex rigidity of the abdominal musculature which is associated with certain acute abdominal diseases and which is in itself a valuable indication of the presence of one of these conditions; but my purpose in this paper is to describe a method whereby it is possible to diminish the voluntary contraction of the abdominal musculature frequently observed in nervous and apprehensive patients. The method is also valuable in those patients in whom, owing to pronounced lumbar lordosis or to increased intra-abdominal tension, the abdominal wall is tense and stretched.

The customary routine, before abdominal palpation, is to attempt to obtain a maximum degree of relaxation by placing the patient in the supine position, with the arms at the sides and the head and neck comfortably elevated upon a pillow. The lower limbs are drawn up by semiflexing the hip- and knee-joints, and the