fore must be put in the most useful position at as early a date as possible. In most cases flexion to 75 degrees with the hand midway between pronation and supination is the optimum position. Formal excision of the joint is rarely called for, but may be necessary in severe infections to save the limb.

Fractures of the Forearm

One or both bones may be involved, muscles and tendons are often severely damaged, and nerve injury is not uncommon. The preservation of a useful limb therefore presents very considerable difficulty. For transport to the casualty clearing station immobilization on a straight splint, supporting the hand as well as the forearm, is sufficient. At the casualty clearing station, after the routine treatment of the wound, bone fragments, especially those driven into muscle, should be removed. It is better to leave a large gap between the bone ends and a wound that heals readily than to leave small fragments of bone, which inevitably necrose and delay healing for months. The gap can subsequently be filled by bone grafting, which was carried out successfully in many of these cases. Repair of divided nerves and tendons should be effected if possible, though this is not often the case.

For transport to the base a straight Thomas splint with a strapping extension, the forearm being in full supination, is the best method. At the base the straight Thomas splint may be used for the first fourteen days, after which time a bent Thomas splint may be applied. Plaster-of-Paris, though of value later, is not very suitable in the early stages owing to the difficulty in controlling the fractured bones without extension.

Fractures of the Wrist and Hand

Wounds of the carpus are often explosive in type, with a large exit wound and with great damage to tendons: the wrist-joint is commonly involved. In the forward area, after dressing, a simple flat splint is applied. At the casualty clearing station, when the wound has been excised, the carpal bones are carefully examined, and any that are badly comminuted are removed. Closure of the capsule of the wrist-joint is usually impossible, and the wound cavity is best packed with gauze smeared with petroleum jelly. Primary amputation should only be carried out where tendons, vessels, and nerves have been completely destroyed.

Splinting of wrist-joint injuries with extensive wounds is difficult, but for transport to the base a simple flat splint extending from the roots of the fingers to the elbow-joint is generally adequate. At the base, provided healing is proceeding satisfactorily, the hand and wrist should be put on a Jones cock-up splint or in plaster, the wrist being dorsiflexed. On the whole this type of wound gives poor results, but even if partial use of one finger and the thumb can be obtained this is better than any artificial limb.

At a joint meeting of the Yorkshire Regional Committee and the Sheffield Regional Committee of the British Hospitals Association, held in the General Infirmary at Leeds, approval was given to the creation of two divisions to cover Yorkshire (excluding the Cleveland area) and parts of Derbyshire, Lincolnshire, and Nottinghamshire: one division to be centred on Leeds and the other on Sheffield. Power will be given to the divisional authorities to split these two divisions into regions and subregions. The object is to co-ordinate hospital services and avoid duplication and overlapping.

THE ENTRANCE INTO MEDICINE*

BY

HUGH LETT, C.B.E.

President, Royal College of Surgeons of England

The medical curriculum to-day is very different from what it was forty-five years ago when I became a medical student. At that time *x* rays and radium had not been discovered, abdominal operations were few, and the surgery of the brain and thorax was rudimentary; nothing was known of vaccines, vitamins, or hormones, little of blood diseases, and the work in the clinical laboratories was confined to what we now regard as the simplest examinations. There were no ante-natal departments or special departments for diseases of children or infant welfare, for orthopaedic, for genito-urinary, or for chest surgery, to mention only some of those there are to-day.

Fundamental Principles

But although so much more is now known about every branch of medicine and surgery and there is so much more to learn, the fundamental principles of medical education are the same. Now as then there are things that must be known, some that ought to be known, and others with which the student should have some acquaintance if only to satisfy examiners. But that is not all, for in addition to acquiring a certain amount of medical knowledge the student is taught to cultivate accuracy, observation, and deduction, and to reason carefully, so that he or she may be able to bring a trained intelligence to bear upon medical problems.

Medical education should mean much more than the acquisition of so many facts and theories; it should train the mind in such a way that it may be ready to take the initiative, to adjust itself quickly to new situations, to apply to each case in a free and understanding way the principles, the routine, the discipline which have become second nature. The years spent as a medical student are only an introduction to a study which must continue throughout professional life.

The Pre-clinical Years

The curriculum is long and you may feel impatient at the time that must elapse before you reach the wards and come into contact with sick people. You may feel that the time given to chemistry, physics, and biology is wasted, and that a great part of what you are taught in anatomy and physiology is quite unnecessary; yet these subjects have a very important bearing on clinical work. A preliminary knowledge of them is essential if you are to understand the various methods of examining your patients, many tests that are carried out, and certain forms of treatment.

Let us consider for a moment some of the things we owe to physics. First of all the *x* rays, which have proved of such immense value in the diagnosis of disease in every part of the body, and radium, which like the *x* rays has done so much for inoperable cases of malignant disease. Physicists discovered the ultra-violet and infra-red rays, and, while both have their value in treatment, the former play an important part in the demonstration of ultra-microscopic viruses. I need hardly refer perhaps to the importance of a knowledge of electricity, but as one instance I may mention the recent discovery that waves of electricity pass over the brain, and that alterations in these waves tell us of the presence of disease or a tumour. At present it is possible to localize only those lesions that are on the surface, but there can be no doubt that before long the method of electro-encephalography will enable us to determine the exact site of a tumour lying deeply in the

* An address (abridged) given at the London (Royal Free Hospital) School of Medicine for Women, University of London, on June 1, 1939.

brain substance. Lastly, we must not overlook the remarkable and ingenious developments of certain optical instruments by which we can now see the interior parts of the body.

Biochemistry has taught us about vitamins, the vital importance of the secretion of the ductless glands, and has given us many remarkable drugs which have revolutionized the treatment of a number of diseases at home and in the Tropics, so that now with the help of these drugs we expect a recovery in many cases which formerly would have been regarded as hopeless.

Lister by his discovery of the principles of antiseptics laid the foundations of modern surgery, and it is impossible to over-estimate the importance of his work; but the extraordinary developments of the last forty years could never have taken place had it not been for our physicists and biochemists who made these developments possible. Much of the valuable experimental research that is being done to-day could never have been attempted had it not been for the advances that have been made in physics and biochemistry. And so, if we recognize how much medicine and surgery owe to them, it is not difficult to understand why physics, chemistry, and biology form the first part of the education of the medical student.

Anatomy and physiology teach us the structure of the normal human body, the functions of its various organs, and the part they play in the life of the healthy individual. It is obvious that we must have a clear understanding of the body in health before we attempt to recognize the changes that take place when it is diseased, and before we can understand why these changes give rise to the various symptoms and signs that show illness and disordered function.

An important advance in medical education was made recently when it was decided that while the student is still working at anatomy and physiology he should be taught something of the elementary principles of medicine and surgery and learn certain methods of examination which previously were taught only in the clinical years. This elementary clinical instruction helps you to appreciate more clearly the importance of anatomy and physiology, for the gap that previously existed between them on the one hand and medicine and surgery on the other is now bridged, and they are no longer treated as independent parts of the curriculum.

The Clinical Years

When you reach the wards and begin clerking and dressing you will find that among other duties you have to take histories of the patients allotted to you. A good history includes an account of the patient's illness and his previous health, his present symptoms, and finally a description of what you have been able to discover after careful examination, not only of the part that is principally affected but of the whole body.

Now you may feel that all this note-taking is very troublesome and that it is unnecessary. I remember one student who said that he really did not see why he should spend his time taking notes for the benefit of the hospital. I may say, perhaps, that from my recollection of his notes any benefit that the hospital could have derived from them was remarkably small! But he was approaching his work in the wards from the wrong angle. It did not occur to him that the taking of histories was an important part of his education. Careful, systematic note-taking and a routine examination of the whole body in every case will teach you method, develop your powers of observation, and impress the details of cases on your memory. The importance of examining the whole body is sometimes overlooked, and the omission has been followed by serious results.

Try to make your own diagnosis in every case. At first you will find it very difficult and often impossible,

but as your knowledge increases you will be able to do so more and more frequently and with greater success. Do not be too much afraid of making mistakes. We all make them; but if we make our mistakes in spite of having exercised care and thought, and if when we have made them we try to find out where our reasoning or observation was at fault, we may learn more from our mistakes than from our easy or sometimes fortunate successes.

Try so far as possible to make a preliminary diagnosis before the results of the x-ray examinations and the clinical laboratory tests are available. In many obscure cases such a diagnosis will not be possible, nevertheless it is a sound practice to try to make it. Some day you may be in such a position that these examinations cannot be made, and you may have to make a diagnosis when you and your patient are far from an x-ray apparatus or a clinical laboratory. But an even more important reason for following this method is that by doing so one acquires keener powers of observation, reasoning, and deduction. The rapid progress made in medicine and surgery has brought a certain danger in its train: the possibility that too much reliance may be placed on the result of tests in the laboratory and examinations in the x-ray department at the expense of careful clinical examination. This can never be replaced.

A full and careful study of the patient, body and mind, is the essential foundation on which our healing art is built. The training of our students must be directed above all to the cultivation of careful clinical observation and an understanding and appreciation of signs and symptoms. In this way only can they acquire that vital yet intangible something we call clinical instinct. Clinical instinct, born of observation, nourished by experience, and educated by deduction, must and always will remain one of the greatest and most important gifts a medical man can have.

The Approach to the Patient

It is perhaps remarkable that in the past the medical student should have received so few hints on the attitude to be adopted towards patients, the way to approach them, to manage them, to obtain their confidence. On entering practice he has had to rely on his natural gifts and instinct, or perhaps, if he has been fortunate, upon his ability to copy an admired clinical teacher. And yet the importance of understanding the patient and appreciating his mental condition and outlook has long been recognized, even by laymen. Montaigne writes:

"The Physition hath neede of many parts, divers considerations and severall circumstances to proportion his desseigne justly. He ought to know the sick man's complexion, his temper, his humour, his inclinations, his actions, his thoughts and his imaginations."

The introduction into the curriculum of psychology and some knowledge of the psychoneuroses and of the important part the mind plays in good and bad health should be a very material help to both doctor and patient. But the physician who in the approach to his patient cultivates sympathetic understanding, cheerfulness, and confidence may be said to have acquired already the fundamentals of bedside or consulting-room psychology.

Sentimentality must of course be avoided, but if you can develop a sympathetic understanding of your patient's anxieties and difficulties you will find that you can help him greatly, not only in his illness but in his convalescence. You will be able to do much to reassure him, to remove great and little discomforts, both physical and mental, and so make his path easier.

A quiet cheerfulness is also a great asset. The gloomy physician "maketh the sick man sicker." Sir Benjamin Brodie said that it was a good rule in the practice of an art, as in the common affairs of life, to look on the favourable side of the question so far as it was consistent with reason to do so. But the cheerful attitude

does not mean that aggressive, bouncing cheerfulness which can be so very irritating, especially to the sick man. "A Physician's first entertainment of his patient should be gracious cheerful and pleasing." If you should be anxious about your patient try to conceal your anxiety from him, and remember that while you are making your examination and talking to him he will be searching your expression to learn whether you really believe in the assurance you are trying to give him.

As for confidence, this can only come with experience and hard work, by learning from what you have seen and read, and by thinking over the reasons for your successes as well as your mistakes. Nothing is more ugly than conceit or vanity, but they are very different things from that confidence in which, said the prophet of old, shall be your strength. Those who are naturally nervous may find this confidence extremely difficult to acquire, but I think they would be much helped by reading an address which Dame Louisa Aldrich-Blake gave to this School nearly forty years ago. Here is one extract:

"From the very beginning of one's career one must school oneself to look upon nervousness as a disability one must do one's utmost to conquer. For those whose temperament makes them need this discipline most the process is a very unpleasant one. I suppose the first step is to look it in the face as an enemy."

SCIENTIFIC TREATMENT OF DELINQUENCY

The seventh annual general meeting of the Institute for the Scientific Treatment of Delinquency was held at its new house, 8, Portman Street, W., on May 22, Dr. E. T. JENSEN presiding. The annual report stated that the year had been noteworthy for two important developments in the Institute's educational work. One of these was a week-end course for doctors, and the other the recognition of the Institute by the University of London as a centre for extension lectures. The subjects of research which have occupied the year have included the treatment of sex offenders, general investigations on intelligence in relation to crime, the aetiology of exhibitionism, and temperament tests. In the psychopathic clinic 166 new patients were treated during the year, of which number 106 were drawn from the courts and twenty-four from medical practitioners and clinics. The number of cases described as psychoneurotic was fifty-three, and the number of character cases (including psychopathic personalities and sex perverts apart from neurosis) was fifty-six. Theft was the chief type of offence represented (seventy-six), sex offences coming next (forty-six). The Institute applauded the Criminal Justice Bill as a humane and progressive measure. "We believe that the work of the Institute must have played an important part in demonstrating the feasibility of and need for such [scientific] treatment, and we are grateful indeed to Sir Samuel Hoare for what is, in effect, our charter."

Young Offenders

Dr. DENIS CARROLL, co-director of the Institute, gave an address. The medical attitude towards the treatment of delinquents, he said, had changed. Formerly the doctor was concerned mainly with the question of ascertaining the criminal's responsibility for his actions, but now doctors had discovered that it was possible to treat delinquents by methods originally intended for the treatment of mental disorders. The most important group concerned consisted of young offenders. There were two peak periods associated with delinquency, one of them round about 14, produced by the stress of leaving school added to the emotional stresses incident on puberty, and the other at about 16 to 17, associated with the well-known difficulties of adolescence. One of the hardest

problems to come up against was the juvenile who had formed the habit of crime. He had seen children of 8 or 9 who had been delinquent from their earliest years. It was not a question of heredity. He did not believe there was such a person as the born criminal, at any rate not in the sense that the child's instincts were more criminal than those of older people, but no doubt strength of instinct might be inherited, and this was capable of being turned to good or bad use.

Among juvenile delinquents some 10 per cent. had gross mental defect and 50 per cent. were backward. Roughly speaking, there were three or four times as large a proportion of backward children among delinquents as among the ordinary child population. At one time, under the influence of the Italian school, it was thought there was such a thing as a physical criminal type. So far was this from being true that no physical factor of any kind could be found which might be regarded as specific of delinquency. In a very small group it had been discovered that on the whole, but not invariably, the asthenic type of physique was shown, but that amounted to no more than an interesting fact without significance. The offences which children who had suffered from chorea or sleepy sickness committed were not different from those associated with violent temper. It was not uncommon to find that a child who, for example, had had a fractured leg, with permanent shortening, by reason of his lack of capacity came to have a feeling of inferiority which might stimulate him to delinquent courses in order to "get even." But not many delinquents committed an offence merely because they wanted a certain glamour; when such children came to be studied their condition was much more complicated. A good deal of nonsense was talked about glandular conditions, but so far as was known there was no specific relationship between glandular disorders and crime. Some failures of development of the sex glands were associated with personal changes, and some pituitary abnormalities with gross changes in appearance, and there again it might be that a feeling of inferiority was at work.

Dr. Carroll went on to speak of nervous conditions, hysteria, and certain obsessional states—the kind of thing which always made a person want to go back to see if he had turned the taps off. A huge field was covered by these neurotic symptoms or, in a mild form, neurotic character traits. In such conditions offences might be related to the disease in the sense of being a symptom. A man with paranoic tendencies might believe that his neighbour was about to stab him, and anticipate the event by stabbing him in self-defence. If the neurotic state could be cleared up such persons would go back to the average level of capability to resist delinquent impulses.

It was common to speak of poverty as a cause of crime. Poverty was often associated with crime, but to describe it as a cause was a different matter. The relation was the same as that of bad housing. In such conditions the home was likely to be badly run, and in the sensitive child some form of delinquent behaviour might be produced. Unsuitable work and undesirable associates might stimulate delinquent response in certain subjects.

In conclusion Dr. Carroll described the elaborate procedure through which cases were put, the mental and physical examinations, the intelligence tests and the investigation of the social history. After the diagnosis had been made a decision was taken as to what should be done and how much supervision was necessary, and a report was made to the court or other body which had referred the case. The Institute arranged for any necessary treatment, medical or psychological, to be carried out. Re-education of the social sense depended on getting a thoroughly good rapport, for which the delinquents had not very much capacity, so that these methods were limited in their application. Where a persecutory element was to the fore the best results were got by analytical methods. The subject was encouraged to talk freely. Some of the best results had followed from this so-called superficial analytical treatment, always providing that there was no