Pregnanediol excretion was found in all cases having a secretory endometrium.

On the basis of Venning and Browne's findings estimates were made as to the normality of ovarian function.

The expenses of this investigation were partly defrayed by a grant (to A. M. H.) from the Moray Fund.

BIBLIOGRAPHY

- Browne, J. S. L., Henry, J. S., and Venning, E. H. (1937). *J. clin. Invest.*, **16**, 678.
 Venning, E. H. (1938). *J. biol. Chem.*, **126**, 595.
 — and Browne, J. S. L. (1936). *Proc. Soc. exp. Biol.*, N.Y., **34**, 792.
 — (1937a). *Amer. J. Physiol.*, **119**, 417.
 — (1937b). *Endocrinology*, **21**, 711.
 — Henry, J. S., and Browne, J. S. L. (1937). *Canad. med. Ass. J.*, **36**, 83.

TREATMENT OF GONORRHOEA BY M & B 693

BY

JAMES SOMMERVILLE, M.B., Ch.B.

Medical Officer, Bellahouston Venereal Diseases Centre,
Glasgow

In May, 1938, I began to treat a series of cases of gonorrhoea with the preparation which is now known as M & B 693. In this series 152 consecutive cases come under review; from these are excluded thirteen early defaulters and irregular attenders. Failures—cases which showed no material improvement after approximately two to three weeks' administration of the drug—numbered five. It should be noted here, however, that those "failures" cleared rapidly when irrigation was instituted after the drug had been withdrawn. Relapses occurred in eleven cases: six under fourteen days and five between twenty-one and forty-two days, all in alcoholics. Re-exposures numbered five, and the defaulter rate was about 30 per cent.

The following represents the detailed results in 134 cases. I began these investigations by using a dosage of two tablets three times a day. I excluded adjuvant irrigation therapy, as I had had a long-standing suspicion that the high incidence of complications which occurred during treatment at all the Glasgow centres—somewhere in the region of between 30 and 40 per cent.—was due to the act of irrigation, and also because I wished to see what the drug could do unaided. I began, therefore, with a dosage of two tablets thrice daily, later increasing it to two tablets four times daily to obtain a more even concentration during waking hours. Precautions with regard to sulphur intake were emphasized. As the results with two tablets four times a day were more encouraging I was emboldened to try three tablets four times a day; but my first case produced a rash with temperature, and I did not feel justified in proceeding further with this dosage.

Later, to find if a smaller dosage given in the same way would have as much benefit and, I hoped, with lessened toxic effects, I instituted a daily dosage of one tablet four times a day, and until recently this has been my routine treatment. The tablets were given until the discharge had completely ceased, and maintained for a period varying from one to two weeks thereafter. Following this there was a rest from treatment for three or four days, followed by a week's treatment, and then

two weeks' rest; after which a prostatic smear was taken, and, if this was satisfactory, a week later a urethoscopic examination was carried out and the urethra dilated, with a subsequent examination and blood Kahn test. Throughout the course of treatment the patients were so far as possible seen daily, but this quite often proved impossible, so that "dry" readings were more numerous than "G.C.-negative" ones. Thus it may safely be assumed that the figures obtained represent periods longer than would be got by exact findings under "indoor" supervision.

Table I shows the results obtained with varying dosage, and the extraordinary fact emerges that one tablet four times a day produced my best results, surpassing those with double that amount.

TABLE I

Dosage	Urethral Smear G.C.-negative		Urethra Dry	
	No. of Cases	Average in Days	No. of Cases	Average in Days
2 tablets 3 times daily ..	9	3.33	22	5.73
2 " 4 " " " ..	12	3.00	17	3.80
1 tablet 4 " " " ..	40	1.92	33	3.37
3 tablets 4 " " " ..	1	2.00		
Total ..	62	2.33	72	4.19

Thus it will be seen that sixty-two cases were gonococcus-negative in an average of 2.33 days, and if we read "dry" as "G.C.-negative" we get an average of 3.33 days for the total 134 cases.

As cases may be gonococcus-negative but not dry I have extracted my total "dry" observations.

105 cases were dry in 7 days or under, averaging 4.21
 16 " " " 14 " " " 10.7
 5 " " " 21 " " " 16.9
 1 case was dry in 26 days.

Taking detailed dosage the following results were got:

29 cases with 2 tablets thrice daily were dry in average 7.2 days
 29 " " " four times daily, " " " 5.62
 69 " " 1 tablet " " " " " 4.86

The remaining seven cases were gonococcus-negative but not dry before defaulting.

Complications.—No complications were encountered during treatment, and where complications were noted before treatment these cleared up rapidly.

Intolerance.—This was observed in fifty-two cases, or approximately 39 per cent. Table II reveals the frequency of each toxic symptom noted and its average time incidence in days. In some cases more than one symptom occurred.

TABLE II.—Intolerance

Symptom	No. of Cases	Average Day Incidence
Vomiting ..	3	3.5
Nausea ..	30	3.7
Headache ..	26	4.0
Vertigo ..	6	4.3
Migraine ..	1	5.0
Temperature ..	5	6.2
Rash ..	3	6.6
Constipation ..	2	7.0

No case of breathlessness or cyanosis was found. With the dosage of one tablet four times a day the symptoms

of headache and nausea only were observed. All the cases of rash and temperature occurred with the larger doses.

In view of the suggestions and observations with regard to the delayed method of treatment, I have extracted my findings on seventy-three cases with the same dosage—namely, one tablet four times a day. I have followed the method adopted by Cokkinis and McElligott, and although my numbers are small they at least give some guidance. From Table III it is evident that there is no material difference between early and late treatment.

TABLE III

Duration of Disease before Treatment (Days)	No. of Cases	G.C.-neg. (Days)	Dry (Days)
1-3	21	1.9	4.65
4-5	4	3.25	6.5
6-7	17	1.63	5.41
8-10	6	3	5.16
11-14	8	2.8	6.5
14-21	11	2	4.3
Over 21	6	2.33	2.62

From those cases in which it was possible to obtain an accurate estimation of the incubation period I have extracted Table IV; it would seem that the duration of this period has apparently no effect on the action of the drug.

TABLE IV

Incubation Period in Days	No. of Cases	G.C.-neg. (Days)	Dry (Days)
1-3	24	1.63	4.87
4-5	13	2	3.9
6-7	18	2.54	5.22
8-10	3	3	6.3
Over 10	5	2.5	5.4

Conclusions

M & B 693 is an exceedingly potent remedy for gonorrhoea, if not the most potent yet obtained, giving in this series 90 per cent. apparent early cures.

It is equally effective in early and late cases, and therefore delay in starting treatment, with its potential subsequent onset of complications, can be avoided.

Moderate dosage—namely, one tablet four times a day—would seem to give better results than greater doses. Intolerance under these circumstances, therefore, is almost negligible.

The high incidence of complications obtaining during treatment under the old irrigation therapy has been completely abolished.

Summary

152 cases of gonorrhoea under treatment with M & B 693 are reviewed. Thirteen of these patients defaulted or attended irregularly, and five failed to respond to the drug.

Of the remaining 134 cases eleven relapsed, due to alcohol in all instances.

Early cure was obtained in 90 per cent. of cases. This cure was possible in, on an average, under five days.

Apparent failures cleared up quickly with subsequent irrigation.

Toxic effects occurred in 39 per cent. of cases, and these were generally slight.

No complications during treatment were encountered.

Additional Note

Since these cases were reviewed I have altered my technique in an effort to overcome the incidence of failures, and still maintain freedom from complications. New cases have been given M & B 693 tablets with a daily dosage of one tablet four times a day for one week only, followed by daily irrigation with 1 in 8,000 potassium permanganate for one week, and then lavage on alternate days for another week. So far, in fully fifty cases this method has met with complete success, there being no failures and no complications during treatment. I hope at some future date to review these results fully.

My thanks are due to Messrs. May and Baker for original supplies of the drug, and especially to Dr. R. Forgan for suggestions as to dosage in the early part of my investigations. I am also indebted to my colleagues Drs. A. D. MacLachlan and Albert Sharman for their help and ready compliance with my suggestions, and to the clinic attendants for the increased work which they undertook with enthusiasm.

VACCINIAL ENCEPHALITIS IN A CHILD AGED 14 WEEKS

BY

E. SAKOSCHANSKY, L.R.C.P., M.R.C.S.

AND

H. J. TRENCHARD, M.B., Ch.B.

*House-physicians, Royal South Hants and Southampton
Hospital*

The points of special interest in this case seem to be the mildness of the disease, its onset on the tenth day after vaccination, its simulation (before admission to hospital) of many common childish ailments (except for torticollis), and the otherwise apparently normal progress after vaccination.

Vaccinial encephalitis is generally considered a complication of vaccination in older children and adolescents. Hewlett (1933) states that the young infant seems to be wholly immune from this complication. J. F. M. Scott (1932), who has collected twenty-two cases in the first year of life, illustrates the rarity of vaccinial encephalitis at this age by the following figures. Out of a total of 569 cases collected from different parts of Europe only forty-one (7.2 per cent.) occurred under 1 year. Gunnar Olin (1938) states that little or no risk accompanies vaccination under the age of 1 year: he computes the incidence of post-vaccinial encephalitis in Sweden from 1924 to 1936 as 1 in 30,246 of infants primarily vaccinated during the first year of life.

Case Report

The patient was a female, aged 14 weeks on admission, the first child of healthy parents. It had been breast-fed from birth, and the only previous illness was a heavy cold lasting a few days when 3 weeks old. There was no family history of tuberculosis. The father had never been vaccinated, and the mother was vaccinated as a child. The following is a chronological account of the illness.

On February 25, 1939, the child, then aged 13 weeks, was vaccinated with calf lymph—two marks on the left arm. This resulted in normal vesicles on March 2, and normal pustules