of headache and nausea only were observed. All the cases of rash and temperature occurred with the larger doses.

In view of the suggestions and observations with regard to the delayed method of treatment, I have extracted my findings on seventy-three cases with the same dosage—namely, one tablet four times a day. I have followed the method adopted by Cokkinis and McElligott, and although my numbers are small they at least give some guidance. From Table III it is evident that there is no material difference between early and late treatment.

TABLE III

Duration of Disease before Treatment (Days)	No. of Cases	G.C.-neg. (Days)	Dry (Days)
1-3	21	1.9	4.65
4-5	4	3.25	6.5
6-7	17	1.63	5.41
8-10	6	3	5.16
11-14	8	2.8	6.5
14-21	11	2	4.3
Over 21	6	2.33	2.62

From those cases in which it was possible to obtain an accurate estimation of the incubation period I have extracted Table IV; it would seem that the duration of this period has apparently no effect on the action of the drug.

TABLE IV

Incubation Period in Days	No. of Cases	G.C.-neg. (Days)	Dry (Days)
1-3	24	1.63	4.87
4-5	13	2	3.9
6-7	18	2.54	5.22
8-10	3	3	6.3
Over 10	5	2.5	5.4

Conclusions

M & B 693 is an exceedingly potent remedy for gonorrhoea, if not the most potent yet obtained, giving in this series 90 per cent. apparent early cures.

It is equally effective in early and late cases, and therefore delay in starting treatment, with its potential subsequent onset of complications, can be avoided.

Moderate dosage—namely, one tablet four times a day—would seem to give better results than greater doses. Intolerance under these circumstances, therefore, is almost negligible.

The high incidence of complications obtaining during treatment under the old irrigation therapy has been completely abolished.

Summary

152 cases of gonorrhoea under treatment with M & B 693 are reviewed. Thirteen of these patients defaulted or attended irregularly, and five failed to respond to the drug.

Of the remaining 134 cases eleven relapsed, due to alcohol in all instances.

Early cure was obtained in 90 per cent. of cases. This cure was possible in, on an average, under five days.

Apparent failures cleared up quickly with subsequent irrigation.

Toxic effects occurred in 39 per cent. of cases, and these were generally slight.

No complications during treatment were encountered.

Additional Note

Since these cases were reviewed I have altered my technique in an effort to overcome the incidence of failures, and still maintain freedom from complications. New cases have been given M & B 693 tablets with a daily dosage of one tablet four times a day for one week only, followed by daily irrigation with 1 in 8,000 potassium permanganate for one week, and then lavage on alternate days for another week. So far, in fully fifty cases this method has met with complete success, there being no failures and no complications during treatment. I hope at some future date to review these results fully.

My thanks are due to Messrs. May and Baker for original supplies of the drug, and especially to Dr. R. Forgan for suggestions as to dosage in the early part of my investigations. I am also indebted to my colleagues Drs. A. D. MacLachlan and Albert Sharman for their help and ready compliance with my suggestions, and to the clinic attendants for the increased work which they undertook with enthusiasm.

VACCINIAL ENCEPHALITIS IN A CHILD AGED 14 WEEKS

BY

E. SAKOSCHANSKY, L.R.C.P., M.R.C.S.

AND

H. J. TRENCHARD, M.B., Ch.B.

*House-physicians, Royal South Hants and Southampton
Hospital*

The points of special interest in this case seem to be the mildness of the disease, its onset on the tenth day after vaccination, its simulation (before admission to hospital) of many common childish ailments (except for torticollis), and the otherwise apparently normal progress after vaccination.

Vaccinial encephalitis is generally considered a complication of vaccination in older children and adolescents. Hewlett (1933) states that the young infant seems to be wholly immune from this complication. J. F. M. Scott (1932), who has collected twenty-two cases in the first year of life, illustrates the rarity of vaccinial encephalitis at this age by the following figures. Out of a total of 569 cases collected from different parts of Europe only forty-one (7.2 per cent.) occurred under 1 year. Gunnar Olin (1938) states that little or no risk accompanies vaccination under the age of 1 year: he computes the incidence of post-vaccinial encephalitis in Sweden from 1924 to 1936 as 1 in 30,246 of infants primarily vaccinated during the first year of life.

Case Report

The patient was a female, aged 14 weeks on admission, the first child of healthy parents. It had been breast-fed from birth, and the only previous illness was a heavy cold lasting a few days when 3 weeks old. There was no family history of tuberculosis. The father had never been vaccinated, and the mother was vaccinated as a child. The following is a chronological account of the illness.

On February 25, 1939, the child, then aged 13 weeks, was vaccinated with calf lymph—two marks on the left arm. This resulted in normal vesicles on March 2, and normal pustules

on March 6. On the 7th she vomited in the evening. Castor oil was given, and she slept till morning. The bowels acted in the morning of the 8th. The mother stated that the child had a peculiar cry, and held her head to the left side. The child slept all day. On examination in the evening she appeared drowsy, resented being disturbed or placed on the right side, and did not recognize her mother or smile. She fed well, however. The head was held turned to the left, and she seemed to resent it being straightened. There was no rise of temperature, and no external cause was found for the torticollis. No enlarged glands were seen, the tympanic membranes were normal, and there was no strabismus, nystagmus, neck rigidity, or Kernig's sign.

On March 9 the child was admitted to hospital. It was conscious, but seemed very quiet. There were two black scars on the left arm, due to vaccination; these were not surrounded by any inflammatory zone, and no enlarged lymphatic glands were palpable in any part of the body. The fundi were normal and the pupils reacted normally to light. The head and eyes were directed to the left, and there was a weakness of the left lower face, causing a drooping of the corner of the mouth. The movements of the limbs appeared normal. The knee-jerks were present and equal, but the ankle-jerks were not obtained. The plantar reflexes were extensor and equal on the two sides. There was a slightly positive Kernig's sign and a suspicion of neck rigidity. The tympanic membranes were normal, and nothing abnormal was discovered in the throat, heart, lungs, or abdomen. Lumbar puncture was performed, and showed a clear fluid under moderate pressure. The pathological report on the fluid was: cells (lymphocytes), 60 per c.mm.; globulin, slight excess; protein, 0.06 per cent.; chlorides, 710 mg. per 100 c.cm.; sugar, low (the fluid had been kept overnight before examination); culture, no growth; Wassermann reaction, negative; Lange test, 0012321000.

On March 10 a temperature of 99.5° was recorded, but the physical signs remained unchanged. Treatment consisted of 10 c.cm. whole blood intramuscularly from the mother and 10 c.cm. whole blood from an adult vaccinated seven years previously. On the 11th, in addition to the signs previously noted, there was a right sixth cranial nerve paralysis, and on upward movement of the eyes a coarse lateral nystagmus occurred. The movements of the left upper limb seemed weak, the muscles especially affected being the deltoid, flexors of the arm, and extensors of the wrist. The arm was held motionless. Five c.cm. of serum was given intramuscularly (serum from an adult vaccinated four weeks previously from the same batch of lymph). The next day the signs were unchanged, and 7.5 c.cm. of serum was given. On the 13th the general condition seemed better. The sixth cranial nerve palsy was not so marked and movement of the arm had improved. Five c.cm. of serum was given.

Weakness of the left face had disappeared on March 14, but some weakness of the right side of the face seemed to be developing. Movement of the fingers of the left hand was present, but the arm was held immobile by the side unless stimulated. Kernig's sign was positive, and some neck rigidity was present. Five c.cm. of serum was given. By March 16 the squint and paresis of the left arm had recovered. There were no focal signs, but the neck rigidity and Kernig's sign were still present.

On March 22 the temperature, which had been running between 98° and 99°, rose to 100.6°. The general condition seemed good, but the neck rigidity and Kernig's sign, which had nearly disappeared, returned. Lumbar puncture showed a clear fluid under moderate pressure. It contained fifty-four cells per c.mm. and 0.04 per cent. of protein, but no excess of globulin. No growth was found on culture and there were no tubercle bacilli. On the 28th no neck rigidity was present, but Kernig's sign was still slightly positive. The child had begun to cry and to behave more normally. Lumbar puncture showed fluid under normal pressure. Examination showed: cells (lymphocytes), 6 per c.mm.; protein, 0.02 per cent.; globulin, no excess; chlorides, 710 mg. per 100 c.cm.; deposit, no tubercle bacilli; culture, no growth; Lange test, no reduction.

On April 1 no abnormal physical signs were observed, and the patient was discharged from hospital.

The serum used was obtained from a youth aged 17, who had been vaccinated four weeks previously with lymph from the same batch as that employed in vaccinating the patient.

We wish to express our thanks to Dr. Jeffreys, under whom the patient was in hospital, for permission to publish this case; to the medical officer of health and the Public Health Department of Southampton for tracing with such dispatch an adult vaccinated with lymph from the same batch as that used for the patient; to Dr. Gleave, pathologist to the Royal South Hants and Southampton Hospital, for the pathological reports and for the preparation of the serum; and to the donor who so readily gave his blood for the serum.

REFERENCES

- Hewlett, R. T. (1933). *Recent Progress in Medicine and Surgery*, edited by Sir J. Collie, p. 145, H. K. Lewis, London.
Olin, G. (1938). *Bull. mens. Off. int. Hyg. publ.*, **30**, 743 (quoted in *J. clin. Res.*, 1938, **25**, 32).
Scott, J. F. M. (1932). *Med. Annu.*, p. 569.

Clinical Memoranda

A Case of Injury to the Pancreas

An intra-abdominal injury affecting the pancreas alone is a rare occurrence, and seems to warrant the publication of this case.

CASE HISTORY

A carter aged 48 was standing behind his van when a lorry reversed into him, crushing him against the tailboard of the van. When released he collapsed, and was immediately taken to hospital. On admission he was pale, sweating, and severely shocked, with an almost imperceptible pulse. He complained of abdominal pain and was tender on deep palpation in the epigastrium and right flank. His condition improved rapidly, and six hours later his colour was good, his pulse was 78 and of good volume, and he felt much better. He still complained of some upper abdominal pain and was tender in the epigastrium and right hypochondrium, but there was no marked abdominal rigidity and no dullness in either flank. For the next twenty-four hours there was little change in his condition. After this time, however, his pain became more severe, he vomited several times, and his pulse rate began to rise steadily. Six hours later, just thirty-six hours after admission, his pulse had risen to 120, he had developed generalized abdominal rigidity with tenderness most pronounced in the upper abdomen, and he looked drawn and anxious. Laparotomy was decided upon, the diagnosis being ? rupture of small intestine in the region of the duodeno-jejunal junction.

Under spinal anaesthesia his abdomen was opened through an upper right paramedian incision. The peritoneal cavity was found full of blood-stained fluid, with areas of fat necrosis scattered everywhere. Some of these areas in the great omentum were an inch in diameter. The peritoneum over the pancreas was oedematous and the pancreas itself appeared dark and swollen, but no actual tear in the peritoneum could be found. No evidence of trauma to the intestines, stomach, spleen, or liver was seen. Two drainage tubes down to the pancreas—an upper between the stomach and transverse colon and through the lesser sac, and a lower below the transverse colon—were inserted in the wound and a third placed in the pelvis through a suprapubic stab wound. The incision was closed throughout with interrupted sutures, with tension sutures of thick silk through the rectus sheath, to minimize the danger of subsequent bursting of the wound. The patient's condition remained good throughout the operation. When the abdomen