TREATMENT OF CARCINOMA IN THE PHARYNX AND LARYNX AND ITS RESULTS*

BY

LIONEL COLLEDGE, F.R.C.S.

Surgeon, Ear and Throat Department, St. George's Hospital

The introduction of irradiation has rendered the problem of treatment in various situations infinitely more complicated, because the results of its use are neither so good that the method can claim to supplant simple surgical treatment altogether nor so bad that they can be dismissed as negligible. In certain situations other than the pharynx and larynx the matter is much simpler; for example, in the floor of the mouth irradiation has taken the place of surgery almost entirely, whilst in the stomach the abdominal surgeon does not need to concern himself with it. Unfortunately, it is also true that in many situations, such as the oesophagus, where the results of surgery are bad, those of irradiation also are poor, and, conversely, many of the best results obtained by irradiation are in situations where the tumour is most amenable to surgery. This applies to the larynx and pharynx. If surgical treatment is chosen, then an operation suited to the site and extent of the tumour must be designed; if irradiation is chosen, then the type of radiation and the method of its application must be settled.

It is proposed to deal first with tumours in the oropharynx—that is, tumours of the tonsil and fauces—and then to deal rather more fully with the lower pharynx—that is, the laryngopharynx and the larynx itself. My own endeavour will be chiefly to indicate what can be expected from surgery, and I will leave it to others to show the best that can be obtained by irradiation. A great variety of tumours may be encountered in the situations under consideration, but I think that this discussion will have more value if it is limited to epithelioma, which is the common form of malignant tumour.

In the oropharynx, epithelioma usually begins either on the lower part of the anterior pillar or in the angle between the lower pole of the tonsil and the side of the tongue, and is usually seen spreading over the tonsil and fauces and into the side of the tongue. There may or may not be a palpable gland in the neck; often there is not, but if the cervical glands are enlarged the size is often in inverse proportion to that of the primary growth.

Where the growth is comparatively early without fixation and there are no palpably enlarged glands in the neck, the tumour may be removed by diathermy, as Norman Patterson has long advocated. Such cases must be kept under observation lest enlarged glands appear in the neck. If a gland dissection has to be combined with removal of the primary tumour it should precede the diathermy operation by ten or fourteen days to avoid difficulties from the changes which diathermy produces in the neck. Patterson records that out of fifty patients treated by diathermy nine died and thirteen survived for more than three years.

In more advanced infiltrating growths, where the tongue is more invaded or the growth has become adherent to the inner surface of the mandible, diathermy is not

applicable alone, and the operation required includes removal of the mandible from the canine fossa to the condyle. This is naturally a mutilating operation and must be followed by a careful reconstruction. Dickie records that in thirty cases so treated there were seven deaths and fifteen recurrences in less than five years, but eight patients survived for more than five years and one for over eleven years. Radiation employed in such cases may take the form either of radium applied at a distance or of the implantation of radon seeds.

The treatment by irradiation is most attractive, because when successful the pharynx is almost normal in appearance, but the question remains whether in treating epithelioma it gives a sufficiently high proportion of lasting good results to displace surgery. Biopsy must take an important place in making this decision, because sarcoma and Group IV epithelioma, which in its clinical course may be said to have more affinity to sarcoma than to keratinized epithelioma, give uniformly bad results with surgery and are radiosensitive.

In the lower pharynx the growths to be considered are those on the lateral wall of the pharynx, the epiglottis, the aryepiglottic folds, the fossa pyriformis, and the post-cricoid area. Of these various types the growths on the lateral wall of the pharynx, the epiglottis, the aryepiglottic fold, and in the post-cricoid area may be treated by lateral pharyngotomy combined with a subsequent plastic operation if in an early stage. For these tumours in a more advanced stage and for tumours of the fossa pyriformis almost always operation must be a pharyngolaryngectomy. That operation includes clearance of the glands on at least one side of the neck, and removal of the whole larynx and either a portion or a complete segment of the pharynx. A somewhat elaborate plastic operation is required later to reconstruct the pharynx, the patient being left with a laryngectomy.

The classification of cases in these groups into those suitable for lateral pharyngotomy or for pharyngolaryngectomy and those which are inoperable must be decided by clinical experience, which takes into account the extent of the primary growth, the invasion and degree of fixation of lymphatic glands, and the general condition of the patient.

Gluck and Soerensen record fifty-eight operations by subhyoid pharyngotomy, but this method is applicable only to carcinoma of the epiglottis. In 334 cases of extrinsic cancer treated by laryngectomy combined with resection of the pharynx with seventy-seven deaths, they obtained a permanently good result in 10 per cent. In Gluck's clinic, patients whenever possible were submitted to operation, so that no attempt was made to make the statistics appear favourable by selection of cases. Of my own patients treated by pharyngotomy or pharyngolaryngectomy between 1920 and the beginning of 1935 the following are some particulars:

TABLE I

	No.	Died	Recurrence	Remain Well
Pharyngotomy :				
Private patients	13	0	7	6
Hospital "	26	9	10	7
Total	39	9	17	13
Pharyngolaryngectomy :				
Private patients	7	1	4	2
Hospital "	9	2	3	4
Total	16	3	7	6

* Read in opening a discussion in the Section of Oto-rhinolaryngology at the Annual Meeting of the British Medical Association, Plymouth, 1938.

Of these cases five of pharyngotomy and two of pharyngolaryngectomy have remained well for more than eight years. That is, out of fifty-five operations nineteen were successful, or 29 per cent. Against this must be taken into account that some cases have to be rejected as inoperable and that some of the operations are of a mutilating character; but, again, the operation has succeeded in some cases after the failure of radiation.

On the other hand, at any rate until recent years, the results obtained by treatment with x rays or radium in this class of lower pharyngeal or extrinsic laryngeal tumour were almost negligible and the condition of many patients was aggravated. It is in this class of case, in which the operations are particularly difficult and require ceaseless care and patience in the subsequent management, that an alternative treatment by x rays or radium would be welcome if a reasonable percentage of successful results can be obtained.

In the group of intrinsic cancers of the larynx, including tumours of the vocal cords—a few being on the ventricular bands or having origin in the ventricle, and some being of subglottic origin—the operative treatment is either by laryngo-fissure (with which partial laryngectomy may be included) or by total laryngectomy. The position and extent of the growth here again determine whether simple removal of a vocal cord on one side is done, or whether the front of the larynx or the whole larynx has to be removed.

Gluck and Soerensen record, for intrinsic cancer, 141 cases of laryngo-fissure with nine deaths from the operation and ten recurrences, 122 patients remaining well for at least three years. They also record 501 cases of total laryngectomy for intrinsic cancer. This number includes a series of 100 with two deaths, and in the last 181 cases there were twelve deaths from the operation. During the last ten years there were 30 per cent. of lasting good results.

The following are particulars of operations on my own cases of intrinsic cancer.

TABLE II

	No.	Died	Recurrence or Untraced	Remain Well
Laryngo-fissure or partial laryngectomy:				
Private patients	16	1	1	14
Hospital	7	1	2	4
Total	23	2	3	18
Total laryngectomy:				
Private patients	42	2	9	31
Hospital	33	12	7	14
Total	75	14	16	45

Of these cases nine of laryngo-fissure and eighteen of laryngectomy have remained well for more than eight years, including three for more than fifteen years.

In this group, out of ninety-eight operations seventy-three were successful—that is, 74 per cent. Here, again, some of the operations are of a mutilating character, but in early cases where only a vocal cord is removed the voice is often normal and much better than after treatment by radium. This group also includes patients in whom the cartilages had been attacked by growth or in whom perichondritis had supervened, rendering them unsuitable for treatment by radiation.

It is in this group that irradiation has had most success, and therefore that line of treatment is the most attractive,

but it has to be shown that it can produce a large percentage of good results. I suggest that if radiation be employed it should be in the form of the radium beam, and that needles should no longer be used.

The figures for the operations have been given separately for the private and hospital patients because, although the numbers of each are approximately equal, the results in private practice are very much superior owing to the better condition, both general and dental, of the patients, the segregation of the patients in private rooms, the avoidance of changes in nursing staff, and the constant personal supervision of the surgeon.

In introducing this discussion I wish to avoid undue emphasis upon any particular method of treatment. This is particularly dangerous in discussing malignant disease, because in practice individual cases have to be considered, and, whatever the treatment, the prognosis of any one case is always doubtful, however favourable it may appear on the one hand or unfavourable, short of being quite hopeless, on the other. There is therefore, when failure ensues, often a feeling of regret that different treatment was not instituted, especially when irradiation fails to cure an operable case. Operation, too, might have failed, and I am concerned only to bring forward some points in the hope that the ensuing discussion may help to clarify the position.

I have to thank Mr. Arthur Pereira, registrar at St. George's Hospital, for his help in following up the fate of the patients and in preparing the statistical tables.

THE DIAGNOSIS OF UNDESCENDED TESTICLE

BY

DENIS BROWNE, F.R.C.S.

Surgeon to the Hospital for Sick Children, Great Ormond Street, London

There are few conditions upon which more contradictory diagnosis and advice can be obtained from the medical profession than those which fall into the vague and elastic category of "undescended testicle." I know a parent who was told that her son's testicles:

(a) Needed immediate operation if he were not to be practically a eunuch. (A surgeon of great experience.)

(b) Should be left for five years to see what happened. (Another reputable surgeon.)

(c) Should be treated by hormone injections. (A physician.)

(d) Should on no account be operated upon, as the operation invariably failed. (The family general practitioner.)

(e) Were normal. (Myself.)

(f) Were held up owing to a displaced vertebra. (An osteopath.)

It is hardly necessary to say that the last diagnosis and the treatment that followed are now held responsible for the present obviously satisfactory condition of affairs. Admittedly this is an extreme case, but others like it are only too common.

When I first began to see numbers of these cases I was continually in doubt as to what advice to give, but latterly they appear to me to belong to certain well-defined groups. It may be of interest to put forward my present classification: I am far from thinking it a final one, but it is certainly much better than anything I had to start with. One of my difficulties was that the testicle in almost all its varied positions is extremely mobile, so