SURGICAL PROCEDURES IN GENERAL PRACTICE

This is one of a series of articles contributed by invitation

INJURIES TO THE ELBOW-JOINT

BY

S. L. HIGGS, F.R.C.S.

TENNIS ELBOW

Much has been written but little is known with certainty about this common minor ailment. It is an annoying condition, not only to the patient but also to the doctor, as treatment so often fails to effect a cure, with consequent dissatisfaction to all concerned.

Diagnosis

The condition may sometimes be mistaken for arthritis of the elbow or some other painful disorder; but the diagnosis may be made in a few moments, as the signs and symptoms are definite and vary but little. The patient complains of pain at the outer side of the elbow and sometimes down the forearm on gripping with the fingers, as in shaking hands or opening a door, and on such movements as lifting quite small objects—a teapot or even a tumbler. Tennis and other games requiring a firm grip become impossible.

On examination the appearance of the arm is quite normal. The elbow should be moved passively, with the muscles relaxed. No pain or stiffness is experienced, so that the joint itself is thereby excluded as the site of the trouble. On active extension of the wrist against resistance, especially if this movement is combined with supination, the typical pain occurs. If this active extension is maintained the site of the pain can be localized. There are three usual situations, all of them beneath the common origin of the extensor-supinator group of muscles. The first and most usual is over the radio-humeral joint; the second, rather higher up, in the region of the external epicondyle; and the third, and least frequent, rather more forward beneath the bulky part of the muscle origin. The moment the active contraction is relaxed the pain disappears. An exactly analogous condition is very occasionally found on the inner side of the elbow just below the tip of the internal epicondyle at the origin of the flexor-pronator group of muscles.

The condition is rarely seen in patients under 35. The trouble is provoked by vigorous use of the arm, as in playing tennis, fly-fishing, and gardening. Although sometimes seen in industrial workers, it is the occasional over-use, as in games, rather than regular activity which seems to determine an attack. There is often a history of "rheumatism," lumbago, sciatica, and so on, suggesting that an infective factor may be at work. Frequently the history is given of a tennis elbow lasting two or even three years, but rarely longer. The trouble seems to run a certain course and then to disappear spontaneously.

These observations seem to throw some light on the nature of the disorder and to indicate the line of treatment. When the elasticity of youth has passed, and at an age when focal sepsis is common, unaccustomed vigorous use of the forearm muscles may lead to trauma to their deep fibres of origin. Instead of clearing up in the manner of an ordinary strain the painful symptoms persist, owing

it may be supposed, to a low-grade infective process becoming established in the damaged tissues. In a case explored by operation eight weeks after the onset of the trouble a patch of granulation tissue was discovered—an observation which lends support to this theory.

In course of time the active stage of inflammation passes into one of fibrosis and adhesions, and ultimately the lesion heals and symptoms disappear. In these later stages a chronic inflammatory area resembling wash-leather may be discovered at operation.

Treatment

The multitude of methods that have been advocated only indicate the uncertainty of the results. Many patients are seen who have already tried a number of remedies—massage, "bone-setting," and the like—and are disappointed at their lack of success. Some account of the nature of the trouble should therefore be given and its chronic character and ultimate spontaneous disappearance explained. The advice, however, will depend chiefly on the duration of the trouble when the patient is first seen.

At the onset, or if it has been present for a few weeks only, it is sometimes possible to cut short the attack by rest, heat, and avoidance of the painful movements. The arm should be kept in a sling and heat applied by antiphlogistine at night. In addition diathermy may be given. The elbow-joint should, of course, be moved through its full range each day to prevent stiffness, but the forearm muscles must be kept out of action. Focal sepsis should be sought and an x-ray examination of any crowned, dead, or suspicious-looking teeth carried out and apical infection dealt with if discovered. Massage, movements, and electrical stimulation appear to make matters worse rather than better, and should be avoided. Most patients do not seek advice at the early stage, but only when the condition has been present for some months.

In the early chronic stage, which may last a year or more, treatment is most difficult. This is rather to be expected if we believe that a localized chronic inflammatory process is at work. During this period there is no actual contracture of the affected muscles, and this would explain the frequent lack of success of rapid manipulative methods. It should be pointed out that a very certain cure within a short space of time can be effected by means of an operation. The majority, however, will not welcome this suggestion, as they regard a tennis elbow as a minor nuisance not worthy of such drastic treatment. Nevertheless there are some who find a disabled arm such a hindrance to sport or work that they will agree to operative measures, especially when other means have failed, and for them a speedy cure is assured.

The Operation

The whole arm is given the usual thorough preparation. Then, to render the operation field bloodless, an Esmarch bandage is bound round from the hand upwards and finally made fast by a few turns about the middle of the upper arm. The lower part of the bandage is then unwound and the elbow held in mid-flexion with its inner side resting on a sandbag. In no circumstances

may the rubber bandage be kept on for more than half an hour, but far less time than this should be sufficient.

A slightly curved incision four inches long is made on the outer side of the elbow behind the external epicondyle. In this situation the resulting scar is hardly noticeable. The skin is retracted forwards and then the muscles of the extensor-supinator group are reflected from the epicondyle and the capsule of the radio-humeral joint. In some cases definitely pathological tissue is visible beneath the muscles. No special search is made, and since the object is not so much to remove anything but to relieve tension the muscles are stitched back again with a few interrupted No. 0 catgut sutures. It is as well to take off the Esmarch bandage and pick up any bleeding points before sewing up the skin. The arm is kept in a sling until the stitches are removed one week after the operation, and then gradual return to active use is allowed. The arm is usually strong and comfortable once more by the end of a further two weeks.

For the majority of patients other measures must be tried. Success, by operation or otherwise, is achieved by relieving tension at the painful muscle origin, and the method described by Cyriax is giving promising results; treatment can be carried out by a masseuse familiar with the technique. The patient's elbow is flexed to a right angle and fully supinated. Deep friction is then applied to the muscles in front of the external epicondyle over the tender area. This is continued for four or five minutes, after which the elbow is extended in full supination and the forearm adducted forcibly. Some "cracks" may or may not be elicited. The treatment is carried out three times a week, and in a short time relief should be experienced if the method is going to be successful.

In the late chronic stage, where adhesions have formed, a tennis elbow can usually be cured in a dramatic manner by a single manipulation. The suitable case can be detected in the following way. The fingers and wrist are flexed and the forearm pronated. The elbow is then extended. By these means the affected muscle group is put on the stretch. If the range of extension is limited and causes pain at the elbow we may assume that contractures have formed and that the case will respond to forcible stretching. The manipulation (Mills's method) can be done without an anaesthetic, less painfully after an injection of 2 c.cm. of 2 per cent. novocain into the area of maximum tenderness, but best of all under anaesthesia from gas or from an evipan injection. Only a moment's relaxation is required. The muscles are put on the stretch in the manner described. With one hand the operator grasps the flexed fingers and wrist, and with the other the elbow, placing his thumb behind the radio-humeral joint. The elbow is then forcibly extended, and simultaneously the flexion of the hand and pronation of the forearm are increased to the full extent. A "crack" is usually heard and felt in the region of the muscle origin at the elbow. No after-care is necessary, and the patient should be encouraged to resume active use of the arm.

DISLOCATIONS AND SPRAINS

Next to the shoulder-joint the elbow is most frequently dislocated. Unlike the former it occurs more often in children than in adults. It is an injury fraught with serious consequences because of the great damage that inevitably occurs to the ligaments and their periosteal attachments and to the neighbouring muscle insertions. By far the commonest type is the posterior dislocation, due to a fall on the outstretched arm, causing hyper-

extension at the joint and levering of the coronoid process backwards over the articular surface of the humerus. More rarely the elbow is displaced sideways, either inwards or outwards, but this is usually accompanied by fractures. Anterior dislocation is rare.

The diagnosis is as a rule quite obvious from a consideration of the deformity and position of the bony points. What is not so evident by clinical examination alone is whether or not there is associated bony injury. An x-ray examination is therefore most necessary for the purpose of both treatment and prognosis. If, however, x-ray apparatus is not available, then to avoid delay and suffering the elbow should be reduced, but radiographs must subsequently be taken. Frequently, in a posterior dislocation, a fracture of the coronoid process or radial head will be revealed, and in a lateral dislocation more serious fractures involving the condyles may be seen.

Treatment

In fresh cases reduction is easy, the ligaments and soft structures being so torn that little resistance is offered except by the bony contours. It is more satisfactory in every way for the reduction to be carried out under a general anaesthetic, affording good relaxation and ample time. In a posterior dislocation the coronoid process of the ulna, which has slipped back beneath the lower end of the humerus, must be made to retrace its path. An assistant steadies the upper arm, and the forearm is grasped and pulled firmly but gently in the line of the limb. If the dislocation is of several days' duration, or difficult to reduce, then hyperextension will lever the coronoid process over the humeral surface and the joint will be felt to slip back into place. The arm is then put up in fairly full flexion, but as the swelling may give rise to pressure symptoms this must not be overdone or forced. A period of rest in the flexed position is most necessary, not only to allow the swelling to subside, but also to prevent recurrence of the displacement to save the torn ligaments from further trauma during the first stage of repair. A sling, with early movement, is not a wise line of treatment. It is better to immobilize the elbow in a moulded plaster. A four-inch or six-inch plaster bandage, depending upon the size of the arm, is immersed in water and, keeping it wet, a "slab" of the requisite length from upper arm to wrist is prepared. It is then put on straight over the skin to the back of the upper arm, elbow, and forearm, and as it sets it is fixed in position by a lightly applied domette bandage. This should leave the elbow region free in front, in order to prevent pressure and provide for inspection of any swelling. The whole is supported by a sling. In uncomplicated cases this splint should be retained for a fortnight and then replaced by a collar-and-cuff sling for a few days until the elbow gains the position of a right angle. This is followed by an ordinary sling for at least a further week.

The complications to be feared are the formation of new bone around the joint at the site of the torn-up periosteum, ossification in the ligaments, and myositis ossificans in the tendon of the brachialis anticus. The danger of these may be reduced but not eliminated by careful treatment, and therefore when the case is first undertaken it is advisable to inform those concerned of these possibilities and to record them in writing.

During the period when a sling is worn, and also later when it has been discarded, restoration of movement should be left to the patient's own efforts. Massage,

stretching, and forcible measures of any kind must be absolutely forbidden, as such misguided attempts, instead of hastening matters, may only lead to further trauma and increased stiffness and even to periarticular ossification that would otherwise not have taken place. Return to normal movement is usually slow, but no anxiety need be felt provided that the range steadily improves. Should this become stationary or the stiffness increase, then an x-ray examination must be made to determine whether ossification is occurring in the neighbourhood of the joint.

Treatment of Complications

The radiographs may show a shadow above the coronoid process of the ulna, indicating the onset of myositis ossificans in the brachialis anticus muscle. Complete fixation of the joint for a long period is essential. This is best carried out by applying a wet plaster slab direct to the back of the arm with the elbow flexed to a right angle, the plaster being held in place by an evenly applied bandage. Finally a few turns of a plaster bandage surround the whole. In this way immobilization is assured, with maximum strength and minimum weight of plaster. The hand is left free and can be used. At intervals of one month fresh radiographs are taken after removal of the plaster case in order to observe progress. The extent of the ossification continues for a time, reaches a maximum, and then begins to wane. It may be six months or more before the condition becomes stationary, and during the whole of this time the plaster fixation must be continued. Absorption of the new bone is rarely complete. Usually a branch-like growth of well-defined bone persists. The elbow is given its freedom, but no treatment is desirable. With time and active use a certain range of movement will return. In most cases this range is adequate but not complete. If, however, the limitation is serious even after a full six months of activity, and it is clear that the bony growth has formed a block, then this should be removed by operation. Too early surgical intervention will only lead to recurrence of the ossification, so patience and careful judgment based on the x-ray findings are necessary.

Ossification in the ligaments, or bony spurs at the periosteal attachments, also require rest and avoidance of any form of irritation by massage or by passive movements. Complete immobilization in plaster is not necessary, but a sling should be worn for a month or more until there is radiographical evidence that the new bone formation has ceased.

Treatment of Old Dislocations

Although reduction of a dislocation is very easily accomplished soon after the injury, with every day's delay it becomes more difficult. In these old unreduced cases another method must be employed, as forcible stretching with the elbow in extension may not only fail to effect reduction but may also cause further injury to the soft tissues. To obtain the necessary purchase the upper arm is attached by means of a loop of broad firm bandage to a radiator or to some fixed point on the wall. Strong traction is then applied to the fingers and thumb with the elbow bent at a right angle. It is difficult, however, to get the necessary grip, and Böhler's method of applying the pull through gauze stuck to the patient's fingers and thumb with "mastisol" is most valuable. By these means steady force can be applied and the coronoid process of the ulna made to retrace its path beneath the lower end of the humerus. The after-treatment by

a period of fixation followed by increasing activity is the same.

Reduction by open operation is rarely either necessary or satisfactory, as it entails so much soft-part injury and subsequent stiffness.

Sprains

A "sprained" elbow should not be regarded so lightly as a similar injury to other joints. An x-ray examination is advisable, since the general swelling and stiffness may conceal a fracture of the head of the radius or other bony damage, neglect of which may lead to serious consequences. Strapping, active use, and physical treatment may suffice for other sprains, but in the case of the elbow rest and an unhurried and unaided return to activity is the right policy. It cannot be too strongly emphasized that stiffness must not be treated by manipulation, massage, or forcible stretching.

LONDON FEVER HOSPITAL OPENING OF THE NEW BLOCK

H.R.H. the Duke of Kent on July 14 opened a new isolation block, which has been added at a cost of £30,000, to the London Fever Hospital in Islington. This replaces an older building which was in its time considered the most up-to-date isolation block in the world.

The New Isolation Block

The new building, different from the old in almost every respect, is the result of many years' experience of fever hospital administration. It is a three-story structure, with twenty-three beds, so designed that each patient is isolated, and it will act principally as a clearing house for incoming patients whose symptoms have not been clearly defined. Doubtful cases will be observed until they can be placed in their appropriate category, when they will be transferred to the corresponding ward. There may also be patients with more than one disease who will need single-room accommodation.

Each of the three floors of the block has a wide veranda in front, continuing around both ends of the building and forming a service corridor behind; thus there are no closed spaces which in old-fashioned buildings were responsible for a certain amount of room-to-room infection. Except for the top floor, with its two four-bedded wards, each floor is of the same design, arranged in single rooms. The rooms are separated by glass partitions, which give the nurse a view of the whole length of the floor from the central kitchen, though curtains afford privacy when necessary for adult patients. A portable bath sink enables each patient to have a bath in his own room. Each patient has his own thermometer, torch, spatula, basin, treatment tray, and doctor's, nurse's, and visitors' gowns. The washing and sterilizing plant in the central kitchen is cast in one piece of stainless steel. The crockery is washed in one sink, and after rinsing transferred in specially designed racks, to be sterilized in a steam-jacketed tank. There are many such interesting contrivances and safeguards in this modern building.

History of the Hospital

The London Fever Hospital was the first voluntary fever hospital in this country, and is still the only one of its kind in the London area. It was formed as a result of the deplorable social and hygienic conditions which existed in London 140 years ago. Like many other movements of the time it originated in a tavern—the Thatched House Tavern—where a meeting of citizens was held in 1802 to institute means of dealing with the contagious diseases then so prevalent. Its original home was in Gray's Inn Lane, after which it moved to a site now occupied by King's Cross station. It has been on its present site in Islington for