CHRONIC MASTITIS AND BREAST CANCER**A FAMILY HISTORY OF FIVE SISTERS**

BY

W. SAMPSON HANDLEY, M.S., F.R.C.S.*Consulting Surgeon to the Middlesex Hospital*

(WITH SPECIAL PLATE)

This history of a family of five sisters pursued by breast cancer covers a period of about nineteen years, during which, with regrettable gaps, hesitations, and refusals of advice, I have been jointly responsible with their own doctors for their defence against the family foe. It is known that their mother died at the age of 69 of cancer of the liver, and her mother at the age of 37 of abdominal cancer, probably gastric, since it was attended by persistent vomiting.

In advising this family I have been guided by the following convictions.

- (a) Chronic mastitis is a precancerous condition.
- (b) Chronic mastitis is often amenable to deep x-ray treatment without operation, and consequently deep x-ray or radium treatment may be an effective prophylactic against cancer.
- (c) The protection afforded by x rays is neither absolute nor permanent, and cases so treated should come every three or six months for re-examination.
- (d) Cases of chronic mastitis in immediate danger of cancer present recognizable clinical peculiarities, and should generally be treated by mastectomy.
- (e) Breast cancer should be treated by radical operation, with simultaneous radium to the internal mammary glands and subsequent moderate prophylactic x-radiation. Exceptionally, Keynes's radium method may be preferable.

Five separate breast cancers in the family have been treated by radical operation—four by myself and one by Mr. Turner Warwick; and in addition two simple mastectomies have been done—one for incipient cancer, the other for precancerous mastitis. In the result, two of the sisters have died of breast cancer and three are well at the time of writing. The two sisters who died survived operation eleven and nine years respectively. These results, in the face of intense predisposition to the disease in a bilateral form, offer encouragement to my belief that routine mastectomy for chronic mastitis—a policy equally distasteful to patient and surgeon—can be safely rejected in favour of x-ray treatment followed by periodic examination, with mastectomy in reserve if danger signals show themselves.

If I have described these cases at length it is partly because close personal observation of such a group over such a long period is perhaps unique, but mainly because in these histories the whole important question of the prevention of breast cancer is involved. To avoid a running commentary upon the case histories I here submit some general remarks upon the nature and treatment of chronic mastitis.

Clinical Characters of Chronic Mastitis

It is rare for the induration of chronic mastitis to affect the whole breast. The upper and outer quadrant of the breast is the seat of election, as for carcinoma, and the fact emphasizes the genetic link between the two conditions. The indurations of chronic mastitis affect the breast by lobes, and consequently, like the lobes themselves, they are sector-shaped, broader or narrower according to the number of adjacent lobes affected, often multiple in the same breast, and sometimes symmetrical in the two breasts.

For their appreciation a special technique of examination is necessary, resembling that of a cat treading its

bed preparatory to lying down. A displacement of the swelling upon the pectoral, produced by alternate pressure of the two hands lightly laid upon the skin, enables its outline and character to be accurately determined as it slips to and fro under the fingers. Except in the cystic form of mastitis, it is not a defined lump but a finely granular induration. None of the signs of contraction or adhesion is present. The glands, if enlarged, are soft and often tender.

Chronic Mastitis—a Precancerous Condition

It is impossible within the scope of this paper to marshal the evidence that chronic mastitis tends to end in carcinoma. In my book (Handley, 1931) I believe I have proved that chronic mastitis and eventually carcinoma arise within local areas of lymph block in particular lobes of the breast. The lymphatic block results from lymphangitic surface infections, often unnoticed, of the main ducts near the nipple. A state of papillomatous hypertrophy within the smaller ducts is seen in the later stages of chronic mastitis, as Lenthal Cheatle showed, and this is the immediate precursor of carcinoma. All the stages of the process, from simple lymph stasis through chronic mastitis to early carcinoma, can often be demonstrated in the same breast.

Hormones and Chronic Mastitis

Normal physiology—the rapid development of the breasts at puberty and the recurrent monthly congestion and tenderness of the breasts just before each menstrual period—demonstrates the important influence upon the breast of the ovarian hormones. Lewis and Geschickter (1934) showed that the injection of oestrin into the breasts of male monkeys produced a feminine type of breast, a condition of gynaecomastia. Lacassagne (Cheatle, 1935), experimenting on young male mice, found that a course of oestrin injections produced in some weeks a cystic mastitis, terminating in carcinoma. Cramer and Horning (1936) showed that heavy doses of oestrin produce degeneration of the adrenals, together with an initial stimulation, followed by subsequent destruction, of the anterior pituitary. Quite recently they have found that these changes in the anterior pituitary produced by oestrin can be kept in check if the thyrotropic hormone of the anterior pituitary is injected in suitable doses. They further find that the administration of this thyrotropic hormone to females of a strain of mice in which half the females die of spontaneous mammary carcinoma affords apparently complete protection against the disease (Cramer and Horning, 1938). The therapeutic hint of this striking piece of work is obvious, but it will take some years to test the suggestion it contains. The lymph-stasis theory of chronic mastitis and cancer to which I have referred is perfectly consistent with the view that hormones play an essential part in the causation of these diseases. It is obvious that local areas of lymph obstruction must necessarily be areas where the hormone supply is defective, for the regular flow of the tissue fluid through these areas is more or less arrested. Hormones may fail to reach such areas, or may accumulate in them to excess. Lymph stasis supplies the localizing factor which determines the site where the cancer arises.

Treatment of Chronic Mastitis

It has been fashionable of late years to treat chronic mastitis empirically by oestrin. The experimental evidence just detailed would seem to show that the treatment is irrational, and possibly dangerous. I have recently seen a case of cystic mastitis in which the patient herself was convinced that her breast trouble dated from a course

of oestrin injections. Whether treatment with anterior pituitary thyrotropic hormone will succeed in preventing human cancer is a question yet to be decided. It may prove difficult, as in the case of tuberculin, to get the remedy to the actual seat of the disease. Meantime we have at our disposal a method of proved value.

X-RAY TREATMENT

In 1910 I advocated the x-ray treatment of chronic mastitis, and ever since, in hundreds of cases, I have advised its use instead of mastectomy, with good results. Pain promptly disappears, and in a few weeks obstinate mastitic indurations often completely vanish. The proof that by this treatment many breast cancers have been prevented is naturally very difficult, but the evidence, which I reserve for another paper, raises a presumption that such is the truth.

Danger Signals in Chronic Mastitis

The characters of chronic mastitis in young women have just been described. As the cancer age approaches, the disease may show one or more of the following danger signals, all of them dependent upon the formation of adventitious fibrous tissue in the affected lobe or lobes. These signs cannot here be described in detail. They are:

NIPPLE CHANGES

1. Retraction of one or more of the duct orifices upon the nipple, without retraction of the nipple itself.
2. Serous or blood-stained discharge from the nipple.
3. Deviation of the axis of the nipple.
4. Elevation of the nipple.
5. Appearance upon the nipple of white or black dots from which plugs of degenerate epithelium can sometimes be squeezed.
6. Flattening of the nipple.
7. Retraction of the nipple.

BREAST CHANGES

1. Appearance of multiple hard, elastic, or fluctuant lumps in the mastitic area (cystic mastitis).
2. The presence near the nipple in the mastitic area of radiating tortuous cords which are thickened main ducts.
3. Adhesion of the breast to the skin or the pectoral fascia is rarely seen except in fully developed carcinoma.

Case Histories

The case histories, subsequently to be given in detail, are summarized in the following table:

	Date of Birth	Right Breast	Left Breast	Result
Sister A	1871	Mass appeared in 1930 just before death. (Second primary carcinoma)	Primary carcinoma, 1919. (Radical mastectomy)	Died 1930 with signs of pyloric obstruction due to a primary gastric growth or to secondary abdominal masses
Sister B	1875	Primary carcinoma, 1925. (Radical mastectomy)	Primary carcinoma, 1937. (Radical mastectomy)	Alive and well, 1938
Sister C	1879	Proliferative mastitis and incipient carcinoma, 1933. (Simple mastectomy)	Proliferative mastitis, 1930. (Simple mastectomy)	Alive and well, 1938
Sister D	1885	Mastitic induration, 1930. (Surface radium treatment)	Mastitic induration, 1929. (X-ray treatment)	Alive and well, with normal breasts, 1938
Sister E	1886	Second primary carcinoma, 1927. (Radical mastectomy)	Primary carcinoma, 1923. (Radical mastectomy)	Died in 1932 of recurrence in supraclavicular glands, sternum, and spine

SISTER A; BORN 1871; UNMARRIED

In 1919 I operated upon this patient for a carcinoma in the lower outer quadrant of the left breast. The growth had been present since 1918, was 2½ inches in diameter, and was adherent to skin and muscle, though clinically the axillary glands were not enlarged. Microscopical section showed a regular spheroidal-cell carcinoma with considerable tubule formation, but the axillary glands were free from growth.

She remained in good health until 1929, when she began to suffer from arthritis of the spine and hip. Radiographs showed no evidence of secondary bone deposits. In November of the same year I again saw her; she was then suffering from a neoplasm of the stomach, with pyloric obstruction. Her condition was hopeless and did not admit of radiography of the stomach. It was impossible to say whether the neoplasm was primary or secondary. She died in January, 1930; her doctor, Dr. A. M. Barlow, reported that a few weeks before death a lump (? a second primary carcinoma) appeared in the right breast.

On the whole it appears likely, since her mother possibly and her grandmother probably died of gastric carcinoma, that death was due to a primary neoplasm of the stomach.

SISTER B; BORN 1875; MARRIED

The patient was first seen in 1920. She had had an abscess in the left breast after a miscarriage, and a narrow sector-shaped mastitic induration containing two small nodules, thought to be cysts, was found above and internal to the nipple. The left nipple was elevated, but in view of the history of abscess the importance of this sign was minimized. The right breast was normal. I asked to see her in six weeks' time, but it was not until 1925 that she again appeared, with a typical carcinoma of the right breast. Vague induration could be felt in the left breast at the site of the cysts, but it was otherwise normal. A right radical mastectomy was done, and microscopical examination of the specimen showed a carcinoma with no involvement of the axillary glands.

In 1927 she was well, but a thickened duct could be felt a short distance above the left nipple. Deep x-ray and surface radium treatment was given during 1927 and 1928. In 1930 a precautionary left mastectomy was advised, but the patient declined operation, and another course of deep x-ray treatment was given. In 1934 thickened ducts had again appeared, and operation was urged but without success.

In 1937, five years after the last x-ray treatment, a hard lump adherent to the skin appeared in the upper and outer quadrant of the left breast just outside an area of x-ray telangiectasis present above the nipple since 1931. Mr. Turner Warwick performed a radical mastectomy, and pathological examination showed a cellular spheroidal-cell carcinoma without axillary gland invasion. The patient made a good recovery from the operation and is at present well. It is interesting that the carcinoma appeared outside the area of telangiectasis. I have never seen a carcinoma develop beneath such an area, but I can recollect at least one other case in which a carcinoma appeared just beyond the margin.

The history of this case shows that the protection against carcinoma afforded by x rays is not absolute and permanent. It also emphasizes the need for three-monthly or six-monthly re-examinations of the abnormal breast, and for repeating the x-ray course at least every two or three years. It is to be noted that the breast was watched and treated for fourteen years before its removal was urged, and that frank carcinoma did not supervene until three years later. This is prima facie evidence that it is possible to distinguish the onset of the dangerous stage of chronic mastitis.

SISTER C; BORN 1879; MARRIED

In 1927 this patient was sent to me by Dr. Barlow for pain in and under the left breast. She had cardiac weakness and irregularity following influenza, with a bruit over the lower sternum, and was subject to indigestion. All remaining teeth had been extracted a year before.

On examination a sector-shaped mastitic induration was found in the upper quadrant of the left breast. In it near the nipple could be felt tortuous thickened ducts. The left nipple was higher than the right and pointed forwards and outwards, while the right pointed normally—that is, forwards, outwards, and downwards. Owing to these signs I regarded the breast as a dangerous one, but in view of the cardiac condition x-ray treatment was advised in preference to mastectomy. A month later the left breast appeared almost normal, except that some thickened ducts persisted above the left nipple. On this occasion some thickened ducts were felt above the nipple of the other (right) breast. In June, 1928, both breasts were much improved. In May, 1930, there was a slight duct thickening above the left nipple, and further x-ray treatment was given to both breasts by Dr. Russell Reynolds. The left nipple was still raised and deviated.

In September, 1930, following a blow on the left breast, a small lump a quarter of an inch in diameter and adherent to the skin was found in the left upper and outer quadrant. I diagnosed carcinoma. The pathologist reported that the lump was a cyst and that the breast only showed chronic mastitis. A simple mastectomy was done. Re-examination of the slide has convinced me that the duct epithelium was in a precancerous condition.

In March, 1933, a mastitic swelling containing thickened tortuous ducts appeared above the right nipple, and I urged a simple mastectomy, with removal of the glands. This was done. The breast showed advanced fibrosis with microcystic distension of the smaller ducts and proliferation of their epithelium. At certain points the breast showed diffuse papillomatosis of the ducts, still innocent (Plate, Fig. 1). At one point there was seen a duct distended by epithelium, with definite infiltration of the surrounding tissues—an incipient carcinoma (Fig. 2). Glands were not infected. Subsequently a course of x rays was given.

This case confirms the evidence of the preceding one as to (1) the controlling power of irradiation in chronic mastitis; (2) the need for periodic examinations in such cases; (3) the value of the danger signals which herald approaching malignancy.

SISTER D; BORN 1885; MARRIED, WITH TWO CHILDREN

The patient first presented herself in 1929 because of discomfort and pain in the left breast of twelve months' duration. Examination showed a slight mastitic thickening in the upper inner quadrant of the left breast, which was tender. There were no other signs in either breast. The induration disappeared after a short course of x-ray therapy.

In 1930 an indurated area was noted in the region of the axillary tail of the right breast, but this also subsided under surface radium treatment. The breasts were normal when the patient was last seen in October, 1937.

SISTER E; BORN 1886; MARRIED

This patient was operated upon by me at the Middlesex Hospital in December, 1923, for an advanced carcinoma of the left breast when aged 37. Radium tubes were introduced in the upper intercostal spaces and a patch of radio-dermatitis developed a month later over the inner end of the third space.

She does not appear to have been seen again until September, 1927. At this time two patches of radio-dermatitis were present at the inner ends of the third and fifth left spaces. In the right breast in the upper and outer quadrant was a defined granular sector-shaped thickening, an area of chronic mastitis. The right nipple seemed less prominent than normal, and was flame-shaped in contour in the plane of the skin, with the point directed towards the mastitic thickening. There was no definite adhesion to the skin, but a very faint depression appeared over the induration when the breast was displaced. I advised removal of the right breast, but as an alternative, owing to the reluctance of her relatives, offered x-ray treatment, with re-examination in two months. The axillary glands were not enlarged. There was no recurrence

on the left side. I wrote to her medical adviser, Dr. Barlow: "There seems to be some danger that this patch will develop into a carcinoma, and removal of the right breast would be a wise precaution, since the second breast in a patient who has had carcinoma is four times as likely to develop carcinoma as the breast of a normal woman."

By October, 1927, after one dose of deep x rays which gave her acute pain in the breast and made her ill for some days, the patient had made up her mind to undergo operation. A faint but unmistakable skin dimple could now be elicited over a small lump beneath the edge of the right great pectoral at the periphery of the breast. The mastitic swelling had become less definite, probably as the result of the x-ray dose. The flame shape of the nipple was still evident. A diagnosis of second primary carcinoma was made, and immediate radical operation was urged and performed. Histological examination showed a spheroidal-cell carcinoma, without invasion of lymph glands, considered to be a second primary carcinoma.

In June, 1928, she was free from recurrence though suffering from anginal attacks, but she then escaped from observation. In May, 1930, I was informed that she was having radium treatment for enlarged glands in the neck. A sternal recurrence was subsequently treated by buried radium. In 1932 spinal secondary deposits appeared, and she died in 1934, after being bedridden for two years.

In the history of this case the regrettably long intervals between examinations are to be specially noted.

Commentary

The history of this unfortunate family recalls the high incidence of breast cancer found in certain inbred laboratory strains of mice. Dr. W. E. Gye (1936-7) makes some interesting observations on these strains. He says:

"Inbred strains of mice are very useful in some branches of experimental cancer research. The discovery of the carcinogenic property of oestrone was made with such a strain, and the analysis of the chain of events which ends in malignancy of the breast is simplified by the use of pure strains of high and lower cancer incidence. But, as Dr. Cramer has pointed out, inbred mice are not normal animals, and in high cancer lines the clinical picture of cancer is distorted by the predominating influence of one aetiological factor—inherited susceptibility in a particular organ—to such an extent as to minimize the importance of other factors. Conclusions drawn from observations on such strains cannot, therefore, be applied without reserve to cancer of normal mixed stocks. In high cancer strains, for example, multiple tumours of the breast are of frequent occurrence, whereas in mixed stocks single tumours are the rule. If a cancer in a high strain is excised surgically, the animal, though cured of this particular tumour, is liable to develop a second or a third independent growth in another part of the same breast or in the breast of the opposite side. The inherited predisposition to cancer of the breast is not affected by the removal of the first tumour, and surgical treatment in these exaggerated artificial circumstances is ineffectual and appears to be irrational. In mice of mixed strains surgical removal of a tumour is not followed by the appearance of fresh independent new growths; if local recurrence takes place it is due to incomplete removal of the primary tumour. The differences between cancer in pure line strains and in mixed stocks may have their counterpart, exhibited less strikingly, in human surgery."

How far are these conclusions supported for human carcinoma by the facts I have related? The fact that three of the five sisters are still living, free from pain and from signs of recurrence, partly as the result of radical mastectomies for developed cancer, partly as the result of simple mastectomies for threatening precancerous conditions, shows that in the human race (which fortunately has only two breasts) surgical treatment may be worth

while, however strong the constitutional predisposition to breast cancer. Though I believe radiation to be a valuable prophylactic, it cannot in all cases be an efficient substitute for surgery; indeed, the impermanence of the protection it affords is manifest in the case records.

Dr. Gye's statement that "in mice of mixed strains surgical removal of a tumour is not followed by the appearance of fresh independent new growths" is far from being applicable in the ordinary case of human breast cancer with no specially strong family history. My experience is that I see an increasing number of cases of second primary carcinoma of the opposite breast. I hope to diminish that number in future by advising routine post-operative irradiation of the remaining breast, and by even more careful following up of cases.

Summary

This paper records the history of five sisters, descendants of a cancerous mother and grandmother, who have been under my observation for nineteen years. There is evidence that three of them have suffered from double primary carcinoma of the breast. One sister was treated by irradiation for bilateral chronic mastitis, which has not recurred. The remaining sister was treated for chronic mastitis by irradiation, at first with success, but after three years a left simple mastectomy seemed necessary, and three years later a right simple mastectomy. The right breast showed an incipient carcinoma. The danger signals which herald the onset of carcinoma in such cases are described, and it is held to be possible to distinguish between cases of chronic mastitis which may safely be treated by irradiation and cases in the precancerous stage needing mastectomy. Two of the sisters have died of breast cancer, surviving operation eleven and nine years respectively, and three remain well. Five radical mastectomies and two simple ones have been done.

REFERENCES

- Cheatle, Sir G. Lenthal (1935). *Brit. J. Surg.*, **22**, 710.
Cramer, W., and Horning, E. S. (1936). *Lancet*, **1**, 247.
— (1937). *J. Path. Bact.*, **44**, 633.
— (1938). *Lancet*, **1**, 72.
Gye, W. E. (1936-7). Imperial Cancer Research Fund Report, p. 18.
Handley, W. Sampson (1910). *Practitioner*, **84**, 463.
— (1931). *Genesis of Cancer*, H. K. Lewis, London.
Lewis, D., and Geschickter, C. F. (1934). *Ann. Surg.*, **100**, 779.

Dr. Henry Herd, school medical officer for Manchester, in his annual report for 1937, records the transfer to his department of a child guidance clinic which was formerly run on voluntary lines. The present tendency to make use of the school clinics for early consultation, in preference to awaiting the next routine visit of the medical officers, is shown by the rising ratio of "special" to "routine" inspections. During the year under review some 33,000 routine inspections were undertaken, as compared with nearly 49,000 special inspections. The nutritional grading required by the Board of Education was continued during the year. An analysis of the very different findings of a group of officers confronted with the same batch of children, the grading being based on general observation only, bears out Dr. Herd's view that the economic test, which reveals that a child can, or cannot, be receiving an adequate diet on the income available for feeding him, is still the only reliable and practicable method capable of application on a large scale to the assessment of nutrition. The scope of this report is as wide as ever, and it includes a paper on the physical fitness of schoolboys, with special reference to nutrition and the mechanism of heredity.

DYSFUNCTIONAL UTERINE BLEEDING TREATED WITH PROGESTERONE*

BY

T. N. MACGREGOR, M.D., F.R.C.S.Ed., M.C.O.G.
*Gynaecologist, Deaconess Hospital, Edinburgh; Late Tutor
in Clinical Gynaecology, Royal Infirmary, Edinburgh;
Crichton Research Scholar*

(WITH SPECIAL PLATE)

Although towards the end of last century it was suggested that irregular uterine bleeding was associated with abnormal ovarian function, it is only within comparatively recent times that this has been confirmed by scientific data. The dysfunctional nature of irregular uterine bleeding was disclosed when the endometrial structure and hormone findings in such cases were correlated with the endometrial changes and hormone concentrations occurring cyclically during the normal menstrual process. It is known that menstrual rhythm is founded on the harmonious interaction of the gonadotropic hormone of the anterior pituitary and the oestrogenic and luteal hormones of the ovaries. Several factors, however, particularly dysfunction of one or more of the other glands in the endocrine chain and abnormal nervous stimuli, may affect this intimate relationship.

The present communication records a study of thirteen cases of irregular uterine bleeding that was essentially associated with a disturbance of the pituitary-ovarian mechanism.

Types of Bleeding

Two types of bleeding only are considered—namely, "ovular" and "anovular."

Ovular Bleeding.—By this term is implied bleeding which may occur at or near the time of ovulation, about the middle of the cycle. This phenomenon, which has been noted by many observers (Hartman, 1932; Hain, 1934), is probably due to a disturbance of the hormonal equilibrium at the time of ovulation. In such cases it is presumed that between the time of rupture of the Graafian follicle and the formation of the corpus luteum there is a transient lowering of the blood concentration of oestrin. Consequent on the fall of blood oestrin there is diminished endometrial stimulation, which is sufficient to induce partial disintegration of the endometrium with bleeding. When the corpus luteum is mobilized, however, the endometrial stimulation is restored and the haemorrhage subsides.

Anovular Bleeding.—As distinct from anovular menstruation, this term is synonymous with that of metrorrhagia haemorrhagica. The pathological findings in this condition are cystic degeneration of the Graafian follicles with no recent or forming corpus luteum in the ovary, myohyperplasia of the uterus, and marked proliferation of the endometrium with cystic dilatation of the glands (Plate, Figs. 1 and 5). The cells lining the glands are cubical, with centrally placed nuclei; no glycogen is found in the cells (Figs. 2 and 6). This characteristic pathological picture may result from (a) an upset in the pituitary-ovarian mechanism or (b) a local pelvic lesion interfering with the normal process of ovulation.

Mechanism of Bleeding

Ovular bleeding, it has been suggested, is associated with a lowering of the blood concentration of oestrin at

* A Report to the Therapeutic Trials Committee of the Medical Research Council. From Dr. Douglas Miller's Gynaecological Clinic, Royal Infirmary, Edinburgh, the Deaconess Hospital, Edinburgh, and the Institute of Animal Genetics, Edinburgh University.

W. SAMPSON HANDLEY : CHRONIC MASTITIS AND BREAST CANCER : A FAMILY HISTORY

FIG. 1.—Sister C. Right breast; Diffuse innocent papillomatosis of the ducts. Magnification $\times 120$. (Photo by Bland-Sutton Institute.)

FIG. 2.—Sister C. Right breast; Incipient carcinoma. Duct distended by epithelium, infiltration beginning. Magnification $\times 200$. (Photo by Bland-Sutton Institute.)

MAX GREENBERG : PARASITIC FOETUS SUCCESSFULLY REMOVED BY OPERATION

FIG. 1.—Parasite after removal.

FIG. 2.—Child at age of 10 months.