while, however strong the constitutional predisposition to breast cancer. Though I believe radiation to be a valuable prophylactic, it cannot in all cases be an efficient substitute for surgery; indeed, the impermanence of the protection it affords is manifest in the case records.

Dr. Gye's statement that "in mice of mixed strains surgical removal of a tumour is not followed by the appearance of fresh independent new growths" is far from being applicable in the ordinary case of human breast cancer with no specially strong family history. My experience is that I see an increasing number of cases of second primary carcinoma of the opposite breast. I hope to diminish that number in future by advising routine post-operative irradiation of the remaining breast, and by even more careful following up of cases.

Summary

This paper records the history of five sisters, descendants of a cancerous mother and grandmother, who have been under my observation for nineteen years. There is evidence that three of them have suffered from double primary carcinoma of the breast. One sister was treated by irradiation for bilateral chronic mastitis, which has not recurred. The remaining sister was treated for chronic mastitis by irradiation, at first with success, but after three years a left simple mastectomy seemed necessary, and three years later a right simple mastectomy. The right breast showed an incipient carcinoma. The danger signals which herald the onset of carcinoma in such cases are described, and it is held to be possible to distinguish between cases of chronic mastitis which may safely be treated by irradiation and cases in the precancerous stage needing mastectomy. Two of the sisters have died of breast cancer, surviving operation eleven and nine years respectively, and three remain well. Five radical mastectomies and two simple ones have been done.

REFERENCES

- Cheatle, Sir G. Lenthal (1935). *Brit. J. Surg.*, **22**, 710.
Cramer, W., and Horning, E. S. (1936). *Lancet*, **1**, 247.
— (1937). *J. Path. Bact.*, **44**, 633.
— (1938). *Lancet*, **1**, 72.
Gye, W. E. (1936-7). Imperial Cancer Research Fund Report, p. 18.
Handley, W. Sampson (1910). *Practitioner*, **84**, 463.
— (1931). *Genesis of Cancer*, H. K. Lewis, London.
Lewis, D., and Geschickter, C. F. (1934). *Ann. Surg.*, **100**, 779.

Dr. Henry Herd, school medical officer for Manchester, in his annual report for 1937, records the transfer to his department of a child guidance clinic which was formerly run on voluntary lines. The present tendency to make use of the school clinics for early consultation, in preference to awaiting the next routine visit of the medical officers, is shown by the rising ratio of "special" to "routine" inspections. During the year under review some 33,000 routine inspections were undertaken, as compared with nearly 49,000 special inspections. The nutritional grading required by the Board of Education was continued during the year. An analysis of the very different findings of a group of officers confronted with the same batch of children, the grading being based on general observation only, bears out Dr. Herd's view that the economic test, which reveals that a child can, or cannot, be receiving an adequate diet on the income available for feeding him, is still the only reliable and practicable method capable of application on a large scale to the assessment of nutrition. The scope of this report is as wide as ever, and it includes a paper on the physical fitness of schoolboys, with special reference to nutrition and the mechanism of heredity.

DYSFUNCTIONAL UTERINE BLEEDING TREATED WITH PROGESTERONE*

BY

T. N. MACGREGOR, M.D., F.R.C.S.Ed., M.C.O.G.
*Gynaecologist, Deaconess Hospital, Edinburgh; Late Tutor
in Clinical Gynaecology, Royal Infirmary, Edinburgh;
Crichton Research Scholar*

(WITH SPECIAL PLATE)

Although towards the end of last century it was suggested that irregular uterine bleeding was associated with abnormal ovarian function, it is only within comparatively recent times that this has been confirmed by scientific data. The dysfunctional nature of irregular uterine bleeding was disclosed when the endometrial structure and hormone findings in such cases were correlated with the endometrial changes and hormone concentrations occurring cyclically during the normal menstrual process. It is known that menstrual rhythm is founded on the harmonious interaction of the gonadotropic hormone of the anterior pituitary and the oestrogenic and luteal hormones of the ovaries. Several factors, however, particularly dysfunction of one or more of the other glands in the endocrine chain and abnormal nervous stimuli, may affect this intimate relationship.

The present communication records a study of thirteen cases of irregular uterine bleeding that was essentially associated with a disturbance of the pituitary-ovarian mechanism.

Types of Bleeding

Two types of bleeding only are considered—namely, "ovular" and "anovular."

Ovular Bleeding.—By this term is implied bleeding which may occur at or near the time of ovulation, about the middle of the cycle. This phenomenon, which has been noted by many observers (Hartman, 1932; Hain, 1934), is probably due to a disturbance of the hormonal equilibrium at the time of ovulation. In such cases it is presumed that between the time of rupture of the Graafian follicle and the formation of the corpus luteum there is a transient lowering of the blood concentration of oestrin. Consequent on the fall of blood oestrin there is diminished endometrial stimulation, which is sufficient to induce partial disintegration of the endometrium with bleeding. When the corpus luteum is mobilized, however, the endometrial stimulation is restored and the haemorrhage subsides.

Anovular Bleeding.—As distinct from anovular menstruation, this term is synonymous with that of metrorrhagia haemorrhagica. The pathological findings in this condition are cystic degeneration of the Graafian follicles with no recent or forming corpus luteum in the ovary, myohyperplasia of the uterus, and marked proliferation of the endometrium with cystic dilatation of the glands (Plate, Figs. 1 and 5). The cells lining the glands are cubical, with centrally placed nuclei; no glycogen is found in the cells (Figs. 2 and 6). This characteristic pathological picture may result from (a) an upset in the pituitary-ovarian mechanism or (b) a local pelvic lesion interfering with the normal process of ovulation.

Mechanism of Bleeding

Ovular bleeding, it has been suggested, is associated with a lowering of the blood concentration of oestrin at

* A Report to the Therapeutic Trials Committee of the Medical Research Council. From Dr. Douglas Miller's Gynaecological Clinic, Royal Infirmary, Edinburgh, the Deaconess Hospital, Edinburgh, and the Institute of Animal Genetics, Edinburgh University.

the time of ovulation, and it is probable that a similar factor operates in metropathia haemorrhagica. Clinically and experimentally it has been demonstrated that endometrial stimulation can be maintained for long periods, provided an adequate amount of oestrogenic hormone is administered. The withdrawal of the hormone, however, is followed by endometrial disintegration. In metropathia haemorrhagica, therefore, there must be some factor which initiates a lowering of blood oestrin.

I have several times confirmed the observation made by many investigators that the large follicular cysts found in the ovaries in cases of metropathia haemorrhagica contain a fluid which has oestrogenic properties. There are therefore in these cases depots of oestrin in the ovaries which supply a continuous amount to the general circulation and thus endometrial stimulation is sustained. Gradual intracystic tension, however, may cause either atrophic changes in the secreting cells or compression of the blood vessels. Whatever the cause, the direct result is a reduction of the hormone passing into the general circulation. Although it was shown by Corner (1927) and Hartman (1932), and subsequently confirmed by others, that oestrin withdrawal causes uterine bleeding, it was not until Markee's (1936) experiments on monkeys were carried out that the mechanism of this phenomenon was demonstrated. He showed that diminution in the blood concentration of oestrin causes vasoconstriction of the endometrial capillaries. Consequent on this vascular spasm the nutrition of the highly proliferated glandular structure is impaired. Partial necrosis results, with consequent shedding of the endometrium.

Diagnosis

The history obtained in cases of anovular bleeding is characteristic. There is usually a period of six to ten weeks' amenorrhoea followed by prolonged bleeding of from four to eight weeks' duration. In cases of ovular bleeding there is a history of frequent loss, which in some cases occurs regularly every fourteen days. It is sometimes necessary to remove endometrium on several occasions in order to distinguish between the ovular and true menstrual bleeding. Although a diagnosis of these conditions may be made on clinical evidence, this should be confirmed by histological examination of the endometrium. In the early years of reproductive life a thorough examination of the genital tract, including measurement of the length of the utero-cervical canal and microscopical examination of the endometrium, furnishes information which is of value in determining the form of therapy most likely to be beneficial. This examination, including in particular biopsy, is essential near the menopause, as the possibility of carcinoma being the cause of the bleeding must always be excluded.

Treatment

Progesterone was the form of therapy used in the present investigation. Initially 1 to 2 mg. were given daily or every second day, but when more abundant supplies became available 5 mg. were given daily. In ovular bleeding the treatment was given two or three days before the expected anomalous bleeding, the aim being to maintain endometrial stimulation during the time of ovulation, and thus tide over the period when the blood oestrin was lowered. The administration of oestrin at this time, by preventing a diminution of blood oestrin, produces the same effect. As is known, however, oestrin at a certain concentration inhibits the anterior pituitary, and accordingly it is possible that this

form of therapy may, by influencing the pituitary stimulus, delay or inhibit ovulation. Theoretically, therefore, the former method of treatment is preferable.

In metropathia haemorrhagica progesterone may be given at any phase. During the bleeding stage 5 mg. were administered daily until the bleeding was controlled. During the amenorrhoeic phase 5 mg. were given daily, the total dosage amounting to 25 to 30 mg. Though oestrin therapy may, by sustaining endometrial stimulation, control the haemorrhage in these cases, the effect is only a temporary one. Indeed, oestrin would only aggravate the condition, and its withdrawal would be followed by still more haemorrhage. Accordingly, this form of therapy is definitely contraindicated.

Data and Results

Thirteen patients were treated with progesterone; five were cases of ovular bleeding and seven of anovular bleeding, whilst one case, though treated with progesterone, was found to come under another category.

Ovular Bleeding.—As the result of treatment three patients have had regular menstruation for the last eighteen to twenty-two months. One patient, studied over a period of eighteen months, had three regular periods after cessation of therapy. She then became pregnant, but aborted at the end of the second month of gestation (Chart I). The other patient, though not completely well.

CHART I.—Case 4; aged 30. Menarche at 12. Menstruation regular 7-8/28 until the age of 25. Since then menstruation has been irregular 7-21/17-21. Biopsy curettage carried out on two occasions indicated that ovulation was occurring but was associated with bleeding. The chart indicates the effect of treatment. Doses of progesterone were given on the following days: 1936, April 7, 8, 9, 10, 15, 18, 21, and 25, 1 mg.; April 27 and 28, 2 mg.; May 24, 27, and 31 and June 3, 14, 17, and 21 and August 10 and 13, 1 mg.; August 30 and September 1 and 4, 5 mg.; September 20, 23, 27, and 30 and October 19, 22, 26, and 28 and November 8, 1 mg.; November 15, 18, 22, 25, and 29 and December 2 and 6, 2 mg.

has now only a brownish discharge, whereas formerly she had frank haemorrhage for a few days between every two periods.

Anovular Bleeding.—In four cases of anovular bleeding menstruation became normal in duration and periodicity. In two cases, however, there has recently been a recurrence of the irregularity. This has been attributed by one patient to a "nervous turn" following the sudden death

CHART II.—Case 6; aged 31; 2-para. Menstruation 4-7/21-28-day type since the menarche at the age of 13 till December, 1935, when period lasted from December 5, 1935, to January 1, 1936. After an interval of four weeks, bleeding began again in the beginning of February and lasted until March 23, 1936, when a diagnostic curettage was carried out. Basal metabolic rate + 7 per cent. Blood-sugar tolerance: 88 mg. per 100 c.cm.; after 50 gm. glucose, 100, 119, 99, 90, and 90 mg. per 100 c.cm. The diagnosis of metropathia haemorrhagica was confirmed by histological examination of the endometrium. Doses of progesterone were given on the following days: 1936, May 10, 11, 12, 13, and 14, 2 mg.; May 31, 1 mg.; June 3, 7, and 29 and July 1, 2 mg.; August 3, 9, 12, 15, 17, 19, 22, and 24, 1 mg.; 1937, August 8, 2 mg.; August 9, 1 mg.

CHART III.—Case 7; aged 31; 1-para. Menarche at age of 12. Periods were regular 4/28 until birth of child two years ago. Menstruation recommenced three months after the birth and was regular 4/28 until April, 1936. The diagnosis of metropathia haemorrhagica was confirmed by histological examination of the endometrium removed on April 30, 1936. Doses of progesterone were given on the following days: 1936, November 26 and 30 and December 2, 6, and 9, 1 mg.; December 13, 15, 17, and 20, 2 mg.; 1937, February 21, 24, and 27, 2 mg.; February 28 and March 1 and 2, 5 mg.

of a near relative; while in the other case the patient sustained a shock on falling off a bus (Charts II and III). In the remaining three cases the uterine endometrium was examined histologically before and after treatment, and it is of interest to note that as a result of the treatment the hyperplastic endometrium has been changed into a pseudo-secretory phase in two of the cases (Figs. 3, 4, 7, and 8). These cases have only recently been under treatment, and two have had two normal periods, whilst the third has had one normal period.

On the other hand, the following case illustrates the wasteful aspect of the indiscriminate use of endocrine therapy for irregular uterine bleeding.

Case 13.—This patient, aged 13, was first seen on August 12, 1936. Menstruation started in February, 1936, and her periods were as follows: the first period lasted five days; the second period, five weeks later, lasted seven days; the next period, one month later, lasted a similar time; the period in May lasted seven days; two periods in June lasted seven days; a period started on July 4, 1936, and continued until the patient was seen on August 12. Thyroid therapy had no effect on the bleeding. Between August 12 and 17 the patient had 10 mg. of progestin. Bleeding stopped on August 28, but re-started on September 3 and lasted nine days; it was again present from September 24 to October 3 and came on once more on October 12. Between October 19 and 26 the patient had 35 mg. of progestin, which did not control the bleeding. Examination under anaesthesia on October 31 showed that the uterus was small, the utero-cervical canal measuring only two inches. Thereafter she was put on progynon B oleosum forte (oestradiol benzoate), and after a total dosage of 45 mg. menstruation came at regular intervals for five days and has continued to do so.

I have indicated that the progesterone treatment in this case was indiscriminate. If the patient had been examined initially under anaesthesia and the hypoplasia of the uterus detected progesterone would not have been given, as it would have had no effect on an endometrium which had not been sufficiently primed with oestrogenic hormone.

Discussion

The histological appearances of the endometrium before and after the administration of progesterone are significant, in that they demonstrate clearly the effect of this hormone on the endometrium. If this were the only action produced, however, this therapy would be of little value, as its effect would be merely a local and temporary one. I have shown that progesterone therapy, in some cases of metropathia haemorrhagica, was followed by a normal menstrual rhythm. This result, it is presumed, has been due to a re-establishment of a normal hormonal balance. The rationale of this form of therapy is possibly as follows: the progesterone brings about an increased elimination of oestrin by urinary excretion or converts it into less physiologically active fractions (Smith, Smith, and Pincus, 1938). This results in a lowering of the blood concentration of oestrin to within normal limits; following upon this, that inhibition of the pituitary which has been associated with a high concentration of oestrin is removed. The pituitary can now act at a normal level. A full

reactivation of ovarian function restores the normal hormonal balance, a continuation of which effects the regular menstrual rhythm.

Comparatively few favourable results will follow the indiscriminate use of progesterone therapy, but in cases where its employment appears to be indicated by the histological structure of the endometrium encouraging results can be obtained. Although radium therapy is successful in the treatment of metropathia haemorrhagica occurring at or near the menopause, a wider use of progesterone seems to be indicated as a more rational form of treatment.

Summary

1. Two types of irregular uterine bleeding are described.
2. The mechanism of bleeding in metropathia haemorrhagica is discussed.
3. The results of treatment with progesterone are given, and the effect of this form of therapy on the uterine endometrium is illustrated by photomicrographs.
4. The rationale of progesterone therapy in irregular uterine bleeding is discussed.

I wish to express my indebtedness and thanks to Dr. Douglas Miller for the privilege of investigating and treating cases under his care, and to Professor F. A. E. Crew for the facilities afforded me in his department.

The preparation of progesterone used in this investigation was progestin organon, and I am grateful to the manufacturers (Messrs. Organon Laboratories), who supplied it to me through the Therapeutic Trials Committee.

REFERENCES

- Corner, G. W. (1927). *J. Amer. med. Ass.*, **89**, 1838.
 Hain, A. M. (1934). *Quart. J. exp. Physiol.*, **24**, 101.
 Hartman, C. G. (1932). - Carnegie Inst. Wash. Publ. No. 433, Contr. to Embryol., No. 134.
 Markee, J. E., et al. (1936). *Anat. Rec.*, **64**, 231.
 Smith, G. van S., Smith, O. W., and Pincus, G. (1938). *Amer. J. Physiol.*, **121**, 98.

BRODIE'S ABSCESS AND ITS DIFFERENTIAL DIAGNOSIS*

BY

JAMES F. BRAILSFORD, M.D., Ph.D., M.R.C.P.

Consulting Radiologist, City of Birmingham Hospitals

(WITH SPECIAL PLATE)

In the collected works of Sir Benjamin Collins Brodie, published in 1865, will be found a paper on "Chronic Abscess of the Tibia" which was presented to the students of St. George's Hospital. In this paper he pointed out that an abscess may occur in the interior of any bone in the body, but according to his experience we meet with it more frequently in the tibia than in any other bone. He gives a brief account of eight cases. All the patients except one were males between the ages of 13 and 34 years. The abscess was in the upper third of the tibia in four and in the lower third in four. The clinical histories indicated that the lesion had existed for periods ranging from five to eighteen years. The account of his first case is as follows:

Some Early Cases

"It was as long since as the year 1824 that I was consulted by a young man, 24 years of age, under the following circumstances. There was a considerable enlargement of the lower end of the tibia, but the ankle-joint permitted of every motion and was apparently normal. The skin was thin, tense, and closely adherent to the periosteum. There

was constant pain in the part, generally of a moderate character, but every now and then it became excruciating, keeping the patient awake at night and confining him to the house for many successive days. It made his life miserable and his nervous system irritable: one effect of which was, that it spoiled his temper, and thus produced another set of symptoms in addition to those which were the direct consequence of the local malady. The disease had been going on for twelve years. He had consulted many surgeons respecting it, and had used a great variety of remedies, but had never derived benefit from anything that was done. Instead of getting better, he every year became so much worse. I tried some remedies without any advantage, and at last recommended that he should lose the limb. Mr. Travers saw the patient with me, and agreed with this opinion. Amputation was performed: and the amputated tibia is now on the table. You will see how much the lower end of it is enlarged, and that the surface of it presents marks of great vascularity. The bone in the preparation is divided longitudinally, and just above the articulating surface there is a cavity as large as a chestnut. This cavity was filled with dark-coloured pus. The inner surface of it is smooth. The bone surrounding it is harder than natural. On observing these appearances, I could not help saying that if we had known the real nature of the disease the limb might have been saved."

In another paper he tells us that the patient died a few days after the operation. The other seven patients he treated by trephining the bone at the site of the point of maximum tenderness, and the release of pus relieved the symptoms and led to rapid cure.

In addition to the signs and symptoms recorded in the account of the first case the following are recorded in the other histories.

"The pain may be of moderate character with periodic exacerbations of excruciating almost intolerable pain and periods of perhaps several months of freedom from all symptoms. The adjacent joint may exhibit periodic effusions after exercise which are absorbed with rest. One patient had tubercles in the lungs and vomicae, dead bone in the ribs, and other complications. In one case a sinus communicated with the knee-joint. The new periosteal bone is softer than normal."

Such is the account that Brodie gave to a condition which he recognized as a distinct entity. The accuracy of his observations and the value of the method of treatment which he devised were such that though thousands of such cases have been treated by hundreds of different surgeons since his time little if any addition has been contributed to the clinical picture.

What Constitutes a Chronic Abscess?

Brodie's clinical work was unaided, or, as some would express it, unhampered, by bacteriology or radiology. It is therefore impossible to define accurately in the present terms of bacteriology or radiology the nature of the lesion which he described. Though in the light of our present knowledge most of his eight cases appear to be of a staphylococcal nature, the history of at least one of them suggests the possibility of a tuberculous infection. Consequently, though we may reasonably refer to any chronic abscess of bone as a Brodie's abscess, it would be preferable, from the point of treatment, to confine the term to abscesses that are due to the staphylococcus. These constitute all but a relatively small minority that are attributable to the tubercle bacillus, typhoid bacillus, pneumococcus, streptococcus, etc. We are also faced with the question of what constitutes a chronic abscess? Brodie stated that the duration of symptoms in his eight patients varied from five to eighteen years. To-day, with the ready accessibility of the harmless and painless radio-

* Paper read at the Annual Meeting of the British Association of Radiologists at Belfast, May 13, 1938.

T. N. MACGREGOR: DYSFUNCTIONAL UTERINE BLEEDING TREATED WITH PROGESTERONE

FIG. 1.—Case 9; aged 21. Menarche at 16; Menstruation 3/28 regular for four years. Since then periods have been irregular. Endometrium before progesterone therapy. (Magnification $\times 50$.)

FIG. 2.—Case 9. Endometrium before progesterone therapy. (Magnification $\times 260$.)

FIG. 3.—Case 9. Endometrium after 25 mg. progesterone. (Magnification $\times 50$.)

FIG. 4.—Case 9. Endometrium after 25 mg. progesterone. (Magnification $\times 260$.)

T. N. MACGREGOR: DYSFUNCTIONAL UTERINE BLEEDING TREATED WITH PROGESTERONE

FIG. 5.—Case 10; aged 25. Menarche at 13; Menses 3-4/28 for three years; 2 to 3 months' amenorrhoea followed by profuse loss. Endometrium before progesterone therapy. (Magnification $\times 50$.)

FIG. 6.—Case 10. Endometrium before progesterone therapy. (Magnification $\times 260$.)

FIG. 7.—Case 10. Endometrium after 25 mg. progesterone. (Magnification $\times 50$.)

FIG. 8.—Case 10. Endometrium after 25 mg. progesterone. (Magnification $\times 260$.)