BRITISH MEDICAL JOURNAL

LONDON SATURDAY NOVEMBER 12 1938

HEPARIN AND THROMBOSIS*

BY

CHARLES H. BEST, M.A., M.D., D.Sc., F.R.S.

Professor of Physiology, University of Toronto

(WITH SPECIAL PLATE)

While a search of the earlier literature reveals the fact that several investigators realized that certain mammalian tissues contain one or more anticoagulants, an active fraction was first isolated in Howell's laboratory in 1916 (McLean, 1916; Howell and Holt, 1918). Howell named the substance "heparin" (*hepar*—the liver), and he and his collaborators have studied its distribution, chemical properties, and physiological significance. For some time heparin was prepared exclusively from the liver tissue of the dog, and it was very difficult to secure sufficient amounts of a product satisfactory for physiological experiments.

In 1928 Dr. McHenry and I, desiring to use heparin in our histaminase work, found it possible to prepare active fractions from ox liver, and subsequently Charles and Scott (1933a) showed that this substance was present in varying amounts in a great many tissues of this species—lung, liver, and skeletal muscle—giving, in this order, the best yield.

Action of Heparin

Under certain conditions, as Howell originally showed, heparin acts as an antiprothrombase, but in other more physiological circumstances it is apparently an antithrombase. The finding of Professor John Mellanby (1934) that the action of a very potent thrombase added to plasma can be inhibited by heparin has been confirmed in my laboratory by Mustard and Jaques (1937). In these circumstances heparin is apparently an antithrombase. Heparin can, of course, be titrated against thrombokinas with considerable accuracy under appropriate conditions, so that thrombokinas may be termed an antiheparin or, conversely, heparin an antikinase. The product described by Charles, Fisher, and Scott (1934) and tentatively termed "antiheparin" turned out to be a typical thrombokinas (Ridout and Best; see Charles, Fisher, and Scott, 1934).

Recent Work on Heparin

The work on heparin in Toronto was started in the department of physiology in 1929. At that time only crude material was easily available, and any advance along physiological or therapeutic lines appeared impossible until a much more highly purified preparation was at hand. Furthermore, although it was known that heparin was an anticoagulant, its action on thrombus formation had been explored only to a slight extent. In one paper Shionoya (1927a), working at the Mayo Clinic, stated that an adequate dose of heparin did not prevent the formation of white thrombi, and this is certainly true when crude heparin is used in rabbits. He subsequently reported

(1927b) that magnesium sulphate markedly retards thrombus formation in the heparinized rabbit.

PURIFICATION OF HEPARIN

Studies in purification were begun at my suggestion by Dr. Arthur Charles, a young organic chemist, in the department of physiology. Charles was presently transferred to the physiological and biochemical section of the Connaught Laboratories, where he had the privilege of working with Dr. D. A. Scott, whose studies on the crystallization of insulin and the preparation of protamine-zinc-insulin are well known to many. Charles and Scott (1933b, 1934, 1936) soon attained success in the purification of heparin, and before very long they evolved a process for the preparation of adequate amounts of satisfactory material from ox lung. Finally, a crystalline barium salt of uniform potency was obtained. An amorphous benzidine compound was also secured which gave the same empirical formula ($C_{25}H_{65}O_{56}N_2S_5$). I shall refer to this compound later.

The crystalline product contains nitrogen, a part of it at least in the form of NH_2 groupings, upon the inactivation of which the potency of heparin is lost. (Howell thought that his purified product was free of nitrogen.) Jorpes (1935), using purified heparin made by the Toronto procedure, noted the presence of sulphuric acid groups in the molecules, and there is some evidence that the loss of this grouping also may interfere with the activity. Tests for carbohydrate are positive but do not indicate the presence of glycuronic acid, which some previous observers had thought was present. An adequate discussion of the chemistry of heparin should, of course, include detailed references to the interesting and very important contributions made by Schmitz and Fischer, Jorpes, Chargaff, and others, but I must proceed with more physiological matters.

Charles has shown that when the crystalline barium salt is treated with excess of ammonium carbonate solution at $65^\circ C$. the barium may be completely removed. The heparin is then precipitated by acetic acid, centrifuged out, washed with ether, and dried. It can be stored as a dry powder. When required a solution can be made from this; tricresol may be used, and the solution sterilized by passing through a Berkefeld or Zeitz filter.

STANDARDIZATION OF HEPARIN

This brings me to a necessary digression—a consideration of the standard of heparin. We have suggested quite recently (Charles; see Murray and Best, 1938) that the material from the crystalline barium salt may be used as a standard. In a very informal discussion with Sir Henry Dale we agreed that it might be wise to make the standard available, if need should arise, as a solution.

* A lecture delivered at University College, London, on June 14, 1938.

I will not discuss here the various physiological methods for the standardization of heparin. They depend on the inhibition of the clotting of blood under certain conditions. A unit has never been defined exactly. It is usually stated to be the amount of anticoagulant which inhibits for twenty-four hours the clotting of 1 c.cm. of blood under certain conditions of temperature and so forth. We have suggested as a provisional unit the activity contained in 0.01 mg. (10 γ) of the barium-free material—that is, 100 units per mg. This would make a convenient unit from both the physiological and the clinical aspect.

A standard of heparin would not only be advantageous from the point of view of those interested in its action, either experimentally or clinically, but would also be of great service to anyone working on any aspect of blood

as a result of subcutaneous administration. In the rat the effect of one subcutaneous dose may be to send up the clotting time to over twenty-four hours, the return to normal occupying from twelve to twenty-four hours or longer (Fig. 2). However, two out of sixty experimental animals died as a result of haemorrhage into their subcutaneous tissues after the administration of heparin. There was not a great deal of haemorrhage, but 2 or 3 c.cm. is a large amount of blood for a rat. It is possible that the damage produced locally would not be sufficient to cause a dangerous or even a serious amount of bleeding in large animals.

When heparin is precipitated with protamine a very insoluble compound is formed which cannot readily be suspended (Best, Charles, and Cowan, 1937). With benzidine, however, a compound can be prepared which is absorbed quite slowly and gives a clotting time of ten

FIG. 1.—Intravenous injection of heparin. (From Jaques, Charles, and Best.)

clotting. We can standardize thrombase or thrombokinase against the standard heparin, and thus help to bring order out of a very confused field. More standards—for thrombase and thrombokinase—are urgently needed.

ADMINISTRATION OF HEPARIN

If a single dose of heparin be given intravenously the clotting time becomes prolonged. The increase in the clotting time depends on the size of the dose (Fig. 1). There is no negative phase—that is, the clotting time does not go below normal after a massive dose of heparin: it comes back fairly accurately to the initial value. Heparin can be given subcutaneously as well as intravenously. The crude material first used was not well absorbed, but purified heparin is absorbed quite rapidly from the tissue spaces, and a good effect can be obtained

FIG. 2.—Subcutaneous injection of heparin. (From Jaques, Charles, and Best.)

FIG. 3.—Subcutaneous administration of benzidine-heparin. (From Jaques, Charles, and Best.)

to fifteen minutes for quite long periods (Jaques, Charles, and Best, 1938) (Fig. 3). It was found advisable to have a little unmodified heparin in the mixture, as certain doses of benzidine-heparin given alone will have no effect at all, but if given with the unmodified heparin the immediate and the prolonged action are both forthcoming.

The best procedure in the administration of heparin is to give a small dose intravenously and to follow this with a constant intravenous injection. The clotting time can be set at any chosen level and maintained for long periods. In experimental animals we commonly give 40 units per kg. as the initial dose, and follow this with a continuous injection of 30 units per kg. an hour. With this dosage the clotting time is usually maintained at from twenty to thirty minutes. Somewhat smaller doses are used by Dr. Murray and his colleagues in the department of surgery.

Studies in Relation to Thrombi

We have carried out a number of studies on heparin in relation to the formation of thrombi. In the first series Murray, Jaques, Perrett, and I (1937) wished to determine the effect upon the mixed thrombus produced when the internal surface of veins is injured. This injury was produced in the first series of experiments by mechanical means, such as clamping the vessel repeatedly with strong forceps. In the second series chemical means were used. A section of the vein was isolated and sodium ricinoleate was injected into the lumen. It was allowed to remain for a few minutes and the blood was then permitted to flow again. After both these procedures a thrombus forms in a large percentage of the cases. Heparin was administered continuously to unanaesthetized dogs for seventy-two hours after the injury had been produced, and healing of the surface of the veins was found to take place in this time with no tendency for thrombi to occur subsequently. The fact that in the best series of experiments (see Table)

Exp. No.	Control			Heparin			Heparin u./lb./hr.	Hrs. Heparin given	
	Occluded	Patent	% Occluded	Occluded	Patent	% Occluded			
68	1	1							0.15 c.cm. Soricin injected in all veins. <i>Control</i> 17% patent <i>Heparin</i> 100% patent
69	2	-							
30	4	-							
72	2	-		-	2		10.3	68	
73	-	-		-	4		10.9	73	
74	1	1		-	2		10.5	72	
75	-	-		-	4		10.0	72	
Total	10	2	83%	-	12	-			

(From Murray, Jaques, Perrett, and Best.)

all the heparinized veins were open means that they were found in this state a week, usually, after the injury. Heparin, however, had been given for only three days.

STUDIES ON THE FORMATION OF WHITE THROMBI

These results were encouraging, but we wished to determine the effect of heparin on the formation of white thrombi. It is well known, of course, that the white thrombus is the nucleus from which the mixed thrombus grows. In certain conditions a thrombus is obtained which consists entirely of platelets, and it was interesting to determine the effect of heparin on this process. To do this the procedure evolved by Rowntree and Shionoya (1927) of the Mayo Clinic was followed. A cellophane shunt between the femoral artery and the femoral vein was made and the blood was allowed to flow for a certain time. Then the cellophane tube can be sectioned to determine whether any thrombi have formed (Fig. A on Plate). This was carried out on quite a large series of animals—monkeys, dogs, cats, and rabbits (Best, Cowan, and MacLean, 1938). In the rabbit it was difficult to observe the effect; although the tube never became occluded, small specks of thrombi could be seen in spite of the presence of heparin. There was no doubt, however, that in the case of the monkeys, dogs, and cats the action of heparin was to prevent the formation of the white thrombi. These, of course, are admittedly artificial conditions: the blood flowing over the glass and cellophane tubing favours the formation of thrombi, and one

cannot conclude from the dose of heparin required in these cases how much would be needed to prevent a similar phenomenon inside the blood vessel.

HEPARIN AND CORONARY THROMBOSIS

Another line of approach to this problem is the study, in which Dr. D. Y. Solandt and I (1938) have spent a large part of our time this year, of the formation of thrombi which may be produced in the coronary artery of dogs. It is possible by dissecting out the left coronary artery in the dog to obtain a length of half an inch or so, to clip off all the branches, and put two clamps on the main stem a quarter of an inch apart. Then, by introducing sodium ricinoleate into the lumen of the vessel and leaving it for five minutes enough irritation is produced to cause thrombi to form. In a comparable series of animals in which this procedure was carried out and the animals were heparinized just before the sodium ricinoleate was injected a thrombus occurred in only one case. In other words, with this isolated exception, heparin prevented the formation of thrombosis of the coronary artery in the same way as it did in the peripheral veins. We were led to study the coronary artery because of the relatively high frequency of thrombosis of this vessel in the human species and also because it is apparently much easier to get a thrombus in the coronary artery in dogs than in some of the other arteries.

My collaborators and I have thus carried out four investigations which may be listed as follows: mechanical injury to veins, chemical injury to veins, formation of white thrombi, and the studies on the coronary artery. In each case an effect of heparin in preventing thrombosis could be demonstrated.

[Professor Best then projected a coloured film, the first part of which showed the flow of blood through a glass cell introduced between the carotid artery and the jugular vein of the monkey. The rapid formation of white thrombi was well illustrated, and the breaking off of large masses to form emboli. The growth of the thrombus downstream could be clearly seen. The thrombi tended to form in the periphery of the cell, where the flow is slowest, but also on a scratch in the axial stream where the flow was very rapid. The action of heparin in preventing the formation of white thrombi was demonstrated. In the second part of the film the passage of a plasma containing platelets but very few other cells through a small glass perfusion chamber was shown. The clumpings of the platelets could be clearly seen, together with the formation of the digitations. (Figs. B and C on Plate.)]

Use of Heparin in Physiological Experiments

I should like now to speak on the use of heparin in physiological experiments. It is obvious that there are a great number of things which one can do if adequate amounts of the purified material are available. For example, my colleague Dr. D. W. G. Murray has studied, among other problems, end-to-end anastomoses of blood vessels. This can, of course, be done without heparin, but the percentage of successful results is much greater when heparin is used.

Another illustration of the use of heparin is in so-called "exchange" transfusion in experimental animals. This subject has recently been studied by Thalhimer, Solandt, and myself (1938). In one experiment, for example, a nephrectomized dog in which the blood urea had become very high was joined by means of a pump, which enabled exchange to be made in each direction of the same amount

of blood, to a normal donor. The result was a precipitate fall of blood urea in the nephrectomized animal and a rise in the normal donor (Fig. 4). (One can bring the blood urea of the nephrectomized animal very quickly back to the normal limits if several donors are used, because the kidneys of the normal dogs do double duty and excrete urine for both.) Both animals were, of course, heparinized. I do not suppose that these findings have any obvious application to clinical medicine, but those of us who are medically trained have visualized very exceptional circumstances in which a completely anuric child might be joined for a few hours to a parent, both being

FIG. 4.—The fall in urea of a nephrectomized animal during "exchange" transfusion. (From Thalheimer, Solandt, and Best.)

heparinized. The clinician, however, who would have to take the responsibility in a situation of this kind might have different ideas on the subject.

Neutralization of Heparin by Protamine

Some years ago it was shown that protamine is an antithrombase (Waldschmidt-Leitz, Stadler, and Steigerwaldt, 1929). If protamine be added to the blood the clotting time goes up because of the antithrombase action of the substance introduced. About eighteen months ago Charles, Cowan, and I (Best, Charles, and Cowan, 1937) published a paper in which we showed that heparin and protamine formed a very insoluble compound which could not easily be resuspended. More recently it has been shown by Chargaff and Olson (1937) that if heparin be given to an animal and be followed by protamine the effect of the heparin may be completely neutralized and the clotting time brought back to normal with great rapidity. This has been confirmed by Jaques and myself. Many results can now be brought forward in confirmation of the fact that protamine can be used to neutralize the effect of heparin *in vivo* and *in vitro* (Fig. 5).

Physiological Significance of Heparin

The first piece of evidence in favour of the view that heparin plays a physiological part is obtainable from a study of its distribution: the work of Howell, and subsequently that in Toronto and elsewhere, has shown its presence in various tissues, including blood. Again, the recent work by Jorpes, Holmgren, and Wilander (1937)

(see also Holmgren and Wilander, 1937) along histological as well as chemical lines has shown its presence in particularly large amounts in the mast cells of Ehrlich, and has suggested the possibility that these cells are responsible for its production. A third point has been brought out by Waters, Markowitz, and Jaques (1938) in my department—namely, that there is a good deal of evidence

FIG. 5.—Neutralization of heparin by protamine. (From Jaques, Charles, and Best.)

that the increased clotting time seen in anaphylactic shock in the dog is due to the liberation of heparin. This has been shown also to hold good for peptone shock.

It has been known for some time that histamine, while it may account for many of the signs of anaphylaxis, has little or no effect upon the coagulability of the blood (Dale and Laidlaw, 1911). When a dog goes into anaphylactic shock there is a very great rise in clotting time—that is, from four to five minutes to perhaps forty-eight hours. This was originally shown by Biedl and Kraus (1909). Samples of the blood can be taken and treated with protamine, and the heparin equivalent estimated by this procedure. The results of experiments of this kind carried out by Jaques and Waters are shown in Fig. 6. These results show that heparin in a concentration of approximately 1.5 units per c.cm. has appeared in the blood of the animal in anaphylactic or peptone shock. This amount of heparin is sufficient to raise the clotting time of the blood from the normal value to such extremely high levels—sixty to seventy hours—that it may be termed incoagulable. When an animal was shocked after hepatectomy, Waters, Markowitz, and Jaques found little or no rise in the clotting time of the blood or in the heparin equivalent. Furthermore, these workers have obtained from the blood of the shocked animal much more heparin than can be detected in the blood of the normal animal, while the liver of the shocked animal contains less heparin than normal liver. The physiological and chemical results provide practically conclusive evidence that the anticoagulant isolated from the blood of the shocked animal is heparin, and it may be possible to prepare this material

as the crystalline barium salt. It appears, therefore, that in anaphylactic shock in dogs histamine, heparin, and possibly other substances are liberated.

FIG. 6.—Liberation of heparin in anaphylactic and peptone shock in dogs. (From Waters, Markowitz, and Jaques.)

We may conclude that several points of evidence now suggest the possibility that heparin is a substance of physiological importance.

The Possibility of Clinical Application

As soon as more highly purified preparations of heparin became available investigations were begun in the department of surgery under the supervision of my colleague Dr. D. W. G. Murray. It was found that none of the preparations available except that which had been through the stage of the crystalline barium salt could be given with complete safety. The fact that this highly purified material can safely be given to human subjects over long periods is well established by the findings on some 350 patients, most of whom had experienced a major operation before heparinization was started (Murray and Best). The intravenous administration of heparin is never begun earlier than two or three hours after the completion of a major operation—a splenectomy, for example—and is continued until the patient is able to move actively about in bed. This may be for three or four days, or it may extend to weeks.

It is of great interest that workers in Sweden (Crafoord, 1937; Hedenius, 1937; Holmin and Ploman, 1938; Magnusson, 1938), using heparin prepared by the method elaborated in Toronto, have initiated clinical studies on the prevention of thrombus formation.

It is hopeless to attempt to secure information regarding the effect of heparin in the prevention of thrombus formation by the indiscriminate heparinization of post-operative cases or by the heparinization of isolated cases. This has not been attempted in Toronto. Murray and McKenzie are selecting, in so far as is possible, those cases in which the hospital statistics show that the inci-

dence of post-operative embolism is relatively high. I suppose the best type of case to study would be those exhibiting that rather rare condition known as phlebitis migrans. If it were possible to collect a group of these cases in one ward and to have them thoroughly studied before and after the administration of heparin a great deal might be learned.

A few words may be said with regard to the practical application of the work on coronary arteries in dogs. If the clinical cardiologist knew when thrombosis was about to occur—that is, if there were definite premonitory signs—he might by the appropriate use of heparin secure a short reprieve, perhaps even a long one, for some of his patients. The material is readily available, and can apparently be given safely to hospitalized patients; but our lack of knowledge, or perhaps the complete absence of premonitory signs, makes it impossible to conduct a clinical investigation along these lines. Heparin might be given immediately after one attack of coronary thrombosis, but clinicians do not speak with a single voice when discussing the spread of thrombosis from the original focus or the significance of intramural thrombi formed as a result of coronary thrombosis. If a clinical investigation of cardiac cases should be initiated, the necessity for studying very large numbers and of heparinizing only alternate cases is obvious.

One of the best methods of determining the clinical role of such a substance as heparin is to push ahead with more studies along physiological and experimental pathological lines in the hope that the clinical applications will become obvious. It is true, of course, that many of the experimental conditions which have been studied are not closely related to those observed in the clinic, but the results have been very stimulating, and one hopes that investigations along these lines will be pursued.

I should like to thank Professor Lovatt Evans for his invitation to lecture on this subject in his department.

REFERENCES

- Best, C. H., Charles, A. F., and Cowan, C. (1937). *Amer. J. Physiol.*, **119**, 272 (Proc.).
 — Cowan, C., and MacLean, D. L. (1938). *J. Physiol.*, **92**, 20.
 Biedl, A., and Kraus, R. (1909). *Wien. klin. Wschr.*, **22**, 363.
 Chargaft, E., and Olson, K. B. (1937). *J. biol. Chem.*, **122**, 153.
 Charles, A. F., Fisher, A. M., and Scott, D. A. (1934). *Trans. roy. Soc. Canada*, **28**, 49.
 — and Scott, D. A. (1933a). *J. biol. Chem.*, **102**, 425.
 — (1933b). *Ibid.*, 431, 437.
 — (1934). *Trans. roy. Soc. Canada*, **28**, 55.
 — (1936). *Biochem. J.*, **30**, 1927.
 Crafoord, C. (1937). *Acta chir. scand.*, **79**, 407.
 Dale, H. H., and Laidlaw, P. P. (1911). *J. Physiol.*, **43**, 182.
 Hedenius, P. (1937). *Lancet*, **2**, 1186.
 Holmgren, H., and Wilander, O. (1937). *Z. mikr.-anat. Forsch.*, **42**, 242.
 Holmin, N., and Ploman, K. G. (1938). *Lancet*, **1**, 664.
 Howell, W. H., and Holt, E. (1918). *Amer. J. Physiol.*, **47**, 328.
 Jaques, L. B., Charles, A. F., and Best, C. H. (1938). *Acta med. scand.*, in press.
 Jorpes, E. (1935). *Naturwissenschaften*, **23**, 196.
 — Holmgren, H., and Wilander, O. (1937). *Z. mikr.-anat. Forsch.*, **42**, 279.
 McLean, J. (1916). *Amer. J. Physiol.*, **41**, 250.
 Magnusson, J. H. (1938). *Lancet*, **1**, 666.
 Mellanby, J. (1934). *Proc. roy. Soc. B*, **116**, 1.
 Murray, D. W. G., and Best, C. H. (1938). *J. Amer. med. Ass.*, **110**, 118.
 — Jaques, L. B., Perrett, T. S., and Best, C. H. (1937). *Surgery*, **2**, 163.
 Mustard, R. A., and Jaques, L. B. (1937). *Trans. roy. Soc. Canada*, **28**, Appendix B, CLVI.
 Rowntree, L. G., and Shionoya, T. (1927). *J. exp. Med.*, **46**, 7.
 Shionoya, T. (1927a). *Ibid.*, 19.
 — (1927b). *Ibid.*, 963.
 Solandt, D. Y., and Best, C. H. (1938). *Lancet*, **2**, 130.
 Thalheimer, W., Solandt, D. Y., and Best, C. H. (1938). *Lancet*, in press.
 Waldschmidt-Leitz, E., Stadler, P., and Steigerwaldt, F. (1929). *Hoppe-Seyl. Z.*, **183**, 39.
 Waters, E. T., Markowitz, J., and Jaques, L. B. (1938). *Science*, **87**, 582.

CHARLES H. BEST: HEPARIN AND THROMBOSIS

FIG. A.—White thrombi forming inside a cellophane tube

FIG. B.—Illustrating growth of white thrombi downstream, from above downwards and from left to right. (From Best, Cowan, and MacLean.)

I

II

III

IV

FIG. C.—I and II: Single platelets, showing digitations. III: The clumping of central refractile areas of platelets. IV: Polymorphonuclear leucocyte. (From Best, Cowan, and MacLean.)

A. EISDELL MOORE: TREATMENT OF DELAYED UNION OF CARPAL SCAPHOID

FIG. 1.—Radiograph showing ununited fracture of right scaphoid, with some vacuoles.

FIG. 2.—Application of leather-covered aluminium "cock-up" splint; dorsiflexed part fitted into centre of palm; arm portion held on flexor surface of forearm by leather casing.

FIG. 3.—Radiograph taken at the end of six months, showing apparent union.