BRITISH MEDICAL JOURNAL

LONDON SATURDAY DECEMBER 3 1938

THE SEPTIC HAND *

BY THE LATE

SIR DAVID WILKIE, M.D., Ch.M., F.R.C.S.

Professor of Surgery, University of Edinburgh

A discussion on the septic hand cannot fail to be instructive, because, in the first place, it makes us clarify and bring into focus our views on acute infections and the principles on which they should be treated, and, in the second, it brings into prominence an oft-neglected but all too common problem of everyday practice.

There has been an unfortunate tendency to regard infections of the hand as among the minor surgical maladies. In point of fact such infections are of primary importance, for not only do they not infrequently endanger life but quite commonly they lead to lasting disability in men who have to earn their livelihood by manual work. Rational treatment of such infections will depend (a) on a sound knowledge of the infective process, and (b) on a familiarity with the anatomical structure of the hand and the relation of this to varied types of infection. We owe a great deal to two men for our modern treatment of hand infections—to Bier for his illuminating studies of the infective process, and to Kanavel for his contribution to the pathological anatomy of the infected hand.

The Infective Process

Mild infections of the hand are of hourly occurrence, although but a very small proportion of them ever reach the doctor. There is obviously a considerable local and general immunity against the stray bacteria found on our hands and on the varied objects which may inflict an accidental wound. Every now and then, however, a germ of some virulence gains access, or a wound with devitalized edges presents a soil in which otherwise harmless germs may flourish, and we recognize that infection has gained a hold. The signs of such a foothold are local pain and swelling, and the general evidence of poisonous absorption, such as rigor, rise of temperature and pulse, and dryness of the tongue and mouth.

In the mildest degree of infection the local defence mechanism appears to be easily capable of dealing with the invading germs. The transient blush, which means that the local vascular and cellular reactions have quickly overcome the invaders and restored the part to normal, has a lesson for us in the treatment of infections. Local hyperaemia permits a rapid concentration of defences and

confines the fight to a little zone around the point of entrance of the germ. If we could restrict the area of infection to the tissues round about the point of entry, if we could have the fight fought to a finish in the outskirts and preserve intact the inner defences and the citadel, we should have an almost painless victory with the minimum of loss.

Bier's great contribution to the study of infection was his demonstration that, by controlling the return circulation of blood and lymph from an infected part and producing a slight lymph stasis in the area, we may not only limit the central spread of infection until such time as the process of immunity is fully under way, but we may by bathing the infected area in lymph dilute the poisons, reducing pain and favouring recovery with a minimum of tissue damage by suppuration or necrosis. In this country his message was understood imperfectly, and "Bier's congestion" was taken to imply a venous stasis induced by elastic constriction verging on the degree

FIG. 1.—Bier's bandage applied to the upper arm. (The bandage is applied direct to the skin and secured by a small bit of adhesive strapping. There is no blueness or swelling of the forearm.)

at which capillary haemorrhage occurs. While Bier freely acknowledged that he got the idea from Thomas's method of treating delayed union of fractured bones and that he applied the short period of intense venous stasis in the treatment of tuberculous joints, he maintained that the method of using long-continued mild blood and lymph stasis in acute infections was a later evolution and a discovery of which he had reason to be proud.

If this discussion does nothing else than induce you to use mild passive hyperaemia in acute infections it will

* Read in opening a discussion in the Section of Surgery at the Annual Meeting of the British Medical Association, Plymouth, 1938.

not have been in vain. I consider that a Bier's elastic bandage is the most valuable piece of apparatus a practitioner can possess, with the obvious proviso that he knows how to use it.

Besides this principle of seeking to localize infection whilst immunity is rising, two others immediately suggest themselves. Can we do anything by way of the blood stream (a) to kill off the infecting bacteria; (b) to raise artificially the patient's immunity to their toxin? The era of effective chemotherapy would seem to have been inaugurated, and the discovery of the sulphonamide group of drugs has opened a new and hopeful chapter in the treatment of virulent infections. The use of antitoxic sera, although somewhat overshadowed during the past two years by chemotherapeutic measures, has now an imposing array of successes to its credit, and in severe infections of known character it should, if available, always be tried. Such measures as inducing free excretion by kidneys, skin, and bowels by copious libations of bland fluid, the replenishing of the liver by giving

pulp infection and the need for early opening when the infection gets a hold. We know the anatomy of the spread of pus in an acute paronychia, and the simple but important means of draining it effectively.

Acute Streptococcal Infection with Lymphangitis

When from a punctured wound or from a scratch virulent streptococci gain entrance to the tissues we may see within a few hours the evidence of their multiplication and their spread. The hot throbbing wound, the ache up the arm, the rigor followed by a sharp rise of temperature, indicate a severe infection. On examination there may be local redness and some oedema of a finger, some swelling of the dorsum of the hand, faint red lines up the forearm, and some tenderness in the axilla, all indicating a virulent and rapidly spreading infection. How is such infection best combated? The patient is put to bed and given copious diluent drinks. A warm moist dressing is applied to the hand and forearm and a few turns of a thin elastic bandage are wound *lightly*

FIG. 2.—Transverse section of upper arm. a = subcutaneous vein with accompanying lymphatics.

glucose in the drinks, favouring metabolism and sleep by sunlight and fresh air, all tend to aid the more specific forms of therapy.

The Pathological Anatomy

Kanavel, by his intensive study and analysis of hand infections and his differentiation of types and the anatomical approach in dealing with each, gave a new impetus and interest to the treatment. As in so many other regions of pathology, it is by the study of early stages that we observe the essential nature and the lines of spread of different infections. In the full-blown massive and oedematous infected hand it is oftentimes difficult to detect the true locus of the underlying infection. We have learned, however, to recognize the signs of tendon-sheath infection, of infections in the various palmar spaces, and of those in the deep forearm space. We now know the anatomical restrictions of the finger-

FIG. 3.—Section of arm with Bier's bandage applied. Note compression of superficial vein and lymphatics (a); the vessels beneath the deep fascia are compressed little if at all.

round the upper arm (Fig. 1). The bandage is applied direct to the skin in such a way that a gentle pressure obstructs the superficial veins and lymphatics and, without producing any blueness or duskiness, causes a mild lymph stasis in the limb (Figs. 2 and 3). The secret of successful application of the bandage is the abolition of pain. If pain is increased the bandage is too tight and must be reapplied. The elastic bandage may be left on continuously for forty-eight hours with advantage.

Recognizing from the physical signs the streptococcal nature of the infection, it is well to give without delay 30 c.cm. of streptococcal antitoxic serum after desensitization. At the same time treatment with one of the sulphonamide preparations is instituted. A careful watch is kept for the development and localization of pus in the hand and forearm, but no great haste is made to interfere locally unless there be signs of a finger-pulp or tendon-sheath infection.

The effect of lymph stasis in localizing the infection is in my experience remarkable, and skill in the application of Bier's elastic bandage is a great asset to all who have to treat such cases. The decision as to when a studied conservatism should give place to active surgical intervention requires nicety of judgment, but as a general rule it may be said that delayed is wiser than premature incision.

Finger-pulp Infection

Owing to its peculiar anatomical structure the finger pulp demands special attention. The dense fatty layer which forms the pulp is broken up into compartments by fibrous septa, which further bind the skin down to the bone. Any infection, with oedema and effusion, produces great tension within the pulp. The terminations of the digital arteries, including the branch to the distal phalanx, traverse the pulp and are exposed to the tension within it. This may be so great as to compromise the blood supply to the phalanx, and necrosis results.

It is obvious, therefore, that special anatomical conditions here determine early intervention. If within thirty-six hours of the onset of pain there is no sign of amelioration an incision made on one side of the distal phalanx is carried well down into the pulp, opening up its spaces. A counter-opening on the other side of the phalanx is sometimes desirable. Even after opening the finger it is wise to continue with the fomentations and the passive hyperaemia.

Tendon-sheath Infections

Spread of infection to a tendon sheath is heralded by severe pain and interference with the function of the finger. This is held in a position of semiflexion. Any effort to straighten it causes acute pain. Definite localized tenderness is met with on pressing over the proximal end of the tendon sheath in the case of the index, middle, and ring fingers, and over the line of the sheath in the case of the thumb and little finger.

As soon as infection of a sheath is diagnosed a general anaesthetic should be given, a tourniquet applied to the arm, and an incision made along one side of the proximal phalanx of the affected finger. The bulging sheath is incised and, if purulent fluid escapes, further incisions may with advantage be made along the side of the second phalanx and in the palm. In the case of the thumb and little finger incisions must be made along the thenar and hypothenar eminences and at the wrist to drain the radial and ulnar bursae respectively. Hot saline baths and active movement of the fingers should be encouraged, as they favour drainage. Tubes and rubber drains are better avoided.

Palmar Space Infections

The infection of one or other of the palmar spaces, so clearly defined by Kanavel, may arise from an infected wound of the palm, by lymph spread from a finger, or by rupture of an infected tendon sheath. When due to the last-named the clinical features may be confusing, but the treatment is seldom in doubt. The spaces, being deeply situated, may contain pus under tension without obvious fluctuation being detectable. The dense palmar fascia offers an unyielding barrier to forward spread. The metacarpals and the interosseus muscles bar the way behind and pus is forced to travel along the lumbricals to the web between the fingers, where it bulges, points, and,

if not opened, bursts. The drainage of these spaces, either from the web back along the lumbricals or by incising the palm—preferably the former—will, as a rule, lead to rapid improvement if no tendon sheath has suffered.

Forearm Infection

In some cases, and particularly those of neglected tendon-sheath or palmar space infections, swelling and oedema of the lower part of the forearm may suggest a deep-seated cellulitis. Suppuration in Parona's space—that is, beneath the flexor muscles and tendons and in front of the pronator quadratus—is diagnosed. When oedema and redness involve the lower half of the forearm and considerable tenderness is elicited over the front of its lower third and above the carpal bursa, then one may rest assured that pus is collecting in the deep forearm space. Without delay this should be opened under a general anaesthetic with a free incision, laying open the abscess on one side and a counter-opening for a drain on the other.

Amputation of Fingers

With severe tendon-sheath infection, even after reasonably early opening, it is no uncommon thing to find one or both flexor tendons sloughed. Not infrequently along with this the first interphalangeal joint becomes involved, with softened ligaments, shedding of articular cartilage, grating, and lateral mobility. We should realize that a stiff or tender finger may be a crippling handicap to the labouring man. It is always a keen disappointment at the end of many weeks of painful dressings for him to find that his trouble and patience have been in vain and that amputation is necessary. Sloughing of tendons, if accompanied as it so often is by disorganization of a joint, should point to amputation without undue delay. Gangrene of the distal part of the finger, met with in severe infections, is another indication for amputation. Amputation should never be performed when the inflammation is at its height. A little delay is beneficial, and helps to avoid any flare-up or central spread of infection.

Indolent Infections

A not uncommon type of infection, especially among field workers, is one associated with brawny induration, little tendency to pus formation, and slight general reaction. The usual signs of tendon-sheath or palmar space infection are difficult to determine, and indeed may never develop. In such cases short-wave therapy has been found of material value, and, if available, should always be employed. In some cases the infection may slowly resolve without suppuration but with a considerable period of disability; in others pus may eventually form in the palm or in the forearm and require evacuation by incision. Complete recovery from such low-grade infections may take many months, but eventually may be complete.

Rehabilitation in Hand Infections

The acute phase of infection commands our interest and attention: all too often the slow tedious phase of convalescence receives but scanty care. Almost without exception infections of the hand and forearm are followed by fibrosis with its attendant stiffness. The more protracted the suppuration the greater will be the resultant disability from cicatricial fibrous tissue. When pus has formed and has localized, generous incisions allowing of

the freest drainage will, by hastening resolution, diminish the subsequent fibrous contractures.

It is very desirable that those who are responsible for the treatment during the acute inflammatory phases should also supervise the subsequent treatment by physiotherapeutic means. While few would challenge the value of complete rest to the part during the early acute phase of inflammatory reaction, few would deny the benefits of active movement and of exercise once the florid spreading stage is over and the early dangers are past. Apart from the sloughing of tendons and the involvement of joints which give permanent stiffness, we have the adhesions in tendon sheaths, the fibrosis in the fascial planes between the muscles, the fibrous plaques which form where tissue was destroyed. If the thought of function be kept uppermost in our minds in conducting the treatment in the stages which follow that of the invasive process, we shall never permit the development of the unbreakable barriers to movement which determine crippledness in the neglected case.

The use of active and purposive exercise in restoring useful movement should govern the tedious period which follows that of nursing care. Massage and passive exercise have their place, but active movement, controlled and guided by the patient's will, is the effective element in treatment. If interest and achievement lend their spur to effort, the patient, progressing from one task to another designed to increase the range of certain movements, passes unconsciously from invalidism back to effective work. The workshop now forms an essential part of the equipment of the convalescent institution. Our responsibility for treatment does not end until the maximum of usefulness has been restored.

Summary

1. Acute infections of the hand should be treated in the first instance on conservative lines.
2. In very severe infections three measures may be employed: (a) lymph and venous stasis (Bier); (b) chemotherapy (sulphonamide); (c) passive immunity (antitoxic serum).
3. In certain areas—for example, finger pulp and tendon sheaths—anatomical considerations make early opening desirable.
4. As soon as the early dangers are past, efforts to restore function should begin, and they should end only when the maximum return of function that is possible has been attained.

At the eleventh British Congress of Obstetrics and Gynaecology, which, as already announced in these columns, is to be held at Edinburgh on April 4, 5, and 6, 1939, the chief subject for discussion will be pain in labour and methods of alleviation, to be introduced by Professor Chassar Moir of Oxford and Dr. John Sturrock of Edinburgh. The official guests of the Congress will be Professor Wagner (Berlin) and Dr. Samuel Meaker (Boston). Other communications will deal with genital hypoplasia, dysmenorrhoea, tubal infection, hyperemesis gravidarum, results of pregnancy toxæmia, trichomonas infection, male sterility, habitual abortion, and myometrial activity. A visit will be paid to the Department of Animal Genetics under the direction of Professor F. A. E. Crew. There will be a demonstration of pathological specimens, and those who desire to exhibit pathological material are asked to communicate with the local secretary of the Congress, Dr. Chalmers Fahmy, 7, Chester Street, Edinburgh, 3, from whom other information concerning the Congress may be obtained.

THE CORRECT NAME OF THE MALIGNANT TERTIAN MALARIA PARASITE

BY

SIR RICKARD CHRISTOPHERS, C.I.E., F.R.S.,
Colonel I.M.S. (ret.)

*Leverhulme Fellow under the Medical Research Council,
London School of Hygiene and Tropical Medicine*

AND

J. A. SINTON, M.D., D.Sc.,
Lieut.-Colonel I.M.S. (ret.) *

*Manson Fellow, London School of Hygiene and Tropical
Medicine*

Not only is the malignant tertian malaria parasite referred to under a multiplicity of colloquial names, such as malignant tertian, subtertian, aestivo-autumnal, and "tropica," but even the zoological name of this species is a constant source of difficulty and doubt to those who have occasion to use it. Thus the name now in most general use is *Plasmodium falciparum*, but many authors employ the names *P. praecox* or *P. immaculatum*, claiming that one or other of these is the correct specific designation. Uncertainty also exists as to whether this species should or should not be placed in the genus *Laverania*, and *L. malariae* and *L. falcipara* are names not uncommonly seen. Efforts to apply the rules of zoological nomenclature seem only to have added to the confusion.

The Present Position

Hitherto controversy has chiefly revolved around the question which of the specific names—*praecox*, *immaculatum*, or *falciparum*—has the strongest claim to validity. Before any detailed discussion is entered into it is desirable first to review very briefly the present position in respect to these names. Grassi and Feletti (1890b) gave the name *Haemamoeba praecox* to a parasite found by them in the blood of birds from a certain malarious region in Italy, and also, later in the same paper, to the then newly discovered asexual stages of the parasite of aestivo-autumnal fever in man, believing the two parasites to be identical. Later Grassi and Feletti (1891, 1892) gave the name *relicta* to the same or a similar parasite in the blood of birds. Sergeant *et al.* (1929) consider that, although these authors gave the same name to the avian and the human parasite in the first place, they have by the subsequent naming of *relicta* automatically designated *praecox* as the human parasite, and that therefore the correct name for the latter species is *praecox*. This would be in accordance with a common procedure in zoological nomenclature—that is, where a name at first given has included two closely related species, one of which is subsequently designated by another name, the original name becomes applicable to the remaining species. Other authors (Schaudinn, 1903; Blanchard, 1905; Stiles, 1929) hold that the context in the original description shows that the name *praecox* was definitely given to the bird parasite, and that it must therefore be invalid for the human one, whatever subsequent procedure on the part of the authors may have been. In this case, if *praecox* be invalid for the human parasite, then some other name is required for it. Here again authors differ. Some hold with Schaudinn (1903) that the first valid name is *immaculatum*, used by Grassi and Feletti (1892); others that this name is also invalid and that the proper one for the parasite is *falciparum*, given by Welch (1897). It is pointed out by Blanchard (1905)

* The junior author desires to thank the Royal Society of London and the London School of Hygiene and Tropical Medicine for the financial assistance which has enabled him to continue his work on malaria.