the calcareous mass attached to the under surface of the liver would appear to have been the remains of the primary cyst. In Case II the primary cyst was not located. In neither case was a history suggestive of anaphylaxis obtained, so that the time of rupture of the primary cyst cannot be known; but it is likely that in each case multiple cysts had been present and unsuspected in the peritoneal cavity for several years.

In Case I the first manifestation of hydatid disease appears to have been in the earlier months of the pregnancy, when the patient began to feel unwell. Presumably in consequence of an examination laparotomy was performed and a hydatid cyst removed from the anterior part of the uterus. In Case II hydatid disease was not suspected until the occurrence of an obstructed labour due to a cyst being impacted in the pouch of Douglas. It is noteworthy that in both cases, although hydatid disease had probably been present for many years, it was not discovered until the fourth decade of life, and then only on account of an associated pregnancy.

Literature

References in the literature to the ubiquitous hydatid cyst in relation to pregnancy and parturition are few. Franta (1902) collected all the recorded cases and dealt with them in three papers. The first was concerned with dystocia due to hydatid cysts in the pelvis, and recorded thirty-six cases. The second contained an account of twenty-two cases in which operation for removal of the cyst was performed in pregnancy. The third dealt with the influence of pregnancy, labour, and the puerperium on hydatid cysts. I am aware of only four recorded instances in this country. Gemmell (1899) described a case of hydatid cyst in the omentum obstructing labour which was treated by abdominal section and removal in the puerperium. Blacker (1908) published a case of hydatid cysts complicating pregnancy in which Porro-Caesarean section was performed. Andrews (1908) also described a case in which pregnancy was complicated by the presence of a hydatid cyst in the pelvis. The remaining case is that of Bell (1925), whose patient had an obstructed labour necessitating craniotomy, the cyst subsequently being removed during a second pregnancy.

Summary

1. Two cases of obstructed labour due to a hydatid cyst are described because of their comparative rarity.
2. Both cases were examples of secondary echinococcosis with multiple cysts in the peritoneal cavity.
3. In neither case was there a history of anaphylactic manifestations such as frequently accompany rupture of the primary cyst.
4. Although hydatid disease had probably been present for several years, it had not been suspected. In the first case it was discovered at the fifth month of pregnancy, and in the second during labour.

I wish to express my thanks to Sir Ewen Maclean and Dr. M. D. Arwyn Evans for permission to publish these cases.

REFERENCES

- Andrews, H. R. (1908). *J. Obstet. Gynaec. Brit. Emp.*, **14**, 333.
 Bell, B. (1925). *Ibid.*, n.s., **32**, 114.
 Blacker, G. (1908). *Ibid.*, **14**, 336.
 Dew, H. (1930-1). *Brit. J. Surg.*, **18**, 275.
 Franta, J. (1902). *Ann. Gynéc. Obstét.*, **57**, 166, 296, 420; **58**, 401.
 Gemmell, J. E. (1899). *British Medical Journal*, **2**, 1603.

EPIDEMIC MYALGIA FIVE CASES IN ONE HOUSEHOLD

BY

C. R. G. HOWARD, M.B., Ch.B.

Department of Medicine, University of Bristol

The following account of an outbreak of epidemic myalgia may be of some value, as it provides a possible differential diagnosis in cases of appendicitis, pleurisy, cholecystitis, and muscular rheumatism.

Survey of Literature

Daae (1874) in Norway is first credited with describing the disease, which, he says, is distinguished by pain in the chest and back, febrile disturbance, and relapses. Gradually it became recognized as a clinical entity, and by 1888 the name "devil's grip" had been associated with it in America. From this time it attracted little attention until 1930, when Sylvest (1930, 1931) gave a full account of an epidemic which occurred on the island of Bornholm, in which 10 per cent. of the population were attacked. He described the onset as sudden, with pain in the hypogastrium and nausea, the pain shifting to the chest and shoulders after a few hours. These symptoms are associated with fever, from 99° to 105° F., and the attack lasts a few days. Relapses are frequent. Sylvest noticed a seasonal incidence in summer and early autumn. The sexes are equally affected, and, although no age is exempt, most cases occur between the ages of 5 and 15. There appeared to be an incubation period of two to four days, and examination of the blood showed an eosinophilia.

Since 1930 there have been a few reports in the literature. Rector (1935) records nineteen cases in which the patient was under 8 years old, and he suggests the use of quinine sulphate in cases that do not clear up quickly. Smith (1937) gives an account of an epidemic at Rugby School, in which there were myalgic pains, chiefly in the back of the neck.

In 1934 the disease received more attention owing to several outbreaks in American towns. MacDonald, Howell, and Cooper (1937) investigated an outbreak in Cincinnati; clinically they found the same picture as did Sylvest, but there were no patients over 15 years of age. They point out that constipation was frequent and relapses after a few days were common features. These workers found no eosinophilia, but some of the patients had a leucopenia. Bacteriological investigations of the cerebrospinal fluid were negative, as also were blood cultures. The Wassermann reaction was negative in all cases. Cooper and Keller (1937) tried to find a specific organism in washings of the throat and nose, but the only constant growth was a small colony type of *Streptococcus viridans*, to which they attached little importance. Later work with animals suggested a filterable virus as the cause, with a possible incubation period of eight to ten days. F. Wolter (1937), investigating a series of outbreaks in Germany, adds the fact that the urine is usually very dark in colour, and this, he suggests, is due to the presence of creatinine, which he attributes to the changes in the muscles. He discusses the connexion between epidemic myalgia and epidemic poliomyelitis, and points out that there is a similar seasonal and climatic incidence.

Case Records

Case 1.—A man aged 26 first complained of epigastric pain, made worse by meals. The following afternoon the pain became very acute and was localized to the right costal margin; there were no clinical signs of pleurisy and a radiograph of the chest was normal. During the next two days the pain in the chest became much less severe, but it was made worse by deep breathing or laughing. Tenesmus was present. On the fifth day there was a relapse, with headache, sweating, and a temperature of 101° F. After 30 grains of aspirin had been taken there was a profuse perspiration followed by considerable improvement. During the next three days the pain in the lower part of the chest became less severe and was more generalized over the back and shoulders.

Case 2.—A woman aged 27 complained of vague abdominal pain with nausea; the next morning the abdominal pain was worse and tenesmus was pronounced; in the afternoon there was a sudden onset of pain in the right costal margin, aggravated by coughing or laughing. On the third day she felt better, but had a relapse in the afternoon, and complained of feeling hot and exhausted. The temperature was not taken. During the next three days the pain became less, and finally disappeared after a treatment with an infra-red lamp.

These patients did not go to bed.

Case 3.—A woman aged 26 complained of a sudden onset of pain in the abdomen and general malaise eight days after exposure to Cases 1 and 2. Her temperature was 103° F., and she went to bed. The next day the pain was localized to the right costal margin and was made worse by coughing or deep breathing. The morning temperature was 101° F., and the evening 102° F. A blood film was taken which proved normal. Tenesmus was present. On the third day she felt better; her temperature was normal and she got up. The following day there was a mild relapse, the patient feeling hot and sweating, but she managed to undertake a railway journey.

Case 4.—A baby aged 15 months had a general upset, being off her food and appearing to have pain in the abdomen, especially when crying; she was sick twice. The following day she was well but fretful. On the third day she had a relapse, the temperature being 102° F., but there were no localizing symptoms. During the next few days the child was off its food but was otherwise well. The motions were normal.

Case 5.—A female aged 17, nurse to Case 4, developed abdominal pain, with general malaise and a temperature of 99° F., ten days after Case 4 had recovered. The next day the pain was localized to the right side of the chest, where it persisted for two or three days. There was no relapse. This was a very mild attack, if one at all.

Commentary

These cases thus closely followed the clinical picture described by Sylvest, the main features being a sudden onset with abdominal pain and malaise, the pain moving to the chest, particularly the right costal margin (Cases 1, 2, and 3), on the second day, later diffusing all over the chest and shoulder. Pyrexia varied from 99° to 103° F., and tenesmus was present in three cases. A relapse on the third, fourth, or fifth day was noticed in four cases. Cases 3 and 5 appeared to have incubation periods of eight and ten days respectively.

The chief disabilities are intermittent attacks of malaise for about a week, combined with the inability to laugh or take a deep breath without excruciating pain in the chest, also the frustration, in spite of a strong urge, of being unable to defaecate satisfactorily.

It is interesting to note that at about the same time as the above cases occurred there were admitted to the local hospital more cases than usual, perhaps, of appendicitis that never came to operation. Unfortunately these were not investigated from the point of view of myalgia.

Summary

A brief review of the literature of epidemic myalgia is presented, together with an account of five cases occurring in one household.

REFERENCES

- Cooper, M. L., and Keller, H. M. (1937). *Amer. J. Dis. Child.*, **54**, 231.
 Daac (1874). Ref. in *J. Amer. med. Ass.*, 1934, **102**, 460.
 MacDonald, R. R., Howell, B., and Cooper, M. L. (1937). *Amer. J. Dis. Child.*, **53**, 1425.
 Rector, J. M. (1935). *Ibid.*, **50**, 1095.
 Smith, R. E. (1937). *Clin. J.*, **66**, 330.
 Sylvest, E. (1930). *Ugeskr. Laeg.*, **92**, 798, 982.
 — (1931). *Ibid.*, **93**, 1155.
 Wolter, F. (1937). *Klin. Wschr.*, **16**, 1087.

Clinical Memoranda

Domiciliary Emergency Treatment of Eclampsia

There is no gainsaying the fact that domiciliary treatment of persons acutely ill of obstetrical complications is the correct one, not only in the homes of the well-to-do but also in the overcrowded homes of the poor. Hospitals have their place in the treatment of disease, but a review of the table below proves beyond doubt that the emergency treatment of eclampsia at home is pre-eminently the best line.

Comparative Statement of Cases of Eclampsia Treated from 1933 to 1937 in Hospital and by Domiciliary Emergency Measures

	1933	1934	1935	1936	1937		Totals
Total admissions to hospital ..	1,416	1,649	1,759	1,729	1,780		8,333
Cases of eclampsia occurring in hospital or admitted after onset ..	13	13	17	26	17	Incidence 1.03% Average death rate 11.6%	86
Deaths ..	2	2	1	5	0		10
Emergency team: Cases treated in own home	4	10	13	5	11		43
Deaths ..	0	0*	0	0	0		0
Hospital and home cases combined ..	17	23	30	31	28		129
Deaths ..	2	2	1	5	0	Average death rate 7.7%	10

* Of the patients attended in their own homes during 1934 one was dead before the arrival of the emergency team.

As our experience increases the pitfalls become apparent: one that I have observed is where there is such a pronounced elevation of the blood pressure in a case of pre-eclamptic toxæmia that even the excitement attending transport and admission to hospital causes the patient to develop an eclamptic seizure, the prognosis of which it is difficult to forecast.

During the past five years 8,333 cases have been admitted to this institution, and eighty-six have suffered from eclampsia either before or after admission, with a mortality rate of 11.6 per cent.; whereas during the same period forty-three cases were treated in their own