A thickened ligamentum flavum may so constrict the dura as to produce pressure on one or more roots of the cauda equina. Spurling has recently drawn attention to this condition. Can it be due to localized extradural haemorrhage producing fibrosis and thickening of the ligament? Lipiodol examination usually reveals a constricted hour-glass type of filling defect on the posterior and lateral aspects of the spinal canal. (Fig. 6, Special Plate.)

Results

A total of eighty-three cases of lumbar intervertebral disk ruptures producing sciatic pain have been operated on at the Massachusetts General Hospital up to June, 1938. The immediate complete post-operative relief of the sciatic pain is commented upon by practically every patient.

The operation is of considerable magnitude, and should be undertaken only by competent surgeons after a thorough trial of conservative methods or after demonstration of the presence of a dangerous amount of pressure on two or more nerve roots. There have been two deaths in our series; both of these patients had severe irreparable nerve damage from large ruptured disks. There are three known failures. The rest of the patients are either entirely well or are very markedly improved.

Summary

Rupture or herniation of lumbar intervertebral disk tissue into the spinal canal causing pressure on one or more roots of the cauda equina is a well-established clinico-pathological entity, and is the cause of intractable sciatica in many cases. The lesion can be accurately localized by radiological examination after injection of iodized oil into the subarachnoid space. Laminectomy and operative removal of the extruded disk fragment give prompt and complete relief of symptoms in most of the cases.

BIBLIOGRAPHY

- Barr, J. S. (1937). *J. Bone Jt. Surg.*, **19**, 323.
 — Hampton, A. O., and Mixter, W. J. (1937). *J. Amer. med. Ass.*, **109**, 1265.
 Hampton, A. O., and Robinson, J. M. (1936). *Amer. J. Roentgenol.*, **36**, 782.
 Kocher, Theodor (1896). *Mitt. Grenzgeb. Med. Chir.*, **1**, 415.
 Love, J. Grafton, and Camp, John D. (1937). *J. Bone Jt. Surg.*, **19**, 776.
 Middleton, G. S., and Teacher, J. H. (1911). *Glasg. med. J.*, **76**, 1.
 Mixter, W. J. (1937). *Ann. Surg.*, **106**, 777.
 Schmorl, Georg (1927). *Verh. dtsch. orthop. Ges.*, **21**, 3.
 — (1927). *Verh. dtsch. path. Ges.*, **22**, 250.
 — (1929). *Fortschr. Röntgenstr.*, **40**, 629.
 — (1931). *Arch. Orthop. Mech.-Ther.*, **29**, 389.
 Spurling, R. G., Mayfield, F. H., and Rogers, J. B. (1937). *J. Amer. med. Ass.*, **109**, 928.

ORTHOPAEDIC ASPECTS OF SCIATICA*

BY

W. A. COCHRANE, F.R.C.S.Ed.

Lecturer in Orthopaedic Surgery, University of Edinburgh

In cases of sciatica a careful eliminative diagnosis leaves a group which is essentially orthopaedic. The physical examination is very important, and represents skilled interpretation of an aggregation of indirect signs which, taken together and supported by an understanding of the local anatomy and bodily statics of the part, can often point to a definite orthopaedic conclusion in the individual case and determine the proper treatment. This is the more likely to be the case if the examiner is able to interpret properly the evidence of stereoscopic films of the low back and pelvic joints.

Vulnerability of Fifth Lumbar Nerve

Orthopaedists try to explain sciatica largely on the basis of referred pain from the sacro-iliac and lumbo-sacral joints and their supporting ligaments. There is also the consideration which Putti has given to the funicular portion of the fifth lumbar nerve as it lies in the foramen. Situated as it is in this fibro-osseous canal, the fifth root, surrounded by a thick venous plexus, practically fills it. The walls of the canal are bounded posteriorly by the posterior spinal articulation, antero-internally by the intervertebral disk, and postero-externally by the lumbo-sacral and ilio-lumbar ligaments. The dorsal primary division of the nerve turns backwards to supply the sacrospinalis muscle, and in its course runs close to the lateral aspect of the posterior articulation, supplying the joint capsule itself. These relations make it possible to assert that the fifth lumbar nerve is in a position of peculiar vulnerability, as it may be affected by the action of trauma of the low back on the lumbo-sacral articulations and by inflammation of them also. It may also be affected by prolapse of the disk or by change in shape and size of the canal secondary to displacement of the fifth lumbar vertebra on the sacrum or to degenerative changes of the intervertebral disk. The slight margin of safety present in the asymptomatic individual by reason, I consider, of the competency of the normal muscular support is further jeopardized by anatomical variations of the lumbo-sacral region, which either cause increased motion at the expense of stability or, by reason of their asymmetry, produce abnormal stresses. In this connexion it is of interest to note that Steindler found anatomic variations in the lumbo-sacral region in 37 per cent. of a large group of cases of sciatica referred to the orthopaedic service.

Circumstances Leading to Sciatica

It may be submitted, then, that many cases of sciatica have their explanation in the operation of a chain of circumstances consisting of: (1) the special anatomic form of the patient, who is either of the unduly heavy or unduly slender build; (2) functional decompensation of the supporting structures of the low back, mainly muscular, often added to by advancing age, unsuitable occupation, increasing weight, debility, and a toxic factor, and leading to faulty posture with exaggeration of the normal lumbo-sacral angle. Such faulty and imperfect use, combined with degenerative changes in the disks and the presence of

G. Bertoni and B. Melli (*Rif. med.*, 1938, **54**, 1526), who record twelve illustrative cases in patients aged from 12 to 35, state that Donaggio's reaction is performed as follows. If a certain quantity of ammonium molybdate is mixed with an equal quantity of normal filtered urine or cerebrospinal fluid, boiled, filtered again, and a watery solution of a basic aniline dye, such as thionin, is added, the colouring solution is completely precipitated by the molybdate and the fluid above the precipitate soon becomes quite clear. In some conditions, however, especially typhoid fever, the urine and spinal fluid acquire the property of checking or entirely preventing the precipitation of the thionin. This phenomenon has been shown by Donaggio to occur when the fluids in question contain a larger amount of organic colloid substances than normally. In typhoid fever the reaction becomes positive at an early stage and remains so throughout the course of the infection.

* Read in a discussion at a joint meeting of the Sections of Neurology and Psychological Medicine, and Orthopaedics and Fractures at the Annual Meeting of the British Medical Association, Plymouth, 1938.

anatomical variations in the lumbo-sacral level, lays the subject open to repeated minor traumata affecting the articulations and their supporting ligaments. The trauma is often the precipitating factor, operating upon an area ill prepared to receive it. It is often a trauma which, if the individual had been a physically well-educated person posturally, would never, by reason of the functional and positional reserve of the parts concerned, have produced evil effects. No part of the body can be used wrongly without the rest of the body suffering, and, conversely, if one regards the problem only as a local one without reference to the re-establishment of correct use of the body as a whole, it is not easy to effect a permanent and satisfactory cure.

Restoration of Good Posture

In cases of sciatica, therefore, that are adjudged to be lumbo-sacral in type, regard has usually to be paid to the element of muscular fatigue associated with prolonged misuse, leading to an abnormal alignment of pelvis and lumbar spine. The low back becomes almost spastically fixed in this position, usually of increased lordosis in the subacute and chronic cases. While manipulation may assist such a case, it is often necessary and better to eliminate gravity, weight-bearing, and motion for a time by a period of several weeks of thorough rest in bed on a firm mattress. The patient lies on his back with a pillow beneath the head and knees. Thrice daily, for half an hour, the dorsal spine is thrown forward by placing a firm pillow beneath it, thus leaving the lumbar spine unsupported. Thrice daily he is rolled over, to lie face downwards on a pillow placed lengthwise beneath the trunk, and three hot fomentations, at ten-minute intervals, are placed on the low back. Weight and pulley traction of 10 lb. may materially assist in relieving the more acute symptoms. In about two weeks the muscles of the back lose their spasticity, and one may then make a posterior plaster-of-Paris shell for the trunk to lie in as a cradle. The time is then ripe for giving spinal massage, and instruction to the patient, in the lying-down position, in how to gain active conscious voluntary control of the muscles of the trunk. He is taught how actively to flatten the lumbar spine, to contract the side-abdominal muscles, and to breathe correctly, using the diaphragm with the chest elevated. He is fitted to a light back-brace which spans the lumbar curve when he gets up, and is taught how to maintain an improved posture, with a lengthened-out waist, when he stands, sits, and walks.

It needs an intelligent co-operative patient to do all this, but if the surgeon knows his business and applies the instruction himself there is usually no difficulty. The brace is discarded in due course, and the final test of recovery is the restoration of habitually good posture. One would emphasize the lack of understanding that exists in the country to-day as to what a trained orthopaedist should be able to do in the proper correlation of the structure and function of the body so far as the locomotory system is concerned. He is there to establish right use wherever there is wrong use; he must be trained, and able to apply in his practice his knowledge of the proper order in which re-establishment of correct use must be made. He must carry it out himself in the re-education of his cases, and not leave it to the physiotherapist. All phases of the treatment must be applied in their correct order. Rest without proper instruction in muscle balance later is of little value, as is the application of support while painful protective spasm of muscles is still present. Indiscriminate massage and exercises, often done

at the wrong time, are all to be avoided as entities in themselves.

In no aspect of his work may the art of the orthopaedist be exercised to greater advantage than in the case of sciatica, occurring either apparently alone or in conjunction with low back pain. It is a specialty within a specialty, a field in itself.

In sprains of the sacro-iliac joint producing sciatica manipulation followed by firm protective strapping may have to be employed, but in these cases also it is not unusual for the muscles to be poor and the posture bad. Consequently they may need thorough rest, postural re-education, and support for a time after the manipulation has done its work.

Semi-sacralization of Fifth Lumbar Vertebra

In these cases of painful semi-sacralization the treatment detailed above will often succeed. In aggravated cases the effect of really prolonged rest in a well-fitted plaster-of-Paris spica has not infrequently been successful in my experience. It may be combined with traction of about 10 lb. to the affected leg. I have had patients who remain well after ten years and more following this method. There are cases of the sort which do require operation. Frequently they have had severe sciatica associated with painful affections of both the accessory lumbo-sacral articulation and the pelvic joint on the same side. The operation of spinal fusion is usually best in these cases, but one should not be in a hurry to operate on them. Far from it: if spinal fusion is to be performed, above all it must be certain that the fusion is done with the lumbo-sacral angle in the normal position and not in extension. I have had successful results by fusing the accessory articulation by a lateral Smith-Petersen approach, as for sacro-iliac joint arthrodesis, taking out the block of bone at a higher level, opposite the accessory joint. I consider this to be the least formidable type of operation for the patient. If required, the sacro-iliac joint could be fused at the same time.

Cases of Rheumatic Type

In regard to the cases which are of true rheumatic type accompanied by rheumatic fibrositis and myositis it is possible in certain cases to effect a cure by making an incision along the line of the posterior portion of the iliac crest, to detach the origin of the muscle and fascia there, and slide them down to a lower level. Heyman has rightly pointed out that such cases may closely simulate those of sacro-iliac pain, with referred neuralgic pain in the leg. Ober's operation of division of the contracted ilio-tibial band high up at its origin is also in this category.

Conclusion

Dr. Harris has rightly pointed out the importance of not neglecting to perform a rectal examination in all cases of difficulty in the diagnosis of sciatica. I well recall a case of very severe sciatica that had eluded diagnosis in which the cause was revealed forthwith by a rectal examination; it was due to an enormous abscess in the pelvis consequent upon the recrudescence of an old healed osteomyelitis of the iliac bone, the result of a bullet wound sustained fifteen years previously.

The fact that sciatica is rare in childhood and adolescence is of decided significance in explaining its aetiology on a basis of a strong muscular reserve of support to what is, after all, potentially rather a weak

part of the body—that is, the lumbo-sacral region of the spine. The penalties of the upright position are many, and this is one of them.

I must not close these remarks without paying a tribute to the Boston school of orthopaedic surgeons for the work they have done on this subject. It is due to them, and chiefly to Goldthwait, that a proper system of examination has been established and that these cases, so far as the orthopaedic aspect of them is concerned, are now capable of scientific analysis, classification, and treatment.

THE XENOPUS PREGNANCY TEST

BY

EDWARD R. ELKAN, M.D.

(WITH SPECIAL PLATE)

The discovery of what is now known as the xenopus pregnancy test is based on experiments conducted by Hogben (1930, 1931), who observed that hypophysectomy produced ovarian retrogression, and the injection of anterior pituitary extracts ovulation, in the female South African clawed toad. Further experiments on these lines were carried out by Charles, Slome, and Zwarenstein in Capetown, and later by Bellerby, Zwarenstein, and Shapiro in London, and it is due to the collaboration of these authors that the initial difficulties which stood in the way of a standardization of the xenopus test were overcome. This is particularly true with regard to Bellerby's work on the laboratory conditions necessary for sustained reproductive activity, which work made it possible to improve greatly the reliability of xenopus as a test animal. While there has so far been no difficulty in obtaining regular supplies of these toads from South Africa our next aim must nevertheless be the breeding of xenopus in this country. Bles attempted this as early as 1905, and sporadic successes have been recorded by other authors (Bles, 1906; Bellerby, 1933; Vanderplanck, 1935; Elkan, 1938). It is to be expected that in the near future we shall have not only a standardized xenopus pregnancy test but also a standardized method of breeding this animal.

Weisman (1938) recently reviewed the tests for the detection of early pregnancy which have been devised since Aschheim and Zondek's discovery eleven years ago. Apart from the Aschheim-Zondek reaction only Friedman's technique has become widely popular; the xenopus test, though in some points superior to both the Aschheim-Zondek and the Friedman techniques, has remained more or less unnoticed despite the enthusiastic reports which have appeared in various journals in recent years. These three pregnancy tests are the only ones to deserve, in Weisman's opinion, the term "excellent." In the course of the last twelve months I have done 295 xenopus tests on 2,112 toads, and I fully endorse Weisman's judgment. Some of the objections to this test raised by earlier observers seem to have become invalid, and as the early detection of pregnancy remains important from the psychological and the gynaecological points of view a method which allows the diagnosis to be made within a few hours should be welcome.

Biological Considerations

Xenopus laevis Daud., a toad of the genus *Aglossa*, is fairly common in all the tropical parts of Africa (Fig. 1,

Special Plate). The toad is exported from Capetown, where the animal dealers seem to have no difficulty in catching as many as are required. My own experience with these exporters does not confirm fears expressed by Crew (1937) and Weisman (1938) as to the availability of the animal. So far supplies seem to be unlimited and export unrestricted, and since the animals survive the passage from Capetown to London I see no reason why they should not be shipped to any part of the country. I have at present tadpoles in my laboratory bred in an open-air tank this summer. It remains to be seen whether these tadpoles can be brought up in sufficient numbers and whether their growth is rapid enough to make the breeding of xenopus an economic proposition. My present tadpoles look very much like young fish. They stand on their heads most of the time, and feed on a mixed diet of infusoria, particularly flagellates, and raw liver emulsion, of which a little is poured into the aquarium every day.

The adult clawed toads received from Capetown vary greatly in size, the smallest ones measuring 2 inches, the largest 4½ inches, from mouth to anus. I find those of medium size most suitable for tests. The small toads are mature, but do not stand the injections so well; the larger specimens tolerate the injections well, but they require very big jars for observation. It is easy to distinguish the males from the females. They are identical in colour, but the external opening of the cloaca in the female has three labia—two dorsal and one ventral—which do not, however, form a receptaculum for semen; these labia are absent in the male (Fig. 2, Special Plate). Fully grown males do not reach the same size as fully grown females. The females also acquire a characteristic shape through the bulging lungs being pushed aside by the ovaries, which protrude on both sides of the abdomen like cushions.

The animals are easy to feed. They will accept daphnia, earthworms, newts, tadpoles (*Rana esculenta*), minced meat, or minced liver, but flatly refuse to touch bread or potatoes. It is a little difficult to say authoritatively how often they should be fed. They will certainly accept food every day, but they never show any signs of being particularly hungry, and if not fed for a week they do not seem to be the worse for it. If they grow at all in the laboratory they do so extremely slowly, even if they are well fed. It is wiser not to feed those specimens which are to be used for tests in a day or two. The animals have a short alimentary canal. They eat whenever they happen to come across anything edible, but do not seem to make much use of what they have eaten. The water becomes dirty from their excreta soon after they have been fed, and must then be changed. It does not seem to matter—at least so far as the tests are concerned—whether feeding takes place once, twice, or three times a week if only the animals are given as much food as they will accept at each feeding time.

During the cold season all the animals are kept at the laboratory in a specially constructed tank, and at a temperature of 23° to 26° C. In the summer only the animals actually needed for tests are kept in the laboratory. The others, particularly those who are having their "resting time," are kept in outdoor tanks, where they become comparatively tame; when it rains they can be seen leaving the water and climbing on to stones provided for them. Hundreds of xenopus can be kept in a comparatively small space, and since the females of this species do not make any noise their presence is not a nuisance to the neighbourhood.