Poly-methylmethacrylate soft-tissue spacers to maintain tissue planes during two-stage revision knee arthroplasty

GN GILLESPIE, J MUTIMER, AJ PORTEOUS
Avon Orthopaedic Centre, Southmead Hospital, Bristol, UK

CORRESPONDENCE TO

GN Gillespie, Specialist Registrar in Orthopaedics, Avon Orthopaedic Centre, Southmead Hospital, Southmead Road, Westbury-on-Trip, Bristol BS10 5NB, UK

Two-stage revision surgery is considered to be the reference standard for deep infections following total knee arthroplasty. Even with the use of preformed mobile poly-methylmethacrylate (PMMA) spacers at first stage revision, the scarring seen at second stage frequently obliterates the suprapatellar pouch and the gutters. Re-forming these planes prior to re-implantation is time-consuming.

Following debridement and after fitting the mobile spacers we mould spacers from antibiotic-loaded PMMA cement and place these in the suprapatellar pouch and in the gutters (Fig. 1). At second stage, it is found that these planes are well preserved, facilitating the procedure.

Figure 1A Anteroposterior (A) radiograph taken after first stage revision demonstrating the PMMA implants and soft-tissue spacers.

Figure 1B Lateral radiograph taken after first stage revision demonstrating the PMMA implants and soft-tissue spacers.

Use of a Rampley's sponge holding forceps to facilitate stapled haemorrhoidectomy

MT CARTMELL, AR DIXON
Department of Colorectal Surgery, Frenchay Hospital, Bristol, UK

CORRESPONDENCE TO

MT Cartmell, Specialist Registrar in Colorectal Surgery, Frenchay Hospital, Bristol BS16 1LE, UK
T: +44 (0)117 918 6503; M: +44 (0)798 912 8975;
E: markcartmell@hotmail.com

In stapled haemorrhoidectomy, a circular anal dilator is inserted and the purse-string suture anoscope then used to reduce three-quarters of the circumference of prolapsing mucosa whilst the suture is inserted. Maintaining the position of the purse-string suture anoscope whilst inserting the suture can be difficult, often requiring an unsighted assistant in a difficult position. We have found use of a Rampley's sponge holding forceps to hold the purse-string suture anoscope on the circular anal dilator (Fig. 1) frees the surgeon's hand or obviates the need for an assistant. In addition, it can be torqued and angulated to facilitated suture insertion.

Figure 1 Rampley's sponge holding forceps holding the purse-string suture anoscope on the circular anal dilator.

Removal of impacted deflated Foley catheter from male urethra

UGOCHUKU OKONKWO

West London Elective Orthopaedic Centre, Epsom General Hospital, Epsom, Surrey, UK

CORRESPONDENCE TO

Ugochuku Okonkwo, West London Elective Orthopaedic Centre, Denbies Wing, Epsom General Hospital, Dorking Road, Epsom, Surrey KT18 7EG, UK
E: ugookonkwo@btinternet.com

After deflating the balloon of a Foley catheter, it can become impacted at the urethral meatus, making complete removal difficult for the clinician and painful for the patient.

Remove the needle from an intravenous cannula leaving the plastic sheath. Discard the needle responsibly. Plug a syringe of lubricating anaesthetic gel into the plastic sheath. Pass the plastic sheath into the urethra alongside the Foley catheter. Inject the gel into the urethra then withdraw the plastic sheath. After the gel has taken effect, apply gentle traction to the Foley catheter.