bandage is put over all, wide enough and tight enough to effect all that is required. The time for removing the sponge and under-covering of linen or cotton may vary from twenty-four to forty-eight hours (never longer), and the removal must be effected so carefully as not to cause any dragging on the flaps. If the linen or cotton be found to adhere to any point of the wound, I prefer leaving it where it is, quite dry, to bathing or tearing it off, and a snip with the scissors is all that is needed for this. The linen is replaced by a clean bit, if all be found favourable, and the sponge is washed, wrung dry, and put on again, and left for two or three days, or even more, in order to make sure that the union is quite firm. The patient is kept in bed, and not allowed to move freely until the wound is healed. For three or four days, very meagre diet is allowed, and no stimulants in ordinary cases.

Ligatures, even of waxed silk, do not generally produce irritation, as will be shown in the following case, which I give in support of what I have said.

Three weeks ago, assisted by Dr. McWatt, I amputated the left breast for cancer. The patient was very stout and the gland large and embedded in fat, a considerable amount of which was removed after the tumour, because of infiltration and in order to admit of adjustment of the flaps. Seven waxed silk ligatures were applied, the edges were brought together with carbolised gut, and the wound was dressed in the way described above. On the second morning, while removing the dressing for the first time, and just as I was finishing, the patient suddenly started aside, and thus caused the forcible separation of the linen from the axillary angle of the wound to which it was adhering. About a teaspoonful of blood followed, and I feared that mischief might be the result of this irritation. Next day, when I called, I found she had had a rigor after I left, and, on removing the dressing, I observed a faint blush over the inner half of the wound. I reapplied the dressing, and, in anticipation of erysipelas, prescribed twenty grains of sulpho-carbolate of soda every six hours. Although there was a faint blush over part of the wound, Dr. McWatt, who was present when I removed the dressing, thought it looked like healing by first intention; and everything about the case, at this stage, certainly afforded more reason for saying that it *had* done so than was found in that reported by Mr. Maunder at the same period after operation. Two days later, the linen cloth next the wound was damp, and, on removing the dressing and carefully examining the cicatrix, I found it healed and firm, save for about the third of an inch. To prevent reopening, I supported the flaps on each side of this unclosed spot with strips of adhesive plaster. A little thin pus continued to pass for a few days, but gave no trouble. Five ligatures have come away through the firmly closed edges without a drop of pus accompanying or following any of these, and the remaining two are clean and dry, though gentle traction is applied to each daily. The patient's pulse never rose above 90, and she has never had the slightest pain, except when she gave the start when I was removing the first dressing.

This is not a selected case, but I give it simply because it is the last I have had. I may further remark that, in minor amputations, I invariably place a bit of sponge right over the stump, and have hitherto found it of much service.

CURE OF AN "ULCERATED LEG" OF FULLY THIRTEEN YEARS' STANDING.

By JOHN COCHRANE, L.R.C.P.Ed., and L.R.C.S.Ed.,
Parochial Medical Officer, Colmonell, Ayrshire, N.B.

THE following case, the details of which I now present, is a somewhat interesting one. When I commenced practice here (Colmonell), in April last, I was called to see a male pauper who had suffered for a long time from an ulcerated leg. On proceeding to the patient's house, I discovered the following state of matters. The leg affected was the right one; and about four inches above the ankle, on the outside of the limb, there was an ulcer, measuring five inches in length and three inches in breadth. The smell from it was something most horrible, and, on examining it more closely, I found the wound gangrenous. The limb, from the knee down to the ankle, was very much swollen and the skin dark-coloured in appearance. The foot also was much swollen and the ankle oedematous. The ulcer also discharged continually much watery fluid. Its edges were exceedingly hard and the surface of a slaty-blue colour. Its depth was fully an inch, and conjoined with these signs was much pain in the affected limb. The pain at times was so severe as to prevent the man from getting sleep at night. The history of the case, obtained from the man and his wife, was as follows. Fully thirteen years ago, the limb was attacked with an inflammatory form of

disease; the symptoms, as described to me, warranting the supposition that it was erysipelas, the result of an injury to the leg. Since then, numerous little ulcers had appeared at different parts of the limb, and had disappeared under treatment, such as it was. Then a large ulcer formed on the anterior aspect of the limb, and, healing at one spot, spread over the limb till it settled in the part which was now affected. At the same time, the leg began to swell and become dropsical. For some years, the man continued to work at his employment, which was a laborious one; and then, when he found the leg becoming worse, he applied for relief from the parochial board. When I saw him for the first time, his leg was as hard as a stone from the knee to the ankle. He said that this hardness had been present for a considerable period, and the limb now was almost three times the circumference of the neighbouring leg and much heavier. There were a few varicose dilatations of the veins near the knee; but what struck me most was this same *hardness* of the tissues and the blackened appearance of the skin. The skin also, from the knee downwards, was affected with psoriasis. This latter, no doubt, was due to the continual irritation from the nasty discharges coming from the ulcerated surface, and from the venous obstruction, which also undoubtedly was the cause of the oedema of the ankle and foot. Possibly, also, the condition of the blood was at fault, owing to the neglect which had been shown to the treatment of the ulcer.

The man had never been put under suitable treatment, and all sorts of absurd applications had been made to the ulcer on the recommendation of friends. At one time continual poulticing was used, which, he said, however, never failed to remove the pain for a time. Then my predecessor here, the late Mr. Wilson, who acted on the homoeopathic system, prescribed some drops, which, of course, did no good either to relieve pain or help the curing of the sore leg. At one time, also, the man was sent to Glasgow Infirmary at the expense of the parish, on the suggestion of a member of the board and with the concurrence of the parish doctor. There he was told by one of the surgeons that it was a hopeless case; and, seeing that there was little prospect of a cure, he left the infirmary and returned here again, to be informed by the parish doctor "that there was no chance of effecting a cure", and that, even if it were possible, it would be dangerous to attempt it. Such is the history of the above case, and I believe the man's statements to be true; inasmuch as he enjoys the character of being a truthful and honest workman, and most of the inhabitants of the district know well how he has suffered so long.

I diagnosed the case to be one of what is called "diffuse hypertrophy of the skin and subcutaneous connective tissue", or "pachydermia". The ulcer was merely a concomitant.

The Treatment I adopted, and it has been so far successful as to reduce the size of the leg, cure the ulcer, remove the dropsical swelling of the foot and ankle, and improve the condition of the skin, lasted over a considerable length of time, nearly five months. First of all, I determined to attack the ulcer and bring it into a healthy condition. I used at the commencement, as a dressing, a diluted solution of permanganate of potash applied on lint, which at first was changed twice in the day. After a few trials of this, I resolved to apply a solution of carbolic acid, in the proportion of about two drachms of the acid to eight ounces of water, along with a little glycerine. This soon told on the sore, as it soon commenced to granulate and new skin to form gradually. Very hot weather setting it, I discontinued the watery solution of carbolic acid, and used instead an oily solution of the acid mixed with glycerine, and this did remarkably well. Occasionally, slight venous oozing appeared at the surface of the ulcer and around its edges; but this was counteracted by the application of a solution of zinc sulphate. Soon the wound began to fill up, the hard edges to disappear, and, after a protracted period of over several months' duration, the ulcer entirely healed over. But from the very commencement, I adopted the measure of bandaging the limb from the foot up to the knee, and this was done daily, and often twice in the day, with few exceptions. The man was also kept as much as possible in the recumbent position, with the foot slightly elevated and the limb laid on a pillow. I also gave him internally iodide of potassium in five-grain doses, combined with bichloride of mercury, one-twelfth of a grain, in compound decoction of sarsaparilla, and the compound tinctures of cinchona and of cardamoms. This he continued to take from the first, and occasionally mild symptoms of iodism were produced, but the dose was easily stopped. This dose of the above drugs he took thrice daily, and at the same time occasional doses of saline purgatives were given to keep the bowels regular. When the ulcer healed, I used, for the psoriasis, the inunction over the affected skin of, first, carbolic acid in solution with oil and glycerine. Then I afterwards tried benzoated oxide of zinc ointment, the nitrate of mercury ointment diluted with lard, etc., and latterly used nothing, but simply bandaged the leg. The leg is now almost as well as its neigh-

hour as regards thickness, colour of skin, etc.; but there is still present some psoriasis, which, I hope, will soon disappear.

The above case is interesting, as showing how rest, conjoined with other auxiliaries, may cure what at first sight might appear to be an incurable disease.

CLINICAL MEMORANDA.

THE CONDITION OF THE COSTAL CARTILAGES IN PHTHISIS.

IN the very able review of the fifth volume of Ziemssen's *Cyclopædia of the Practice of Medicine*, written evidently by an accurate observer, attention is directed to the statement of Freund with reference to the ossification of the cartilages of the first ribs, which is mentioned and, to a great extent, adopted by Rühle in his article on Pulmonary Consumption.

On referring to the article by Hertz in the same volume, I find that Hertz states (I quote from page 359) that he has not been able to convince himself of the actual occurrence of the primary changes in the cartilage as specified by Freund in cases of emphysema; and I should be glad to learn what is the experience of other pathologists in this country as to the condition of the cartilages in phthisis.

In statements of this kind, it is desirable to know whether a large number of observations will support what may seem to some to be a very plausible and pretty theory; but pretty theories are generally to be viewed with more than ordinary suspicion, as their very prettiness is apt to produce a strong bias in the observation of facts. My experience as pathologist to the Hospital for Consumption at Brompton is that, in phthisis, the cartilages are invariably extremely soft, and, during the past year, I can state that there has not been a single occasion in which the saw has been used in opening the thorax; indeed, the greatest caution has to be observed in the use of the knife, in order to avoid wounding the lung beneath, the cartilages being very easily severed. My attention was very soon directed to this point, and that before I was aware of Freund's observations, because this experience was in strong contrast with that which I derived in a large metropolitan hospital to which I was attached for many years, where the saw was frequently called into requisition by the condition of the cartilages. In that, as in other general hospitals, cases of phthisis are generally rejected, although, of course, not completely excluded.

If this appear to be a rough way of answering the question, I can only say that it appears to me to be a convincing answer; but I should be glad to know whether other pathologists are of my opinion in the matter.

REGINALD E. THOMPSON, M.D., Senior Assistant-Physician
to the Hospital for Consumption, Brompton.

SURGICAL MEMORANDA.

AURAL THERAPEUTICS.

IN connection with the report of Dr. Purves's aural *clinique* at Guy's, and with Dr. A. Morison's memorandum in the JOURNAL for December 23rd, I wish to add one or two notes from my experience.

Dr. Purves, alluding to attempts at overcoming adhesions of the membrana tympani to the ossicles or promontory, says "he uses Siegle's exhausting speculum, which, though originally intended as a means of diagnosis, answers very well for forcing the membrane to and fro when used with forcible suction". I have used Siegle's speculum for a long time almost entirely as a means of treatment; as a mode of diagnosis, it is far less efficient. In order to obtain the advantage of both these purposes, I had the largest sized speculum usually supplied with Brunton's auriscope closed at the broad end with a piece of plain glass fixed at the same angle as the reflector, and at the side of the speculum a hole was made and a piece of tubing with a mouth-piece attached.

A Siegle's speculum was thus provided, which could be adapted to the body of Brunton's auriscope, and, by making the patient perform Valsalva's method at the same time that suction was used, considerable outward movement of the membrana tympani was produced and distinctly visible.

This adaptation of Siegle's speculum differs from Dr. Morison's in not requiring a separate Brunton's apparatus, and, being made of vulcanite, is very light and handy.

Dr. Purves alludes to an improvement in the aural speculum by cut-

ting out a piece at the side. I do not know whether Dr. Purves regards this as a novelty; but, in the JOURNAL for July 4th, 1874, there is a paper and woodcut of a speculum I suggested, with an aperture at the side for the use of instruments to the membrana tympani, etc., and adapted to Brunton's auriscope. At that time, I used a speculum cut away at the side, like an anal speculum; but, as I found the tragus or soft parts of the ear projected into the aperture and obstructed the view, I preferred the form depicted in the JOURNAL for the purpose of local applications, etc., to the ear. Both forms I now find useful; the one with the aperture is best for caustic applications, and that with the piece cut out for the use of instruments, probe, etc.

I showed Messrs. Weiss and Sons, Strand, a pattern of these specula some time ago, and they said they had not seen anything like them; but, on a recent visit to their premises, I saw that they had made several in vulcanite from my pattern.

Messrs. Wood and Co., Manchester, have made the originals for me.

F. M. PIERCE, M.D., L.R.C.P.L., Manchester Ear Institution.

REPORTS

OF

MEDICAL AND SURGICAL PRACTICE IN THE HOSPITALS AND ASYLUMS OF GREAT BRITAIN.

HOSPITAL NOTES.

LONDON HOSPITAL: DR. HUGHLINGS JACKSON'S CLINIQUE.

Syphilitic Brain-Disease.—In a man, aged 29, this diagnosis was made on account of a "random succession" of nervous symptoms, though no history of syphilis could be obtained. The importance of such a "random succession" or "random association" of nervous symptoms, in forming such diagnosis, was strongly insisted upon. The patient had, first, numbness of the left arm and left hemiplegia; then impaired mental condition (delusions and partial imbecility); afterwards, paralysis of the left third nerve; and, finally, right hemiplegia and aphasia. Although aphasic, he could utter "Yes" and "No"; these are the most automatic words, and are oftenest retained. Indeed "Yes" and "No" are, Dr. Jackson remarked, in effect, propositions, whatever their philological origin may be. Since the patient's admission, he has become able to utter a proposition but little above these in automatic character, viz., "Thank you". It is well known that oaths (deeply automatic with some people) may be uttered by aphasic patients under excitement when they cannot be spoken. This patient, however, could not only utter the phrase "Thank you", but said it when told. He has no optic neuritis; and, if this generally occur in such cases, it is not that a syphilitic growth as such produces it, but only as a glioma or as any other "foreign body" would. It is important to examine frequently for the onset of neuritis, which may occur at any time in such a case. The outcome of the diagnosis was treatment by mercury, and the patient has been taking one grain of hydrargyrum cum creta thrice daily; he is very much better in his general condition, is less imbecile; his speech is a trifle better; but he has still complete ptosis, and is still in danger. Since this note was made, the patient has not improved: he is now having mercurial inunction. Dr. Hughlings Jackson remarked that he had known paralysis of an ocular motor nerve to completely disappear in a patient who had, as the necropsy showed, cancer of the base of the skull—a recovery he could scarcely have believed possible under the circumstances.

Hemiplegia: Thrombosis.—A man, aged 69, when driving his omnibus eighteen months ago, suddenly went giddy, but drove on for some distance. He was assisted to his home, did not lose consciousness, but got right hemiplegia. Some months afterwards, he had necrosis of the great toes of both feet. From the deliberate onset of the hemiplegia, from the age, and atheromatous state of the vessels, Dr. Jackson presumed a thrombosis in a small branch of the middle cerebral artery, and thought a similar local plugging of arteries in the feet to be probable. There was no albumen in the urine. The man recovered from the paralysis, except for some slight weakness of the right arm. We are not warranted in inferring that the recovery is dependent upon restoration of cerebral circulation by anastomoses of other vessels—(1) because vessels in the corpus striatum have scarcely any anastomoses; (2) because, in some cases of patients who have recovered and afterwards died, Dr. Hughlings Jackson has found local lesion persisting. Recovery is, he thinks, brought about by other parts of the centre becoming sufficient for limb-movements. Certainly, recovery goes so far