was owing to the insensibility of the glottis, or whether the displaced cannula had made an opening into the œsophagus from the pressure of its point against the back of the trachea." It is quite conceivable, no doubt, that the sensibility of the glottis may become impaired, seeing that, so far as respiration is concerned, it is now in a state of disuse; but there occurs to me still one other possible explanation of, at least, some of the cases. In all the three to which I have referred, the original disease was diphtheria; and may the passage of liquids into the trachea not be accounted for by the awkwardness of deglutition, which, we know, does exist during this stage, and which often leads to the regurgitation by the nostrils of part, at all events, of the fluids which are swallowed? May it not, indeed, be a paralytic symptom?

I have mentioned, incidentally, ulceration of the trachea by pressure from the end of the tube, but I am entirely without experience of it. If, as is averred by some, a bloody character of the sputa coughed through the tube be an indication of its occurrence, then, to this extent, I have observed it. But this is just as much an indication of the separation of false membranes, for I have seen blood flow from the mouth in considerable quantity during convalescence from diphtheria affecting the nostrils and fauces, but not at any time implicating the larynx, and therefore never at any time calling for tracheotomy. To mitigate, however, any evils due to the pressure of the tube, it is well to use one in which the cannula is movable upon the shield. Such tubes have, at all events, the merit of being worn with much greater comfort.

The management of false membranes or inspissated mucus obstructing the windpipe below the tube, and so frustrating the object of the operation, is a matter of some difficulty. Frequently Nature effects their expulsion in good time, and, on the whole, it is probably best to trust to what may be done in this way. I have in one case, as has been advised, removed the tube, and introducing forceps down the trachea, laid hold of whatever they could catch between their blades and withdrawn it. But it produced no improvement; on the contrary, it was probably prejudicial. Guersant and others advise sucking the tube, but in diphtheria the practice is a highly dangerous one, and hardly warranted in view of its small chance of success. Speaking of this recommendation, Giraldès says in his admirable *Clinical Lectures on the Surgical Diseases of Children*,—"The counsel is excellent, but, in acting thus you run the risk of contracting diphtheria, a disease, in my opinion, essentially contagious. You find yourselves, then, face to face with a question of feeling (*question de sentiment*). If you have courage to face the danger, your conduct will merit the greatest praise. Give over, on the other hand, to Nature the task of unloading the bronchial tubes, and no one can accuse you, no one can blame you." That the disease is essentially contagious is indicated, I have sometimes thought, by two facts, in regard to which I shall be glad to have the experience of any gentleman present. I have a strong impression, first, that while surgeons, who operate in a case of diphtheria, or remain in pretty close attendance upon it afterwards, often contract the disease, physicians visiting and prescribing for a case requiring no operation do not appear to run the same risk; and, secondly, that when an adult in a family contracts the disease from a child, that adult is most frequently the mother. Maternal affection and grief lead to the frequent kissing of the child in spite of warning.

All of us are familiar with the difficulty often experienced in supplying warm and moist air to the little patient who breathes by a tracheotomy tube; and many must have had experience of the trouble and toil which is implied in keeping up a supply of steam in the neighbourhood of the child's head. I am satisfied that this may be dispensed with for the following much more simple mode of after-treatment. If a sponge—the larger the better, a bath-sponge answers well—be wrung out of boiling-water and placed lightly over the tube, it will be found to be an excellent respirator, furnishing an abundance of warmth and moisture for at least a quarter of an hour; at the end of which time it ought again to be wrung out. The child inspires through the pores of the warm sponge, and, by its expirations, tends to keep the under surface warm even after other parts of it are being cooled. It may be fixed lightly in position by a tape or piece of bandage tied over it, and can be easily removed and re-applied even during sleep. It is, moreover, a grateful application to the part, especially when, towards the close of the second or beginning of the third day, a slight blush of inflammation surrounds the wound, and the necessity for slackening the tapes tying in the tube indicates some swelling of the neck.

I must now conclude these very disconnected observations, and I take leave to do so by commenting in a single sentence on a matter which, in spite of much that has been said and written, is not generally sufficiently borne in mind. It is this, that the successful issue of a tracheotomy case is the relief of asphyxia, and not the cure either of croup or diphtheria. The little child who has been snatched from

a death by suffocation to die a week or fortnight hence from gradual asthenia, a sudden syncope, or an unexpected convulsion, is as real, though not so apparent an example of the benefits of the operation, as that other who ultimately survives both operation and disease. Let us not, then, be either too elated by successful statistics of recovery or too discouraged by the reverse. Provided the operation have been performed without accident and the breathing fairly re-established, there is much good fortune in the one experience and but small blame in the other. If we deal with large numbers, all statistics seem to be pretty much equalised, and recovery is found to take place about once in every three or four cases; but it is well to remember, that this leaves out of sight altogether much genuine success achieved by the operation. By the performance of it, we do nothing towards curing the disease; indeed, we are rather apt to encourage its extension. All that we can do—and, therefore, all that we should pretend to attempt—is to restore Breath, and with it its companion—Hope.

REMARKS

ON

THE PATHOLOGY OF CHOREA.

By H. CHARLTON BASTIAN, M.A., M.D., F.R.S.,

Professor of Pathological Anatomy in University College, London; Physician to University College Hospital, and to the National Hospital for the Paralysed and Epileptic; etc.*

Etiology of the Disease.

It may, perhaps, be safely affirmed that nearly nine-tenths of the cases of ordinary chorea or St. Vitus's dance occur in individuals either—

1. In relation with rheumatic fever;
2. As a sequence of fright; or
3. In association with anæmia, chlorosis, or some other cachectic condition.

We know also that it tends to occur all the more readily in children, especially between the sixth and fifteenth years, who have been born from nervous parents, and who themselves show signs of an undue excitability of the nervous system. Occasionally it occurs in adults, or in persons who seem in other respects healthy, except it may be for some slight menstrual disorder, or for the fact of the coexistence of pregnancy, as in cases specially referred to by Dr. Barnes.

These are the known facts, and the question is how we are to deduce from them a pathogenesis for the disease which shall be broad enough to be compatible with the whole of them.

The "*Embolio Theory*" seems to me to be lamentably defective in this respect. It is (1) so narrow that, even if more were capable of being said for it than now seems possible, it could only be considered to apply to one small section of the total number of cases of chorea, and would necessitate our framing some wholly independent hypothesis in explanation of the remainder. That I am not overstating the case against the theory will be plain to those who recollect that, in the large majority of the cases of chorea supervening in individuals belonging to the second and third categories, there is no evidence of the existence of endocarditis at the time of the onset of the disease; and that this is also true of a certain number even of the cases belonging to the first category—that is, to those by no means uncommon cases in which chorea follows rheumatism where there is no heart-disease, as well as to those rare cases in which it precedes the articular affection. But other strong evidence against the "*embolic theory*" may also be cited. Thus (2), chorea is not sudden in its mode of onset, as it should be if this theory were true. In the large majority of cases, it is quite the reverse, so that it may be said to be the rule for chorea to manifest itself gradually. (3.) The valvular disease which does coexist with a large proportion of the cases of chorea, when the disease is fully established, is usually not of a kind to favour the separation of embolic particles, consisting as it does, for the most part, of a row of small bead-like elevations near the free margins of the mitral valve on its auricular surface. It is, in fact, an incipient endocarditis of low intensity, in which the elevations are in the main composed of a coherent overgrowth of the valve-substance, without the addition of those more brittle fibrinous concretions which form so large a part of

* Concluded from page 38 of last number.

the "vegetations" produced under the influence of rheumatic fever.* Without referring to other grounds, which might be more open to dispute, these alone seem to me to suffice for the rejection of the "embolic theory" as an explanation of any large number of the cases of chorea. It is, however, quite conceivable that, in certain accidental cases, multiple minute embolisms of the corpora striata might occur and give rise to this affection. I regard embolism, therefore, as a possible cause, but cannot consider that it frequently leads to the production of chorea.

What better explanation can, in fact, be found for such a case as this, about which I was consulted not very long since? E. P., a girl aged 14, had about twelve months previously suffered from rheumatic fever and endocarditis, which left a mitral systolic murmur, heard very distinctly when she came under observation. Four months after the rheumatic fever, she was suddenly attacked with choreal movements in the right arm and leg, and at the same time with thickness of speech. After a few days she was observed to be spiteful, and soon showed other signs of an altered mental state. The chorea preserved its unilateral character throughout. Now in this case, looking to the pre-existence of a rheumatic endocarditis, to the sudden onset of the chorea, to its limitation to the right side, as well as to the altered mental state and to the absence of any other distinct exciting cause, I think we are warranted in supposing that small embolic particles were more or less simultaneously swept away along the most frequent channels into the left middle cerebral artery, and thence distributed partly to the corpus striatum and partly to the convolutions.

The problem before us is now, from this point of view, decidedly simplified. In the cases belonging to our first category, chorea by far the most frequently supervenes as a post-rheumatic complication; and, if the causative relation between it and heart-disease may be dismissed, then we are thrown back upon the blood-state or general constitutional condition following rheumatic fever as that which favours the occurrence of chorea. This post-rheumatic state is eminently an anæmic and debilitated condition, so that, from this point of view, the cases of chorea pertaining to our first category are found to agree in their conditions of origin pretty closely with those belonging to the third category above mentioned. So far, therefore, we seem to learn that chorea is most prone to occur in individuals of a certain age whose nervous tissues are unduly irritable, partly from their inherent constitution, and partly under the influence of anæmia and other allied though more special blood-states variously induced. The degree of influence of these several factors must, of course, vary much in individual cases.

What, however, is to be said concerning those numerous cases pertaining to our second category—those in which fright seems (to Dr. Dickinson as well as to myself and others) to act as the immediate and undoubted exciting cause? To be brief, it appears to me that these cases may be ranged under two heads; viz., (a) cases in which, with an irritable nervous child, the fright causes such an enormous downrush of molecular motion from centres above to those below in the corpora striata as to set up in them more or less immediately, without the intervention of an altered blood-state, "functional disease" if not structural lesions; and (b) cases in which the disturbance of the molecular life of the lower centres occasioned by the shock was not of itself sufficient to set up the disease, but where the results of the shock have been profound, wider, and more lasting. In these cases, a state of lowered health is gradually established; and in two or three weeks, or even a much longer time after the mental shock, the symptoms of chorea may begin to manifest themselves.

The first or immediate mode of action of fright (although different authorities may have their preferences as to the exact mode of stating it) is probably that which is most commonly entertained. But, whilst admitting the difficulty of the question, I am strongly disposed to think that the second or remote mode of action of fright—that is, by its induction of an altered blood-state—also obtains in certain cases where chorea follows some severe mental shock with a short or not very prolonged interval. My reasons are these. First, I have seen several well-marked cases, and Dr. Broadbent refers to others, in which the most profound and lasting alterations in general nutrition have set in even in adults after a very severe fright or terror. Secondly, in some of the cases in which chorea follows fright, patients do, as a matter of fact, become distinctly anæmic and thinner before or whilst the chorea is developing. Thirdly, it is not difficult to understand how this altered condition of nutrition may be brought about by disturbance of the sympathetic system of nerves, either through the abdominal ramifications of the

pneumogastric or through its own connections with the cerebro-spinal system. Fourthly, even in some of the cases of chorea arising from fright, a mitral murmur becomes developed in the course of the disease; and, when such cases have chanced to be examined *post mortem*, the usual endocarditic beading of the mitral valve has been met with. These latter facts have been brought out in a very clear way by Dr. Dickinson in the valuable paper already referred to, although he interprets the mode of origin of the heart-disease (*loc. cit.*, p. 37) in a manner which I am unable to accept. But this part of the subject must be more specially referred to.

The portion of Dr. Dickinson's communication which seems to me the least convincing is this section, in which he seeks to interpret the occurrence of the slight endocarditis which has been met with in no inconsiderable number of the cases of chorea induced by fright, and in which the corresponding mitral murmur has been known to have developed with the chorea. There is to me nothing whatever peculiar in the character of the endocarditis which is associated with these cases of chorea. It occurs in the usual situation; but, as in many other cases of chorea, it is limited in extent, owing, as I believe, to peculiarities in the state of the blood usually associated with this affection. Microscopical examinations of the blood drawn from the tip of the finger in twelve average cases of chorea, as I find from notes taken in 1869, were principally remarkable for the fact of the absence of separated threads of fibrine, and for the absence of an increase in number of the white corpuscles, or of any exaltation of their amoeboid activity. To these peculiarities of the blood I ascribe the fact that the proliferation in the substance and on the surface of the valve is not reinforced by those depositions of fibrine and white corpuscles which together suffice largely to augment the endocarditic lesions occurring during the course of rheumatic fever. My notes of the examination of the blood in many of these latter cases, made also in 1869, show that in them there is always more or less of increase not only in the number, but in the amoeboid activity of the white corpuscles; and that the amount of fibrine which separates in the form of needles or threads is almost always increased, and is often excessive in amount.

If, however, there be nothing peculiar in the nature of the endocarditic lesion pertaining to these fright-engendered cases of chorea, the same may be said respecting the cause assigned by Dr. Dickinson. If irregular action of the heart could so readily induce actual disease of the mitral valve in chorea as Dr. Dickinson seems disposed to think, it may fairly be asked how it is that the much more prolonged irregular action so frequently met with in non-choreal cases of heart-disturbance does not lead to the same result? I certainly am not aware that any such sequence is either with or commonly looked for by the majority of physicians.

I prefer, indeed, to believe that here also the explanation of the endocarditis is to be found in an altered blood-state generically similar to that met with in other cachectic states with which chorea is often associated, though produced in these particular cases under the remote influence of fright. Thus, so far as we actually know at present, acute inflammation of the endocardium only arises under the coexistence of altered blood-states, though these may differ a good deal amongst themselves according as they pertain to rheumatic fever, puerperal fever, scarlet fever, measles, Bright's disease, or certain less definite cachectic states of the system—which may follow rheumatic fever, may be associated with menstrual irregularity, or with some other less appreciable cause.

In accordance with the views already set forth, I should expect that hemichorea would not unfrequently be caused by the direct action of fright, and that so long as the affection continues to be unilateral, it will be less frequently complicated by a supervening endocarditis than when the disordered movements are bilateral. Embolism, also, would seem, where it does act as a cause, to be more likely to be instrumental in producing an one-sided than a general chorea. On this account, therefore, and also because a previous endocarditis of rheumatic type existed, it would not be improbable that the hemichorea in the case detailed by Dr. Stephen Mackenzie, in his communication to the JOURNAL of December 23rd, may have been occasioned by embolism. On the other hand, it also seems to me, that his case is very fairly open to a different interpretation. As he himself says, the child was "of an age, sex, and temperament which specially predispose to chorea"; and she was subjected to the emotional strain and depressing influence almost necessarily resulting from her mother's death. It is further worthy of note, that the choreal symptoms did not set in immediately after the "vascular excitement" which was presumed by Dr. Mackenzie to have been the occasion of the sweeping away of embolic particles from the diseased mitral valve. They first began to show themselves ten days after this date—which is a fact of no small significance.

* Of course, in those cases in which heart-disease has been engendered during the course of rheumatic fever, and this state is followed by chorea, a condition of the valve more favourable for the sweeping off of minute embolic particles would undoubtedly exist.

Pathogeny of the Morbid States or Disordered Actions, in certain Parts of the Nervous System.

This final problem is one not peculiar to chorea; in essence, at all events, it is similar to that which requires solution in many other cases where abnormal blood-states favour the occurrence of local lesions. There are, as we all know, seats of election—that is, particular tissues are apt to become the seats of inflammation, or some other pathological change in different diseases. In typhoid fever, the small intestine is gravely affected; in scarlet fever, the throat and tonsils suffer; in rheumatic fever and other states, the endocardium of particular parts of the heart inflames, just as in lead-poisoning, particular muscles of the body are especially implicated. It seems to me, therefore, a fact of the same order when we find that a certain altered blood state predisposes certain individuals not only to a subacute endocarditis, but to an irritation of certain specially active parts of the brain, which goes on to produce what we may call a subacute inflammation.

The intimate processes of nutrition in different tissues of the body and in different parts of the same anatomical system, must vary with the chemical composition of such tissues, and is, in all probability, subject to variation under influences which cannot be thus roughly expressed. Hence, a certain blood-state differing from that which is habitual to the individual may easily pass beyond the range of what is compatible with the continuance of an ordinary nutritive and functional activity in the elementary parts of certain tissues. Such tissue-elements become “irritated” (what other word can we use?) and begin to attract more blood. In this we have a state corresponding with what I apprehend to be the first stage of the perverted function which shows itself in the corpora striata and adjacent parts at the commencement of chorea; though, as I have already hinted, a somewhat similar condition may be brought about in different ways, and sometimes would seem to be occasioned by the direct action of fright (*Paralysis from Brain-Disease*, pp. 27-30). At other times, indeed, no distinct cause, either predisposing or exciting, can be assigned (as with other nervous diseases of the same class), though such instances are frequently ascribed to some source of “reflex irritation”. But, howsoever induced, if the irritation continue, the disturbed action outside the vessels communicates itself to the tissue elements inside the vessels, as they come within the morbid area. The white corpuscles, as a consequence, begin to adhere to the walls of the small vessels, so that partial obstructions may be produced, which are, perhaps, thereafter rendered complete by separation of fibrine or allied products. And in certain exceptional cases, such as those which have been encountered by Dr. Aitken and Dr. Tuckwell, small foci of softening may be produced.

After detailing the results of his own observations, Dr. Dickinson writes (*loc. cit.*, p. 16). “Arterial repletion seems mainly concerned in the development of the disease; why, or at what bidding do these vessels thus gorge themselves?” The answer which I should be inclined to give to this question has now been briefly sketched, and I am pleased to find that, in many respects, Dr. Dickinson’s interpretation does not very materially differ therefrom.

Multiple occlusions of capillaries and small vessels may easily give rise to the minute vascular injections which have now, on so many occasions, been met with in the corpora striata and adjacent parts of the brain in fatal cases of chorea (see *Pathological Transactions*, 1869, p. 15). It is, however, especially needful in an affection of this kind in which the only easily recognisable *post mortem* mark is one of hyperæmia, not to confound that which lies at the root of the disease with other hyperæmias more or less secondary in their duration and mode of origin. Certainly, this is a point which should be carefully borne in mind by future investigators.

In brief, then, my view of the etiology and pathology of chorea is this. In common with many who have spoken with greatest authority on this subject, I look (certain rare cases excepted) to an altered and often anæmic blood-state as its predisposing cause in individuals of a certain age and nervous temperament. Secondly, I look to the initiation in such individuals of a disturbed nutrition in the corpora striata and adjacent parts of the brain, tending to issue, and often actually issuing, in what, for want of any other more appropriate term, may be called a subacute inflammation of these centres—often characterised in part by the production of multiple minute thromboses. It will be seen, therefore, that I hold no simple theory which can be embodied in a single phrase, though, as regards the great majority of cases of chorea in which there are local lesions—and these are well marked—it is quite true that I would replace the embolic theory by what Dr. Hughlings Jackson has termed the “hypothesis of thrombosis”.

CONSUMPTION AND CLIMATE.

By RICHARD PAYNE COTTON, M.D., F.R.C.P.,
Consulting-Physician, and late Senior Physician, to the Hospital for Consumption, Brompton.

DR. REEVES of Melbourne, in his recently published work on *Consumption in Australia*,* has, I think, done us a real service. He shows what, I suppose, all of us were well prepared to hear, that phthisis is scarcely less frequent and fatal in Australia than in England; and that, however modified the disease may be at our antipodes, it is nevertheless there, ever ready to claim its victims, and to show us that there is no exception in that country to the universal presence of that dread malady. And yet how many of us are in the habit of sending certain of our phthisical patients to Australia, and of finding that, however often the result may have been unsatisfactory, yet, in many instances, our best hopes have been even more than realised.

Some years ago, I was consulted by a young medical friend, already decidedly phthisical, about his settling in Australia. He went there with my approval, and is now one of the most successful medical practitioners in the colony, and occasionally sends me a letter, saying that now he can equally well both see his patients and hunt the kangaroo. Several similar, and quite as promising cases have at various times come before me; but circumstances have not permitted me to trace for any length of time their subsequent histories.

On the other hand, and quite in accordance with Dr. Reeves’s statements, our *confrères* in Australia not unfrequently send their native consumptive patients to be cured in England. Two such examples have lately fallen under my own observation; in the one, the change was productive of the happiest results; in the other, however, the disease ran on with unchecked rapidity.

How are such circumstances reconcilable? Is it not apparent that there cannot exist, either in the one locality or the other, any special or *specific* influence, and that any benefit to be derived in either instance must be due to other causes? Whether the tubercular diathesis arises here or in any foreign climate, a change from that climate and from every circumstance of life connected with it, a rest from the work and conditions under which it has arisen, a new field of occupation and thought, and especially, as in our relation to Australia, where all this is prefaced by a long sea-voyage, constitute, indeed, the real and only active influences for good.

Early cases of phthisis, more especially those of a strongly inherited origin, where the pulse is not much quickened; the breathing calm; where there has been no recent and never any active hæmoptysis; cases of the “languid” rather than the “florid” type of the disease present the conditions upon which the changes I have just enumerated would be the most likely to tell, and upon which they certainly often do tell very satisfactorily.

And here, in connection with this subject, I would make a passing remark upon the now rapidly dying out opinion, that there is anything in any one climate, wherever it may be, which is opposed to the development or even the increase of tubercular disease. Consumption exists everywhere. There is no favoured spot where it is not. There is no “promised land” for our consumptive sufferers, where they will fail to meet with native consumptive sympathizers.

It is quite true that the form of tubercular diseases, their degree, severity, and duration, may differ considerably in different parts of the world, according to the habits of the people, and the relation of the climate to such habits; but there is no favoured country where tubercle has not a birth-place. Trouble, privation, numerous sources of unhealthiness uncontrolled, and many perhaps uncontrollable, the evil habits of civilised life, the vices of individuals and of society, have, unhappily, no geographical limit, but are everywhere exercising their influence, although, of course, in differing degrees, both in originating and arousing tuberculous diseases. Consumption’s old and special title of “English disease” is as undeserved as it used to be common.

Some years ago (Fothergillian Prize Essay), I published the result of numerous inquiries on this subject. I found that, everywhere, whether in cold, temperate, or hot climates, consumption was known only too well by my professional brethren. Dr. Wilson of New Brunswick stated that “the great majority of the adult aborigines of North America died of phthisis, and that the disease is very common amongst European settlers”.

In more temperate climates—take, for instance, the very health-resorts of the consumptive invalid—the shores of the Mediterranean, Italy, the South of France, Spain, and Portugal—phthisis is to be found. Dr. Burgess says (*Lancet*, July 1850): “There is no part of

* Reviewed in BRITISH MEDICAL JOURNAL, Decembe 30th, 1876.