stricture, and will also give relief to the overdistended cæcum; whilst, as for the remainder—that is, those cases in which colotomy fails in its object—enterotomy should be performed and relief thus afforded, although it may be with but a very imperfect conception as to the exact locality of the source of obstruction.

[Since this paper was written, we have been informed by Mr. Maunder that, after hearing the views expressed therein respecting right colotomy before gastro-enterotomy, he was prepared to act on the suggestion in a case under his care; but the necessity for enterotomy did not arise, as he found the right colon distended.]

THE SILK LIGATURE AS A METHOD OF SECURING THE OVARIAN PEDICLE:

ITS ADVANTAGES AND DISADVANTAGES, WITH BRIEF NOTES OF ITS RESULTS IN THIRTY-EIGHT CASES.

By J. KNOWSLEY THORNTON, M.B.,

Surgeon to the Samaritan Free Hospital for Women and Children.

SOON after commencing to perform ovariectomy, I determined to give the silk ligature a fair trial in a series of consecutive cases. It seemed to me worthy of such a trial, because of its universal applicability, its cleanness and neatness, both at the time of operation and in the after-progress of the case, and because it yielded such good results in the hands of those who only used it in cases in which it alone was applicable.

With the works of Spencer Wells and Peaslee in our hands, in which the whole question of its use is ably discussed, a paper on this subject may seem unnecessary. It differs, however, from the treatment of the subject in either of the works named, inasmuch as it deals only with one of the methods of using the silk ligature which they describe, viz., that in which the ligature or ligatures are made to transfix the pedicle, both ends being then cut off close to the knot and returned with the pedicle-stump free into the peritoneal cavity. I should hope that the old plan of leaving one end of the ligature hanging out at the lower angle of the wound has for ever passed away. It has no advantages to recommend it, and at least two serious disadvantages to contraindicate it. It necessitates the cutting off of the distal end of the stump by ulceration, and, during this slow process, it keeps open a moist highway for the causes of irritation and putrefaction to travel by from the exterior of the wound to its deepest recesses; to the very parts, in fact, which we wish to heal as quickly as possible.

Before proceeding to discuss the details of its application and the difficulties and dangers to be encountered, I will sketch briefly the various methods by which the healing process is accomplished and the ligature disposed of.

It always seemed to me probable that, though the tissues included in the ligature were tightly enough constricted to prevent hæmorrhage, they were not usually so grasped as to completely destroy the nutrition of the distal end of the stump. The following observation confirms the correctness of this view. In one of my cases, I tied a large piece of omentum adhering to the cyst in two places and divided it between the ligatures, thus leaving a stump on the outside of the tumour very like that of the ovarian pedicle. I afterwards injected the cyst with Beale's Prussian blue fluid, and, when the injection was complete, noticed that the distal end of this stump was blue, though not so blue as the proximal part, and there was no appreciable escape of the fluid from its cut surface. Microscopic examination showed that the injection had passed into the minute capillaries, but not into the larger vessels, except where it had passed into them by the anastomoses beyond the ligature.

If the injection would so pass, the blood would also pass; and in this way the vitality of the part would be at any rate partially sustained until the other processes by which the distal end of the stump is nourished had time to be completed.

The five following conditions of the stump after ligature I have myself observed in the dead or living subject. I do not claim any novelty for the observations, but, so far as I am aware, they have not as yet been all recorded side by side.

I. The ligature is buried in the peritoneal coat of the pedicle, the portions on the opposite sides adhere, and vascular connections are thus rapidly established. Cells penetrate between the fibres of the silk and break up the ligature, which is absorbed and disappears. The finer the silk used, and the softer the pedicle, the more likely is this to happen.

II. Lymph is thrown out over the ligature and cut end of the stump; in this new vessels are formed, and the same result is obtained, but more slowly.

III. The stump adheres to some neighbouring peritoneal surface, and from that new vessels pass into it, through which its nourishment is obtained.

It will at once be seen how the passage of some nutriment through the capillaries in the constricted portion must aid either of these processes, and we must also remember that the pedicle is surrounded by the nutritive fluid naturally present between the peritoneal surfaces, and probably always somewhat increased when repair is going on.

The first two of these methods are the most favourable, and are those which we must aim at securing. As a result of either of them, the stump may be in a few months merely represented by a little button of smooth tissue at the site of the pedicle. Such a result I had an opportunity of seeing when assisting my colleague Dr. Bantock at an exploratory operation a few months after ovariectomy. The exploration revealed the above condition of the pedicle, which was not adherent to any neighbouring surface, and was also quite free from the malignant disease of the peritoneum which had caused reaccumulation of fluid. How often such a result is obtained it is of course impossible to say, but I think we may fairly suppose it to have occurred when there is no symptom of any irritation about the pedicle, or at any rate in the majority of such cases (*vide* Cases I, VII, VIII, X, XI, XIV, XV, XVII, XVIII, XXII, etc.). When the stump adheres to some neighbouring peritoneal surface, most commonly some part of the intestine, because they are generally in contact with it, we are by no means likely to get such a good result. We may have no sign of mischief for some time, usually not till after the bowels have been moved. Sometimes, however, some intestinal irritation shows itself much earlier, especially if there be trouble with flatulent distension; indeed, the latter is often a sign of the irritation. When symptoms follow the first action of the bowels, they usually appear from the tenth to the fourteenth day. There is usually some rise of pulse and temperature; this may be considerable, and is, I think, proportionate to the extent of adhesion which has been disturbed. There is often depression of spirits, pain about the pedicle, and sometimes hardness may be detected in the iliac fossa, or by vaginal or rectal examination. All the symptoms may gradually subside, or they may persist for some time with obstruction to the passage of the flatus and fæces or with troublesome diarrhoea and tenesmus. In the former case, fatal obstruction may result; in the latter, the case will end in discharge of pus by rectum or vagina, and all will go well. In the cases given below, examples of each of these results may be found except that of fatal intestinal obstruction.

IV. Fatal hæmorrhage may follow the use of the ligature, and one such case (XX) may be found below; and the same accident has been reported to me by several surgeons as having occurred in their practice. This most formidable accident is due to the escape of one or more of the large veins, which are common at the outer edge of the pedicle (pampiniform plexus), from the external loop of the ligature. It requires great care and some experience to avoid it, when treating a very broad and short pedicle.

V. The ligatures may remain uncovered around the pedicle in cases where diffuse peritonitis with effusion of serum or rapid septicæmia follows the operation; and if the patient survive only a few hours, they will be found just as they were when applied; or if she survive for a longer period, they may be found somewhat loose, with an ulcerated groove round the pedicle, the distal end being in a sloughing condition.

I do not think the fatal result is much influenced by the ligature in this last class of cases, though it is possible that it may become early soaked with the products of putrefaction and be a constant source for their propagation; whereas the tissues might have been able to dispose of the purid or irritative matter had no such absorbent material been present to hold it. Excluding these cases from our consideration, we still have a formidable array of dangers to consider; but, fortunately, experience teaches us that the favourable terminations are the most common after the use of the ligature, as the cases here given will show. There being only five fatal cases out of a total of thirty-eight, and only one of these fairly to be attributed to the ligature; two of the others (Cases II and XIX), being hopeless under any treatment, and the other two sufficiently serious to render it impossible to attribute the fatal result to the ligature, at any rate with certainty.

I will now endeavour to point out certain precautions to be adopted in using the ligature, which will, I think, ensure for it results even more favourable than my own have been up to the present time.

First, as to the material. It should be the pure Chinese silk, without any admixture of cotton. The hemp ligature still finds some advocates, but it has, so far as I can see, no advantages, and the great disadvantage of being a vegetable instead of an animal tissue, and it is slippery and cutting to the fingers when wet. One advantage claimed for it is that it shrinks after tying, and hence is a security against

hæmorrhage. From experiments I have made with it, I do not believe it shrinks at all after it is once thoroughly wet, and this it always is before it is tied; and, if it did so shrink, I should not consider it an advantage in view of what I have already said as to the partial circulation through the constricted portion. The silk, if properly tied, never fails to sufficiently constrict. As to the method of applying the silk, I believe it to be important that the two ligatures should interlock when tied, so as to form a figure of eight, otherwise they may pull apart at the point of puncture and hæmorrhage may result. In a thin pedicle, it is easy to avoid puncturing a vessel, and the interlocking of the ligatures is a matter of less consequence; but, in a thick one, it is often impossible to see and avoid every vessel of sufficient size to be dangerous if punctured. If more than one transfixion be required, the ligatures should all interlock so as to form a chain. This has one disadvantage, tending, in a broad short pedicle, to cause a drag upon the outer loop of the ligature-chain, and hence to increase the risk of hæmorrhage from slipping of some of the large veins to which I have referred. The weight of the uterus and other ovary, aided perhaps by the contractile nature of the tissue of the broad ligament, increases this risk. It is well, therefore, in every case, to examine the pedicle-stump carefully just before closing the incision; and, if any sign of hæmorrhage or slipping be found, to apply a fine silk or catgut ligature by transfixion on the proximal side of the other ligatures, this last ligature to include the veins already referred to. I always adopt this plan now, using either fine silk or catgut wherever the outer edge of the broad ligament feels tight as one passes the fingers along it from the pelvic brim to the edge of the pedicle. In all the earlier cases, I employed very thick silk, and seldom more than a single transfixion, with an accessory ligature if necessary. Thinking this thick silk might be the chief cause of the trouble with the pedicle, I tried a medium thickness, tying the pedicle in smaller pieces and with more transfixions. I found the results satisfactory, and I now use the finest silk that will bear the necessary strain in tying.

Since performing ovariectomy antiseptically, I have, whenever it was possible, sewn the posterior peritoneal cut surface of the pedicle over on to the anterior surface of the broad ligament with fine silk or catgut with a view of covering over the raw surface, and at the same time placing it in immediate contact with a peritoneal surface, from which it might receive its blood-supply quickly. This method has also the great advantage of carrying the stump over to the anterior surface of the broad ligament, and so away from the risk of adhering to the intestines. I am not prepared to advocate this proceeding, however, unless the operation be performed antiseptically, because it has the disadvantage of opening up the cellular tissue of the broad ligament, and it is very probable in some of the cases small vessels are also transfixed in their continuity. In cases where the pedicle is very broad, thick, and short, this sort of flap operation would not be possible; but I think a suggestion made to me by Mr. Lister would answer well in such cases, viz., to unite the cut peritoneal edges of the stump with catgut sutures, placing a few strands of catgut between the sutures to drain, and prevent any possible risk from accumulation of serous fluid and tension.

I am aware that the catgut ligature has been successfully used instead of silk for securing the pedicle, but the results with the silk are so good that I shall not change it for catgut, at any rate, until one can be more certain of always getting a good sample of the catgut. As at present sold, it is too uncertain in its quality.

CASE I.—Both ovaries removed; clamp on left pedicle; right one transfixed and tied in two parts with strong silk, one ligature being made to encircle the whole again. Recovery perfect; no sign of mischief from the stump or ligatures.

CASE II (D).—Right ovary removed; pedicle transfixed and tied in two parts and returned. Case of gangrene of cyst after tapping during pregnancy. Died in sixteen hours.

To save repetitions, I may state that in every case, when no special reference to method or nature is made, the strong silk was used, the ends in all cases being cut off close to the knots, transfixion was employed, and the pedicle always dropped free into the peritoneum.

CASE III (D).—Both ovaries; very complicated case; extensive uterine, vesical, intestinal, and omental adhesions; several transfixions and many ligatures used. Wound continued to discharge for months after the operation, and several of the ligatures came away. Now in good health.

CASE IV.—Left ovary; ligatured as in Case I. On sixteenth day, irritability of bowels, passing off in a few days. Went home well on twenty-third day.

CASE V (D).—Left ovary; an enormous solid dermoid tumour;

strong silk soaked in carbolic oil; right ovary; fine silk also, soaked in carbolic oil. Died on fifth day; congestion of the brain. There was nothing to show that the ligatures influenced the result.

CASE VI.—Right ovary. On fifteenth day, I noted "Bowels do not act very well, and patient is evidently suffering from the irritation which ligatured cases so often have twelve or fourteen days after operation". Went home well on eighteenth day.

CASE VII.—Both ovaries; each two ligatures. No sign of irritation.

CASE VIII.—Left ovary; combination of enucleation and ligatures; three medium silk ligatures used. No trouble of any kind. Left hospital on twenty-first day.

CASE IX.—Right ovary; similar to last, but thick silk used. Died on fifth day. (?) Pyæmia.

CASE X.—Left ovary. No trouble. Went home on twenty-second day.

CASE XI.—Enormous dermoid tumour of right ovary, with very extensive adhesions; smaller one of left ovary; two ligatures on each pedicle. Slow recovery, but no symptom referable to ligatures.

CASE XII.—Both ovaries; two ligatures on each; abscess in left iliac fossa discharged by abdominal wound. Bowels were very irritable for a week from fifteenth day onwards. Slow but perfect recovery.

CASE XIII.—Right ovary; two ligatures; many finer ones on adhesions. Much trouble with suppuration in abdomen. Very slow recovery.

CASE XIV.—Both ovaries; right, strong silk; left, medium silk; each two ligatures. Rapid recovery; temperature never rose above 99.6 deg.

CASE XV.—Left ovary; two ligatures. Good recovery.

CASE XVI (D).—Right ovary; combination of enucleation and ligatures; four transfixions with medium silk; tumour peeled off a large fibroid uterus; abscess in right iliac region discharged by wound. Phlegmasia dolens first in left leg, then in right. Recovery perfect; went home on fifty-fifth day.

CASE XVII.—Left ovary; two ligatures. No after-trouble. Up in a fortnight.

CASE XVIII.—Right ovary; two ligatures. No after-trouble.

CASE XIX (D).—Inflamed cyst united to both ovaries and everything it was in contact with; numerous ligatures. Hopeless case. Died in seventy-two hours from suppression of urine.

CASE XX.—Right ovary; two ligatures; portions of pedicle slipped from outer one, and serious hæmorrhage occurred. I did not discover it for three hours, then found bleeding-point and secured it; but she never rallied. Died in nine hours.

CASE XXI.—Left ovary; several transfixions; laced ligature. Bowels acted on ninth day, and again on twelfth. On fourteenth day, great rise of temperature and pulse, followed by diarrhoea; pus in the evacuations. Not right again till twenty-first day.

CASE XXII.—Both ovaries; right, two ligatures; left, four; laced. No after-trouble. Went home on twentieth day.

CASE XXIII.—Left ovary; several ligatures, laced; firm pelvic adhesion also transfixed and tied in two halves. No after-trouble. Went home on twenty-first day after operation.

CASE XXIV (D).—Right ovary; laced with strong silk; several transfixions. No after-trouble. Went home on twenty-seventh day.

CASE XXV (D).—Left ovary; four ligatures. No after-trouble. Went home on twentieth day.

CASE XXVI.—Left ovary; two ligatures; many others on very extensive vascular adhesions. Nearly died from sickness, commencing on eighth day; afterwards had offensive diarrhoea. Went home well on twenty-first day.

CASE XXVII (D).—Both ovaries; three ligatures on each pedicle. A great deal of pain and trouble afterwards. Went home on twenty-second day.

CASE XXVIII (D).—Both ovaries united into one large universally adherent tumour; two ligatures on each pedicle; many on adhesion. No after-trouble. Went home on twenty-third day.

CASE XXIX.—Both ovaries; three to four laced ligatures on each. Much trouble with bowels, commencing on twelfth day; pain, depression, high temperature; pus in evacuations. Went home on twenty-second day.

CASE XXX (D).—Both ovaries; several ligatures on each. Chronic pyæmic condition followed operation; bowels very troublesome. Went home well on twenty-seventh day.

CASE XXXI (D).—Right ovary; three ligatures. No trouble referable to pedicle. Went out of hospital on eighteenth day.

CASE XXXII.—Left ovary; antiseptic operation; four thick silk ligatures soaked in carbolic oil. Much trouble with bowels from twelfth day; ended in abscess discharged *per rectum*. Went out on twenty-fifth day.

CASE XXXIII.—Right ovary; antiseptic method; two ligatures soaked in carbolic oil. Much trouble with bowels commencing before they had acted, afterwards threatening complete obstruction; gradually passed off; and patient went out on twenty-fourth day.

CASE XXXIV (D).—Left ovary; antiseptic method; two strong silk ligatures soaked in watery solution of carbolic acid (1 in 20); three fine ones in carbolic oil. No after-trouble. Went home on nineteenth day.

CASE XXXV.—Right ovary; antiseptic method; two strong silk ligatures soaked in watery solution of carbolic acid (1 in 20); partial flap operation mentioned above. No after-trouble. Bowels open on eighth day. Patient up on couch on eleventh day.

CASE XXXVI.—Right ovary; antiseptic method; five medium silk ligatures soaked in watery solution of carbolic acid; partial flap operation. No after-trouble. Went out well on seventeenth day.

CASE XXXVII.—Left ovary; antiseptic method; four medium silk ligatures (carbolicised). No after-trouble.

CASE XXXVIII.—Left ovary; antiseptic method; two fine silk ligatures (carbolicised); partial flap operation. Went out quite well on nineteenth day.

In the above cases, I have placed the letter (D) to those in which drainage was employed, as it is often an important element of success in cases treated by ligature. Other details I have purposely left out, as they are such as are common to cases treated by any method, and therefore do not affect the special question of ligature.

I may mention that I have followed the history of all my cases since they left my care, and in no case have I heard of any after-mischief from the ligature. All upon whom I have operated, who recovered, were still in good health a few weeks ago.

ON THE THERAPEUTIC USE OF IODOFORM.

By BERKELEY HILL, M.B., F.R.C.S.,

Professor of Clinical Surgery in University College; Surgeon to University College Hospital, and to the Lock Hospital.

LOCALLY, iodoform, as a dry powder, brushed lightly over the surface with a moistened camel-hair pencil, has been for three years my almost invariable treatment of venereal sores, especially the local chancre. During the last few months, I have often substituted for the dry powder an ethereal solution (one part of iodoform in six or eight of ether). The sore is touched or dabbed with a pencil dipped in the ethereal solution, according to its size and depth, lightly or copiously. The ether quickly evaporates, leaving a thin pellicle of iodoform, that as effectually stays the spread and produces healing of chancres as does the more copiously applied dry powder. Thus the surface is covered more exactly, and the disagreeable smell of the iodoform is too faint to attract attention. The sore is well washed with water and dried before the iodoform is applied, and the surface is lastly protected by a bit of dry lint. When the secretion is abundant, the dressing must be renewed twice daily, but in three or four days the amount of discharge becomes so scant that one dressing *per diem* suffices.

In this way, venereal sores heal quickly. Pain subsides at once; the sore is well in a week or ten days, and the chances of consecutive inoculation or bubo are greatly lessened. In a very few cases, the application of iodoform gives momentary smarting, which is very bearable; even the ethereal solution does not hurt, and usually the patient declares the application to be quite painless. I avoid using iodoform on inflamed sores, or on simple granulating wounds; but indolent non-specific ulcers are rapidly improved by iodoform locally applied.

Lately, I have given iodoform internally with great benefit. It acts more rapidly than potassic or other iodides, and, judging from experience thus far, is as readily borne as are those salts. I have given it in one-and-a-half-grain doses as a pill with extract of gentian. Three pills are given each day, increasing gradually till eight or ten pills are taken in twenty-four hours.

I have used it with excellent effect in cases of obstinate syphilitic ulceration of the tongue, where the dorsum is covered with rugged thickened epithelium, which is constantly splitting into deep fissures, and thus causing continual severe pain to the patient. This affection is often quite insensible to mercury, alkaline iodides, or arsenic—the remedies usually beneficial. In three of these obstinate cases, where I had been treating the patients at intervals for years with the remedies just mentioned with little lasting benefit, iodoform-pills have acted like a charm. Pain, immediately lessened, in two or three days ceased wholly; and the fissures healed rapidly, while the tongue soon shrank to its natural size. How long the relief will endure, time alone will show; but any interval of only apparent cure of this very painful affection is a great blessing to the sufferer, and time is given for the exhibi-

tion of mercury if required. In December last, I had under my care in University College Hospital a patient with ulcerated and protruding gumma of the left testis, non-ulcerating gumma of the right testis, and ulcerating gummata of the skin over the upper end of the right tibia, with other syphilitic affections. Iodoform was administered in pills, and water-dressing applied to the ulcers. Rapid healing and subsidence of the swellings took place, notwithstanding that, when the dose of eight pills *per diem* had been reached and administered for three days, an outbreak of pyrexia, coryza, and idiic acne rendered it necessary to drop the drug completely for a short time. In three weeks, the patient left the hospital almost healed, and continued his treatment as an out-patient. Again, a lady who has during the last two years consulted me occasionally for intensely agonising pain in the head caused by syphilitic pericranial and cranial disease, for which a customary dose was thirty grains of sodium iodide three times daily, was at once relieved of pain by the iodoform pill taken three times daily, though, on the third day, nausea became too urgent to allow the iodoform to be continued in that quantity; it was at first diminished till pain ceased, and then discontinued altogether. This small experience has satisfied me that in iodoform we have a very useful addition to our store of weapons for fighting syphilis. Further observation will enable us to apply it more exactly and when most suitable.

THERAPEUTIC MEMORANDA.

CHLORAL-HYDRATE IN DELIRIUM TREMENS.

A SHORT time ago, I was almost despairing of a case of delirium tremens. The man was most violent, and in a fearful state of excitement; and the remedies adopted appeared only to increase his activity and make him more and more unmanageable. The treatment had been Battley's solution in half-drachm doses; afterwards pure solution of the hydrochlorate of morphia by subcutaneous injection, as much as one grain repeated every two hours. There was no vomiting of the mixtures given on any occasion; these being, in addition to the liquor opii sedativus just mentioned, half-drachm doses of tincture of digitalis given every two hours, etc. After two or three days of the above treatment, and no improvement taking place, I determined to try the chloral-hydrate. Accordingly, at 5.10 A.M. I gave him half a drachm (thirty grains), and the same quantity at 5.40. At 6.0, he had a subcutaneous injection of half a grain of morphia. At 6.10, forty grains of chloral were given; at 6.25, two-thirds of a grain of morphia were injected; and at 6.45 he was asleep. The man slept for eight hours, and awoke without headache or other unpleasant feeling except great thirst. He was now supplied with good nourishing food (beef-tea, etc.), and he was out walking next day. The quantity of the chloral given was one hundred grains, and of morphia one grain and one-sixth, in the space of an hour and fifteen minutes. Previously to the administration of chloral, the pupils were contracted to a point: an indication, of course, that the previous mixtures had been absorbed, but, as we have seen, with the effect only of increasing the excitement. Considering that the preparations of opium given previously had not conduced to somnolency, I attribute this condition to the chloral-hydrate chiefly, if not entirely. In another obstinate case of delirium tremens, in which the usual treatment by digitalis, morphia, etc., was ineffectual, I had recourse to chloral, repeated every ten minutes till one hundred and sixty grains had been taken. The patient then fell over, and, after sleeping for seven hours, was, on awaking, altogether a changed man.

I may add that, during the first two doses, there is always increased excitement, the patient becoming very garrulous—indeed, *intoxicated*, to all appearance; but this soon gives place to thick speech, inarticulate mumbings, and peaceful sleep.

J. FARRAR, L.R.C.P.Ed., etc., Morecambe, Lancaster.

PILOCARPIN, THE ACTIVE PRINCIPLE OF JABORANDI.

A MORE elegant and effective way of introducing the active properties of jaborandi into the system than that proposed by Dr. Coghill as a therapeutical remedy in hydrophobia (*vide* BRITISH MEDICAL JOURNAL of January 5th) would be by the hypodermic injection of the alkaloid named by its discoverer, Merck, pilocarpium muriaticum (muriate of pilocarpine). It consists of transparent white crystals, and forms an easy and clear solution in equal parts of water. A hydrochloric solution of two per cent. is used for subcutaneous injection, one gramme or sixteen minims of which are equivalent to five grammes or eighty minims of the infusion of jaborandi-leaves. About one to one-