THE DIAGNOSIS AND TREATMENT OF AUDITORY-NERVE VERTIGO.

By W. R. GOWERS, M.D.,

Assistant-Physician to University College Hospital, and to the National Hospital for the Paralysed and Epileptic.

[Concluded from p. 420 of number for April 7th.]

Diagnosis from Epilepsy.—The attacks of auditory nerve vertigo resemble those of epilepsy in several particulars. This resemblance is of much significance. A sensation of vertigo often attends the slightest attacks of epilepsy. In these, however, consciousness is usually, but not always, lost, while consciousness is usually preserved during the paroxysms of auditory nerve vertigo. Severe attacks of the latter may be accompanied by sudden, transient loss of consciousness. The sufferer may even fall, and be unaware that he has fallen. A patient whom I once saw immediately after an attack had fallen a hundred yards from the place to which he was going, and had then walked to the house, but could not say how he had got there, and did not remember the fall, of which his clothes showed proof. He retched, vomited, and giddiness persisted for an hour. In this case, there had been many attacks of auditory vertigo, and there was no suspicion of epilepsy.

The common features of the two conditions are instructive. No doubt the significance of the feeling of vertigo is in each case the same. In each, it depends on an "incipient motor process" (Hughlings Jackson). In each case also, there may be a developed motor process. The violence of the movement, in severe attacks of auditory nerve vertigo, is remarkable. It is no mere compensatory action to maintain a disturbed equilibrium; the patient is dashed to the ground with force. In a case, detailed above, the patient fell on one occasion with great violence; while, on another, his grasp at some railings was insufficient to prevent him from being borne down to the ground with force. In a case mentioned by Dr. Ferrier, a patient was turned round three times and then dashed to the ground. Such movement must result from the "discharge" of a motor centre, quite analogous to the discharge which occurs in an epileptic fit. If we locate the centre for equilibration in the cerebellum, we may perhaps regard Menière's disease as a cerebellar epilepsy, reflex in character, determined by an irritation of the semicircular canals,* alone or in conjunction with impressions from some other nerves with which the centre is in connection, as those of the pneumogastric nerve. On this view, it is not difficult to understand the loss of consciousness which may accompany the severer seizures, since the connections of the cerebrum and cerebellum are intimate; fibres from almost all parts of the hemispheres being gathered into the superior cerebellar peduncles.

It has been remarked that ear-diseases may excite actual convulsions, apart from any direct affection of the brain. Such cases have been described by Mr. Hinton, Dr. Brown-Séquard, and others. Many attacks of paroxysmal vertigo present other symptoms resembling convulsive seizures, and rendering it doubtful in which category they should be placed. A premonitory sensation, apparently unconnected with the ear, may be present. In the case of Menière's disease related by Dr. Duffin, some attacks were preceded by such a sensation, which passed from the hypogastrium to the occiput. A case, presenting very interesting symptoms of this character, has been lately under the care of Dr. Ringer in University College Hospital. The patient, a man about forty years of age, was bitten by a dog, about three months before admission, in the back of the right leg, below the calf. The wound was cauterised and healed, but a few weeks afterwards became painful; and one day a sudden pain seemed to start from it, pass up the leg and side of the trunk to the right side of the head behind the ear, and then his head seemed to go round and round "like the wheel of a steam-engine", in what direction he could not say. He had many slighter attacks, and, after the giddiness had subsided, a pain seemed to pass up the leg and down the arm to the palm, backwards and forwards. He was conscious of no auditory defect; but it was found on examination that, while the hearing on the left side was in every way normal, that on the right side was distinctly defective. A watch was not heard so readily opposite the ear as on the left side, and whereas its sound on the left side was much increased in intensity by contact with the

skull, it was not heard at all when in contact with the skull (zygoma or mastoid process) on the right side. There was also some persistent defective equilibrium: in walking, he experienced a constant tendency to go to the left; but he explained this by describing it as the corrective of a tendency to go to the right. The probable explanation of this case I think to be that the equilibrium centre, being disturbed by the morbid state of the auditory apparatus and nerve, was further perverted by the irritation from the bite, which developed sudden overaction in the centre, an "irradiating" action of the sensory centre being set up at the same time.

If the paroxysmal auditory nerve vertigo be thus due to an epileptoid process, it is to be expected that the condition of instability of the nerve-cells, such as we recognise as an important element in the pathology of epilepsy, may influence in some cases the occurrence of this affection. The only way in which such a condition of nerve instability could be recognised as a factor in Menière's disease would be by the coexistence of epilepsy and auditory nerve vertigo. Of course, much care and even hesitation is needed in diagnosing auditory nerve vertigo in the presence of epilepsy; but in two cases, at present under my care, there is, I think, clear evidence of the coexistence of the two maladies.

One of them, a married woman aged 65, began to suffer from epileptic fits at the climacteric period, at about 47; the fits continued for six or seven years, and then ceased. All the fits were the same in character; they were preceded for a day or two by slight fainting attacks, and a slight feeling of faintness was the immediate warning of some of the fits. During the attack, consciousness was lost, and there was slight jerking. She slept afterwards, and on waking had occipital headache. For the last twelve years, she has had no fits. During the last two years she has suffered from vertigo, paroxysmal; the attacks sometimes come on when sitting still, but are more often induced by a sudden movement, especially by looking up. The sensation during an attack is that of falling backwards and towards the right. It always comes on sufficiently deliberately to enable her to take hold of some object and save herself from falling. She never loses consciousness. She frequently vomits with the attacks, and refers them to "biliousness". She was conscious of no defect in hearing; a watch is heard on each side at nine inches from the head. A watch, pressed against the left parietal eminence, is heard slightly, but on closing the ear she ceases to hear it. A watch in contact with the right parietal eminence is not heard at all, with the ear open or closed. (The meatus is free from obstruction.) On each side, a watch in contact with the malar bone, and shielded by the hand from the ear, is unheard. A tuning-fork on each side, in contact with the parietal eminence, is heard, but less loudly when the meatus is closed than when it is open. The effect of movement on the giddiness was tested: a sudden jerk of the head backwards brought on an attack; a sudden rotation of the head to the right also brought on an attack, while a sudden rotation to the left had scarcely any effect. During the six weeks before she came under treatment, she had two more fits, of the same character as the earlier ones. Since the recurrence of the fits, the giddiness has been more troublesome.

In the other case, a girl has suffered from several epileptiform convulsions, and has almost constant vertigo, varied by severe paroxysms on movement. The giddiness is worst after rising in the morning. The sensation is of falling backwards and to the left. She has constant noises in the ears, and is generally, though not always, unable to hear a watch in contact with the head. Occasionally, a watch is heard faintly when in contact with the mastoid process. This patient suffered at night from twitching of the right limbs. A similar symptom existed in the case of the man whose strange postures during sleep have been described, and indicates a tendency on the part of the motor centres to spontaneous discharge. She is also troubled with startings in sleep, as were several other patients whose cases have been detailed. It may be suggested that, in this case, the epileptiform convulsions are really excited by the aural affection. But such an explanation will not apply to the other case mentioned; and I believe that convulsions have not been described as the consequence of anything but a severe ear-disease.

It will be seen from these remarks that the diagnosis from epilepsy must turn, as in the case of gastric disturbance, to some extent on the detection of defective audition, and especially on defect in the power of hearing vibrations conducted through the skull. The definite character of the vertigo is in these cases no criterion. A more important symptom is its persistence between the paroxysms and its long duration in the paroxysms themselves. Its relation to movement is also important. If it be related to change of posture, and can be produced by certain sudden movements of the head, it is probably due to auditory nerve disturbance. In the latter, the attacks are succeeded by

* M. Pierret, in a recent communication on the origin of the auditory nerve (as Dr. Hughlings Jackson has lately mentioned), appears to doubt whether its fibres can be traced with any certainty so far as the cerebellum. The direct evidence contained in Dr. Lockhart Clarke's paper in the *Philosophical Transactions* for 1868 is, however, very convincing, especially his comparative investigations on animals. Indirect evidence is afforded by the fact which he points out, and which I have repeatedly verified, that a large number of the fibres of both divisions of the auditory nerve (the anterior especially) pass into the substance of the restiform body and mingle with its fibres, and the restiform body passes entirely to the cerebellum.

vomiting, much more frequently than are the attacks of *petit mal*. Lastly, there is rarely the objective evidence of momentary loss of consciousness, which is so frequent in minor epilepsy, even in the slightest attack; in auditory nerve vertigo, consciousness is only lost or obscured in severe attacks, in which the giddiness persists for a considerable time and vomiting frequent.

Production of Auditory Nerve Vertigo.—Before speaking of the treatment of this affection, it is worth remark that auditory nerve vertigo can be produced artificially. Quinine produces a sense of confusion with tinnitus; but very definite symptoms may be caused by salicylate of soda. This was shown very strikingly in the case of a patient lately under treatment for acute rheumatism in University College Hospital (under the care of Sir William Jenner). The patient was a woman aged 40, whose hearing was supposed to be unimpaired. She was not subject to giddiness. It was her first attack of acute rheumatism, and there was no cardiac affection. On January 26th, salicylate of soda was commenced in doses of twenty-five grains every three hours. On the 28th, she complained of noises in ears, deafness, and giddiness, which the next day had increased so much that the salicylate was omitted. The following day the giddiness was much less, and on the 31st had almost gone. On February 6th, the same dose was resumed; on the 7th, the same symptoms were complained of. The noises in the ears were constant; a watch was heard only at two inches distant from each ear, and was not heard at all on either side when in firm contact with either the zygoma or mastoid process. A tuning-fork on the vertex was heard fairly well, but the sound was not increased by closing the ears. The giddiness was slight and indeterminate as long as she lay still, but was very considerable and definite when she raised her head or sat up. Objects before her all seemed moving to the right. On the 8th, these symptoms continued, and the salicylate was discontinued. On the 10th, the giddiness was gone, and she could hear the watch at a distance of six inches from either ear, and could hear it, although faintly, in contact with the zygoma or mastoid process, but not when in contact with the parietal eminence. On the 23rd, the salicylate was resumed, and, eighteen hours after its resumption, deafness and giddiness had returned, which again ceased a day or two after the discontinuance of the drug. When the patient was convalescent, a careful examination of the state of hearing revealed very little abnormality, the only difference being that the watch in contact with the skull was not quite so distinct on the right side as on the left. In another case, I have seen similar symptoms of deafness and definite vertigo produced by salicylic acid.

Treatment.—The direct treatment of labyrinthine affections is a subject for the special aural surgeon. I would only here indicate that, when there is evidence of an irritative process, blistering behind the ear sometimes affords very marked relief to the vertigo. Occasionally, there is evidence of a constitutional condition on which the aural disturbance is dependent. In some cases the patient is gouty, and there is reason to believe that a gouty change in the membranous labyrinth is the cause of the morbid action. This was the case in one of Mr. Hinton's patients, and also in one case which has come under my own observation. Marked relief is afforded under these circumstances by colchicum and potash. In rare cases, the vertigo depends on a syphilitic inflammation of the labyrinth. Such a case has been lately described by Moos of Heidelberg.* One of the most severe forms of paroxysmal vertigo I have seen was in a child, with almost complete deafness and every sign of inherited syphilis. In all syphilitic cases, the specific treatment is, of course, required.

It has been just remarked that certain drugs have a marked influence on the organ or nerve of hearing. The effect of quinine induced Charcot to employ it, in full doses, in a case of auditory nerve vertigo, with some beneficial result. I have tried it, but have not found any marked effect was produced on the vertigo. The influence of salicylate of soda upon the equilibrium, which I have described, suggested its use in this disease. Equilibrium is maintained by the balance of opposing impressions; its overthrow is the result of the loss of that balance. In our ignorance of the way in which salicylate produces the disturbance, it is conceivable that it may, in some cases, have such an influence as to counteract the morbid action, and lessen the disturbance of equilibrium. This it has seemed to do in one or two cases in which I have tried it. It does not remove the giddiness, but in some cases lessens its intensity. It has been given in doses of from five to ten grains three times a day. The patient with gastric ulcer, whose case has been narrated above, thought that she was better while taking the salicylate than when taking any other

medicine. Its effect, unfortunately, seems after a time to become less.

One of the objects of this paper is to show that, in auditory nerve vertigo, other morbid conditions may concur with the aural condition in causing the vertigo. The knowledge of this will, in many cases, enable much relief to be given. The gastric disturbances are to a considerable extent under control, and, by their removal, the attacks of vertigo which they excite, although ultimately dependent on ear-disease, may be much lessened in frequency and intensity. The most unremitting care in diet and regimen are, however, necessary. An actual attack, which has been produced by gastric irritation, may often be arrested by a good dose of an antacid.

Lastly, any undue sensitiveness of the grey matter of the equilibrium centre must be lessened. For this purpose, no remedies are so useful as the bromides of potassium and ammonium, and several observers have noted their utility in Menière's disease. It is probable, indeed, that bromide does more good in this condition than any other single medicinal agent. This fact affords another point of resemblance between epilepsy and paroxysmal vertigo. The effect of the bromide is in some instances increased by the addition to it of some other sedative, as belladonna. Other sedatives alone—opium, Indian hemp, gelsemium, hyoscyamus—have seemed to me without effect.

Auditory nerve vertigo is unquestionably, in many cases, a very obstinate affection; but, when the several factors in the individual cases are sought out, and as far as possible corrected, a considerable amount of relief may be afforded. In several cases that I have seen, immunity for long periods has been obtained. The patient, whose severe attacks were described at page 419, has had no severe attack since the commencement of the treatment, now a year ago. Occasionally, he notices slight unsteady feelings, "as if the ground were being tilted up a little". When this is the case, he takes a few doses of bromide of ammonium, carbonate of bismuth, and belladonna, and the symptom always vanishes. In another case, in which there were other indications of an irritable nerve-centre, bromide and bark produced an almost complete cessation of the attacks; and the improvement was subsequently maintained on a course of quinine only, although quinine had at first failed altogether to relieve.

It is on a recognition of the associations of auditory nerve vertigo, and their place in the pathology of the affection, that its treatment can be most effectually based.

CASE IN WHICH MOBILITY AND A PERFECTLY USEFUL LIMB RESULTED AFTER EXCISION OF THE KNEE-JOINT.

By THOMAS ANNANDALE, F.R.S.E.,

Surgeon to the Edinburgh Royal Infirmary, and Lecturer on Clinical Surgery.

At a meeting of the Clinical Society of London in May of last year, Mr. Barwell showed a girl whose knee-joint had been excised three years before by Mr. Cowell. In this case, a "flail" joint had resulted, but the limb was a useful one. Having lately had an opportunity of examining a patient whose knee-joint I excised almost five years ago, I found in her case a condition similar to, but more perfect than, that described in connection with Mr. Cowell's patient. As such a result after excision of the knee is rare, I have thought my case worthy of a special note and illustration.

The interesting result in this case was quite accidental, and I can give no special reason for its occurrence either in regard to the operation itself or the after-treatment, both of which were carried out according to the usual principles. In fact, I at first looked upon the case as one of failure to obtain proper union after excision of the knee-joint; and it was only on examining the patient some years after the operation that I discovered with pleasure and surprise that my case was, instead of a failure, a new and remarkable success, in so far that a movable new joint and perfectly useful limb had resulted. An important fact in the case is, that the patient had the power of flexion and extension, and the new articulation was by no means a "flail" one.

Agnes M., aged 10, was admitted into my wards, suffering from disease of the knee-joint, on October 25th, 1872. Excision of the knee was performed on November 13th, 1872, by means of a semilunar incision across the front of the joint. The articular surface of the femur, a thin slice of the tibia, together with its articular surface and the patella, were removed; care being taken not to interfere with the epiphyseal line of the former. The patient was dismissed with the wound healed on June 22nd, 1873; but osseous union had not taken

* Virchow, *Archiv*, vol. lxi, part ii, p. 313. The patient suffered from attacks of giddiness and noises in the ears; and, later, rapidly increasing deafness. *Post mortem*, there was found a small-celled infiltration of the membranous labyrinth, and especially of the ampullæ.