This inhaler has been made by Messrs. James Coxeter and Son, Grafton Street East, London, and they deserve my best thanks for the efficient manner in which they have carried out my directions as regards its manufacture, besides bringing it out at a very moderate price.

LESION OF THE SYMPATHETIC ON ONE SIDE OF THE HEAD.

By FRANCIS WARNER, M.D.Lond., M.R.C.P.,

Assistant-Physician to the East London Children's Hospital; late extra Acting Physician to the Children's Hospital, Birmingham; etc.

DR. BURNEY YEO'S interesting case of one-sided development of exophthalmos, and the discussion thereupon at the Clinical Society, suggested to me to publish the following case, where a lesion of the vaso-motor system, probably intracranial, produced excessive excitability on one side of the head.

Elizabeth R. came under my care on February 26th, 1876, as an out-patient at the Children's Hospital, Birmingham. The mother, a very intelligent woman, complained that the girl was subject to "fainting attacks and some cough". The patient was a healthy looking girl, rather pale, aged 11. She did not bear the aspect of the nervous temperament, but the mother considered her "rather nervous". When questioned, the girl complained of "headaches" and also a "feeling of numbness at the top of the head". The mother then said that she had noticed that, on excitement, *the right side of the face flushed up to the middle line* much more than the left side. The neck had not been noticed to flush. The following history was obtained.

The girl was apparently healthy and well till aged six years. Then, one day, after a hearty dinner, she went upstairs, and, on returning in a few minutes, was *suddenly convulsed*; she had general twitchings all over the body, no loss of consciousness; she knew all that was passing, and remembered it now. This convulsion lasted an hour. After this attack, there was no drowsiness, no vomiting, but she was fatigued. She was kept quiet a few days, but nothing more happened. In two or three days, the mother noticed the frequent flushing of the right side of the face, the left remaining pale. This she pointed out to the surgeon attending, and she had observed the same tendency ever since. She never had any convulsions but this.

She never had any "fainting attacks" previously, but since had been liable as follows. She was at times unable to speak; sometimes could not open her eyes; never struggled; the hands remained warm; she never bit her tongue. Before these attacks, the mother thought she was more liable to flushing of the right face.

On examination, when looking straight forward, the right pupil was widely dilated, the left of moderate size. The iris of the left eye was light brown, in the right uniformly much darker in colour. The mother had noticed this change of colour as something that had happened during the last few years. Both pupils reacted to light, but the right was much more sluggish; there were no signs of iritis; each eye reacted normally to atropine and Calabar bean. The power of accommodation was distinctly less in range in the right eye as compared with the left, but was not much impaired. There was no affection of any branches of the motor oculi nerve. The fundus was healthy and the circulation was equal in each eye. In these ophthalmic details, I had the advantage of co-operation with Mr. Lloyd Owen.

While talking to and examining the child, the right side of the face and ear distinctly flushed more than the left; and this difference was observable to the median line of the nose and forehead; to touch also the right was warmer than the left. There was no perceptible difference in the tongue on the two sides. The right hand felt warmer than the left. Both radial and carotid pulses were distinctly smaller to the touch on the right than on the left side, but the amount of this difference varied on different occasions.

There was slight drooping of the left eyelid; this the mother had always noticed in the child. There was no enlargement of the spleen. The thyroid gland was normal; the heart was healthy in every respect and acting well; the urine was normal.

The patient remained under observation about three months without undergoing any change to be noted. She was able to run about, but had occasional attacks as follows. One morning, her mother found her in bed not asleep, but apparently sleepy; she was unable to speak or move for some minutes, and the eyelids twitched much; when she got up, she complained of feeling giddy. Such attacks in the daytime never caused her to fall down.

As to treatment, she took bromides and belladonna, at other times

tonics, without any apparent effect. There appears here to have been a localised lesion of the sympathetic centre, a greater susceptibility of excitement in one set of vaso-motor nerves, and this lesion appears to have been a sequel of a series of convulsions six years before the time of observation.

THERAPEUTIC MEMORANDA.

TINCTURE OF ACONITE.

MR. HERBERT JONES has, in the BRITISH MEDICAL JOURNAL of March 3rd, 1877, published an interesting case bearing upon the use of the above tincture. I would supplement his valuable remarks by a few observations. "It is interesting to observe that the effect of this drug differs. It has been known to contract the pupil. In my case, the pupils remained widely dilated for nearly three days, showing that its power over the nervous system varies"—are his words. As he here stops short, I presume to offer a partial explanation of the varying conditions of the pupil. By the case which he cites, the drug was taken in one enormous quantity, and, as a consequence, *extreme symptoms* and almost mortal depression supervened. Of course, the pupil was widely dilated; not, I apprehend, from any specific agency of the medicine, but from its sudden depressant action, almost overwhelming. Now the same exactly holds good of opium and morphia, which in small doses will produce contraction of the pupil—a contraction that will continue even to death if the person be poisoned slowly; whereas, if by a suddenly overpowering large dose, the entire system is depressed, and, along with other manifestations of this condition, the dilated pupil obtains.

In April 1876, I had under my care in the hospital a girl aged 23, an epileptic, who showed an astonishing resistance to the power of aconite tincture, though prepared by one of the first chemists in London, and dispensed by our very careful dispenser. Here is the prescription:—℞ Tincturæ aconiti ℥ viii; potassii bromidi ℥ xii; syrupi ℥ iv; aquæ ad ℥ vi. Sumat ℥ iv ter die in aquâ. An aconite suppository (half a grain) was ordered to be used every night at bedtime. No aconitism was produced, though this treatment was continued for weeks; nor was the epilepsy cured.

G. DE GORREQUER GRIFFITH, Senior Physician to the Hospital for Women and Children, etc.

TREATMENT OF RINGWORM BY CHRYSOPHANIC ACID.

HAVING only just seen the correspondence in the BRITISH MEDICAL JOURNAL on the above subject, I beg to state that I have had large opportunities of studying the treatment of Burmese ringworm, sometimes called "Malabar itch" (which, I presume, is the form of the disease alluded to by Dr. Foulis). Having served for some time in Rangoon, where it is remarkably prevalent at certain seasons, I there not only treated numerous patients suffering from it, but also had a most obstinate and troublesome attack of the disease myself.

I have only found one *certain* remedy, that being Goa powder. Cassia alata I have found useful in some cases, more particularly in the early stages, where the disease is circumscribed; but in other cases it is practically useless, and moreover possesses the disadvantage of losing its specific action during the rains, the very season when most required.

With regard to chrysophanic acid, I have never had an opportunity of using it by itself; but, as I believe that it has never been accurately determined whether Burmese ringworm is the effect of a vegetable or an animal parasite, I am unable to say how it acts (if it be the active ingredient) in Goa powder. Burmese ringworm has no marked effect on the hairs of the affected locality. In some cases where the powder was not procurable, I have found tincture of iodine, and also a fifteen to twenty grain solution of nitrate of silver, of use. Burmese ringworm appears to affect men more than women or children, as I can only call to mind one woman among many male cases in the course of two wet seasons, and have never seen children affected. It is quite non-contagious, and apparently is communicated by the bath-water (at that time of year very unclean, and containing large quantities of vegetable matter in suspension), acting on a skin already rendered prone to irritation by being constantly bathed in the profuse perspiration of a hot, damp, and relaxing climate.

C. J. L. BUSHE, M.B., Army Medical Department, Malleipooram.