THE LUMLEIAN LECTURES ON THE MUSCULAR ARTERIOLES:

THEIR STRUCTURE AND FUNCTION IN HEALTH
AND IN CERTAIN MORBID STATES.

Delivered at the Royal College of Physicians of London.

By GEORGE JOHNSON, M.D., F.R.S.,

Fellow of the Royal College of Physicians; Professor of Clinical Medicine and Senior Physician to King's College Hospital; etc.

LECTURE III.

Renal Asthma: Symptoms and Proximate Cause.—The Dry and Inactive Skin of Bright's Disease: its Proximate Cause.—Is Diuresis a Result of High Arterial Tension?—Is High Arterial Tension a Cause of Albuminuria?—Uræmic Convulsions and other Nervous Symptoms: their Relation to Epilepsy: their Proximate Cause: Principles of Treatment.—Two Forms of Impaired Vision in Bright's Disease: the Proximate Cause of each.—Cholera Collapse and Pulmonary Embolism compared.—Cholera Collapse and Spasmodic Asthma compared.—Conclusion.

THERE yet remain for consideration certain symptoms and results of renal disease which are more or less intimately associated with and dependent upon those impediments and derangements of the circulation to which I directed special attention in my last lecture; and amongst these symptoms there is a peculiar form of dyspnoea, or, as it is sometimes called, *renal asthma*. During the progress of the various forms of renal degeneration which are commonly included under the generic term Bright's disease, a more or less persistent dyspnoea may result from a variety of causes, such as anæmia with a deficiency of the oxygen-carrying hæmatine; bronchitis or oedema of the lungs; fluid in the pleura, or in the pericardium, or in the abdomen; an enfeebled condition of the heart, either with or without valvular disease. But the dyspnoea to which I now refer is of a different character. It is not persistent, but occasional and intermittent, coming on in paroxysms, in some cases soon after food has been taken, and especially after an evening meal; while in other instances it occurs mainly during the night, when it entirely prevents or greatly disturbs the patient's sleep. In some cases, the attack resembles one of spasmodic asthma, and there are loud sibilant *râles* over the lungs, apparently the result of bronchial spasm; but in the more typical cases the phenomena are quite different. The breathing is hurried and laborious, the heart's action is rapid and feeble, and there is more or less lividity of the lips and of the general surface; yet, on auscultation, loud puerile respiration, unmingled with *râles* or crepitation, may be heard over the whole of both lungs; while in other cases, especially after a long continuance of the dyspnoea, fine crepitation may be heard over the bases of the lungs, but there is obviously no deficiency of respired air, nor any change in the physical condition of the air-passages to explain the distressing symptoms. It can scarcely be doubted that the symptoms in question are the result of a form of uræmia; and they occur with especial frequency in the advanced stages of the contracted granular kidney, with which other forms of uræmic poisoning are so commonly associated.

The question then arises, What is the proximate cause of this uræmic dyspnoea? We know that dyspnoea—a distressing sense of suffocation—arises when, from any cause, the free intermingling of the blood with the respired air is prevented. The respiratory process may be interfered with alike either by an obstruction in the air-tubes hindering the access of air to the pulmonary cells, or by some impediment to the onward flow of blood through the pulmonary capillaries. A plug in the pulmonary artery causes a sense of suffocation as urgent and distressing as an obstruction in the larynx, trachea, or bronchial tubes. I have on several occasions had the opportunity of observing the symptoms which result from an embolus or a thrombus in the pulmonary artery. When the obstruction of the artery is sudden and complete, there is usually a combination of pallor with lividity of the surface, the result of systemic venous fulness, epileptiform convulsions consequent on sudden extreme arterial anæmia of the brain, and death within a few seconds or minutes from arrest of the circulation. When the obstruction of the pulmonary artery is less complete, but permanent, and sufficient to destroy life, the respiration is hurried and laborious, and attended with a distressing feeling of want of breath. One poor girl, who was dying from this cause, panted out the statement, "I feel as if

I could not get breath enough"; yet over the whole of both lungs, except over two limited patches where some blood had become impacted in the terminal bronchi, there were unimpaired resonance and loud puerile respiration. The surface of the body in these cases becomes cold and pallid, with more or less lividity of the lips; and the pulse is small and feeble, or even quite imperceptible.

It is obvious that the blood-current through the lungs is the channel by which oxygen reaches the tissues; and, when that current is obstructed, the sense of breathlessness is the result of a call from the tissues for the indispensable vitalising oxygen.

Now, it has occurred to me that the paroxysmal dyspnoea of Bright's disease may be, in part at least, explained by an interrupted circulation through the pulmonary capillaries occasioned by spasm of the pulmonary arterioles, which are stimulated to this excessive contraction by the influence of the impure blood upon the vaso-motor nerves and centre. There are some facts and arguments which support this theory of the pathology of renal asthma.

The panting laborious mode of breathing and the sense of breathlessness are very like the symptoms which result from a plug narrowing but not completely obstructing the pulmonary artery. A consideration of the *juvantia* and *ledentia* lends support to the theory of arterial spasm of the lungs. In some cases, the dyspnoea is speedily and greatly relieved for the time by inhaling the vapour of nitrite of amyl; and this relief is probably due to the well-known influence of this agent in relaxing the muscular arterioles. In many instances, relief is afforded by moderate doses of chloral-hydrate, either alone or in combination with bromide of potassium. Ten grains of each of these compounds may be given twice or thrice in the twenty-four hours. This combination probably acts by lessening the reflex excitability of the vaso-motor centre. If, however, the secretion of urine be very scanty, and the blood consequently much contaminated by retained excreta, the chloral may fail to relieve, and it may even add to the distress. In this condition, the breathing is often most effectually relieved by means directed towards the purification of the blood by promoting the free action of the skin and bowels, while the diet is carefully regulated both as regards quantity and quality, an exclusive milk-diet being often the most suitable. The dyspnoea is almost invariably made worse by opium in any form; and the probable explanation of this is, that opiates, by checking the secretions of the bowels and kidneys, increase the blood-contamination, and thus intensify the exciting cause of the dyspnoea.

My late distinguished friend and colleague Sir William Fergusson often discussed with Sir George Burrows and me the etiology and treatment of this distressing symptom, from which he himself suffered severely during the last few weeks of his life. He asked me if I could explain the dyspnoea, and I suggested to him the explanation which I have just now given. He said that my explanation appeared to be in accordance with his own sensation, which was that of an obstruction somewhere within the chest; and, as the obstruction was evidently not in the air-passages or in any of the heart's orifices, it seemed probable that it might be in the pulmonary arteries. He several times described to me an interesting observation which he had repeatedly made upon himself. He found that, by voluntarily increasing to the utmost the respiratory movements—that is, making forcible inspirations and expirations fifteen or twenty times in succession—he could for a time free himself almost entirely from the distressing sensation of breathlessness. Then gradually, in the course of a minute or two after the cessation of the forced respirations, the former feeling of want of air returned; and he explained the temporary relief by suggesting that the deep inspirations had the effect of sending on a larger volume of blood highly charged with air.

The theory of uræmic dyspnoea which I have here set forth receives some confirmation from one fact which I have frequently observed. I allude to the fact that, when the dyspnoea has been long continued or has frequently recurred, although the respiratory sounds may have been quite pure and free from crepitating *râles* at the commencement, yet sooner or later moist wheezing sounds may be heard over the bases of the lungs at the back. The explanation which I would suggest is the following. In consequence of the impeded passage of blood through the minute pulmonary arteries, there is an unusual distension of the right cavities of the heart and of the systemic venous trunks. The bronchial veins, which open through the vena azygos and the superior intercostal vein into the superior cava, participate in this distension; and, in consequence of the engorgement of the bronchial veins and capillaries, there occurs a passive serous exudation into the bronchial tubes, which, gravitating towards the bases of the lungs, gives rise to the moist crepitating sounds which are heard there. A similar engorgement of the bronchial veins and capillaries occurs as a result of the impeded pulmonary circulation consequent on the various

forms of apnoea, whether the cause of the apnoea be the bronchial spasm of spasmodic asthma, obstruction of the larynx by diphtheritic false membrane, compression of one lung by a rapid and copious effusion, or extensive pneumonic consolidation of one lung. In a case of rapidly fatal laryngeal diphtheria, the contrast between the extreme anæmia of the pulmonary capillaries and the excessive capillary and venous engorgement of the bronchial mucous membrane is very striking. The explanation is obvious. In proportion to the exclusion of air from the lungs, the pulmonary arterioles contract and the capillaries are anæmic, and in the same proportion the blood accumulates in the trunks of the pulmonary artery, in the right cavities of the heart, and in the systemic veins, including the bronchial veins and capillaries; and so the bronchial mucous membrane is congested. The dark colour of the lungs noted in some cases of sudden apnoea—as, for instance, in Dr. Massey's case of hanging quoted in my first lecture—is explained by the engorgement of the bronchial veins and capillaries consequent on the arrested circulation through the pulmonary arterioles.

The hæmoptysis which sometimes follows a severe and prolonged attack of spasmodic asthma is probably explained by rupture of the overgorged bronchial capillaries consequent on the obstruction in the pulmonary arterioles which results from the bronchial spasm. The order of events is, first, bronchial spasm, with resulting partial apnoea; second, contraction of the pulmonary arterioles, with a backward engorgement extending to the bronchial veins and capillaries, which may relieve themselves by a sero-mucous exudation or by actual rupture and hæmorrhage.

Another phenomenon closely allied to this is the oedema of the sound lung, which occurs not unfrequently when one lung has been suddenly consolidated by pneumonia or compressed by a rapid and copious pleuritic effusion. In such a case, moist crepitating sounds may often be heard in the bronchi of the healthy lung: a condition of things which must always be looked upon with anxiety, because an increase of this oedematous effusion into the bronchi may cause a fatal apnoea. There can, I think, be no doubt that the bronchial, and not the pulmonary, vessels are the source of this serous exudation. In consequence of the impervious condition of the inflamed lung, there is a state of partial apnoea; more blood is sent to the sound lung than can be fully aerated; its progress is checked and regulated by the stop-cock action of the pulmonary arterioles; the right cavities of the heart and the systemic veins are engorged; and with this there is bronchial venous and capillary congestion, and a consequent passive serous exudation into the bronchi. In the condition here described, a timely venesection may sometimes rescue the patient from impending suffocation. The abstraction of blood relieves the distension of the veins and the right cavities of the heart, and at the same time it lessens the work of the lung in proportion as the volume of blood to be aerated is diminished.

A case of tricuspid regurgitation, which came under my care some years ago at the hospital, affords an instructive illustration of the effect of an impediment originating at the right side of the heart acting backwards, first upon the bronchial, and secondly upon the pulmonary, circulation. A woman, aged 52, was admitted with general dropsy, albuminuria, a systolic bellows-sound at the bottom of the sternum (*i.e.*, over the tricuspid valve), distended and pulsating jugulars, and the physical signs of bronchitis. A single dose of elaterium, acting very freely upon the bowels, removed at once the dropsy, the albuminuria, and the bronchial râles. After a few days, however, all the symptoms returned, and the patient soon died. We found, as we had expected, dilatation of the tricuspid orifice so great, that the valve was incompetent to close it; while all the other valves and orifices were normal. The lungs were much engorged. In this case, it is evident that the primary cause of all the symptoms was incompetence of the tricuspid valve. There was consequently reflux of blood into the systemic veins, renal congestion and albuminuria, anasarca, bronchial venous and capillary congestion with a muco-serous exudation into the bronchi, and then a secondary obstruction in the pulmonary capillaries consequent on the pressure of the gravitating bronchial secretion.

It is interesting to note that, on one occasion before her admission, she had spat some blood, the source of which was, in all probability, the overgorged bronchial capillaries. We are familiar enough with hæmoptysis as a result of mitral constriction or incompetence, the source of such hæmoptysis being, in most instances, the pulmonary vessels; but disease on the right side of the heart obviously throws back the strain and pressure primarily and chiefly upon the bronchial vessels.

One of the most constant symptoms and results of chronic Bright's disease is a remarkably dry, harsh, and inactive state of the skin; a

state of skin which, in many instances, resists even the diaphoretic influence of the hot-air or the Turkish bath. I formerly supposed, and publicly expressed my belief, that this condition of skin might be due to the fact that the excessive contraction of the hypertrophied subcutaneous arterioles resists the relaxing effect of external warmth which has so powerful a diaphoretic influence upon the healthy skin. My belief was that the deficient cutaneous secretion was a result of a defective blood-supply to the sweat-glands. I take this opportunity of saying that, for some time past, I have abandoned this theory as unsatisfactory and erroneous. It must be borne in mind that the excessive power of resistance possessed by the hypertrophied arterioles is to a great extent, if not entirely, compensated by the increased injecting force of the hypertrophied left ventricle. The inactivity of the skin, therefore, is not to be explained by a defective blood-supply. I now believe that the dry and inactive state of the skin, which is commonly associated with chronic renal disease, is to be sought for in a structurally diseased condition of the sweat-glands themselves. With the able assistance of my colleague Professor Gerald Yeo, I hope ere long to be able to demonstrate as a fact what is now only a plausible hypothesis, but one which is rendered probable by some well ascertained facts. We know that disease of one excretory organ frequently induces disease in one or more allied organs. For example, when, in consequence of obstruction of the gall-duct, or some structural disease of the liver, bile accumulates in the blood, the products of that secretion are vicariously eliminated by the kidneys, and the result is that the secretory tissues of the kidney undergo structural changes. The urine is not only deeply tinged with bile, but it is often found to contain desquamated renal epithelium, tube-casts, and sometimes albumen. The excretion of biliary materials excites a mild form of desquamative nephritis.

It is notorious that suppressed action of the skin by exposure to cold and wet is a very common cause of acute renal disease with albuminuria. The most probable explanation is that the products of the cutaneous secretion, being retained in the blood, are partially excreted by the kidney, and excite structural changes there, as bile-products have been shown to do. There is a very close relationship and interdependence between the cutaneous and the renal secretion. Many of the symptoms of uræmic poisoning may be removed or mitigated if by any means we can promote and maintain a free action of the skin, the prolonged application of the wet pack being one of the most efficacious of these means. Now, there is reason to believe that when, in consequence of degeneration of the kidney, urinary materials accumulate in the blood and are partly excreted by the skin, as they have been proved to be, this vicarious excretion will induce structural changes in the sweat-glands, and thus explain the abnormal dryness of the skin. This, which is at present a theory, I expect soon to see as a demonstrated fact; and if any one or more workers will take up this subject and complete the demonstration before Dr. Yeo and myself have had the opportunity of doing it, we shall be very happy to be thus anticipated.

Another symptom, which some writers on renal pathology have, as I think erroneously, attributed to increased vascular tension, is the copious secretion of urine which is often associated with the small granular kidney. It is possible, though by no means certain, that increased pressure on the capillaries of the kidney might cause a more copious secretion of urine; but, in the granular kidney, the firm contraction of the hypertrophied renal arterioles counteracts the injecting force of the strong left ventricle, and thus prevents an increased afflux of blood into the capillaries of the kidney. There is no reason to suppose that high arterial tension has any direct tendency to cause an increased secretion of urine. In cases of contracted granular kidney, the two conditions are associated, but in the early stages of the lardaceous kidney the copious secretion of urine occurs without arterial tension. It is probable that in both classes of cases the copious flow of urine is caused by the diuretic influence upon the kidney of some abnormal products in the circulation; an influence analogous to that of sugar in cases of diabetes.

Again, it has been supposed by some that vascular tension may be a direct physical cause of albuminuria. But if, as we have good reason for believing, the increased vascular tension is a result of resistance caused by contraction of the arterioles, it is evident that, just in proportion as the renal arterioles contract and resist the blood-stream, the pressure in the capillaries in front of them will be lessened. And it is a matter of daily experience that in those cases of chronic renal disease which are commonly associated with the highest degree of arterial tension—cases of small granular kidney—the amount of albumen in the urine is least, and not infrequently there is none even, in the advanced stages of the disease.

Relaxation of the renal arterioles is more likely to cause albuminuria than their excessive contraction; and the late Dr. Warburton Begbie published some cases in which he believed that the urine became albuminous in consequence of vaso-motor paralysis, resulting in an increased afflux of blood to the Malpighian capillaries.

This view, too, is confirmed by the results of experiments. Vulpian (vol. i, p. 534) describes the effect of dividing the left splanchnic nerve of a dog just above the kidney. The renal arterioles being paralysed, the kidney at once becomes congested, red, and swollen, the quantity of urine excreted by that kidney is increased, and it becomes highly albuminous. The galvanisation of the distal end of the divided nerve stimulates the renal arterioles to contract, and the red and congested kidney gradually becomes pale.

Contraction of the systemic arterioles in general, with relaxation of the renal arterioles, might cause capillary renal congestion and albuminuria; but this is not the condition which exists in cases of chronic albuminuria with high arterial tension. On the contrary, there is evidence that the renal arterioles in these cases are more firmly contracted and more decidedly hypertrophied than those of any other organ or tissue.

An impediment to the circulation causing albuminuria must act in front of the Malpighian capillaries, so as to cause a backward engorgement and a transudation of serum through their walls into the uriniferous tubes. Thus, Dr. George Robinson (*Med.-Chir. Trans.*, vol. xxvi, p. 51) proved many years ago that a ligature on the renal vein of a living rabbit renders the urine albuminous, and even bloody. This experiment has been repeated by Frerichs and others with the same result. Bowman's demonstration of the structure of the Malpighian bodies and their connection with the uriniferous tubes rendered these results quite intelligible. In like manner, distension of the convoluted tubes by morbid exudation within them may, by compressing the intertubular capillaries, cause engorgement of the Malpighian capillaries and escape of serum through their walls.

[To be concluded.]

CLINICAL LECTURES

ON

ANTISEPTICISM IN SECTIONS AND RESECTIONS.

By RICHARD BARWELL, F.R.C.S.,
Surgeon to Charing Cross Hospital.

LECTURE II.

GENTLEMEN,—In the last lecture we saw that division of the femur above the knee, combined sometimes with section of the tibia and fibula just below that joint, forms a valuable substitute for resection of the ankylosed portion itself. The advantages of the one operation over the other are the greater readiness with which the parts heal, the more complete preservation of muscular power, and probably, in young subjects, non-interference with epiphysal growths. In my experience (and I have performed osteotomy in more instances than those now reported), the reunion of bone is very rapid. In Case I, I believe that rare form, immediate union, took place, and that the girl might have gone about in ten days. Generally, the lower end of the femur is firmly knit within six weeks, the tibia within a month. The antiseptic method obviates—I had almost said precludes—suppuration, hence the operation is as safe as any human action can possibly be; it is also nearly painless: none of my patients have suffered pain or loss of health.

The modelling process which took place in C. G. (Case 1*) is very remarkable; no abnormal bend whatever of the femur is apparent, nor is that the only instance. Of John B. (Case II), I said in my last lecture that the appearance of angular incurvation at the femur and tibia was diminishing. Since those words were written, the modification has gone on even more rapidly. It would seem that, as in rachitis the inside of the morbid curve is filled up and strengthened by new bone, so in the retreating angle of these artificial fractures new osseous deposit takes place, while the salient angle is absorbed; thus the divided and bent bone becomes, without losing its direction, very much straightened, and that projection of the knee—a marked peculiarity of fig. 5

(see this JOURNAL, April 28th, p. 507)—gradually diminishes, and may very probably almost disappear.

The experience of Case II is valuable in this wise. Owing to the unsound, though not diseased, condition of the femur and soft parts close to the knee, I felt constrained to divide that bone higher up than in Case I. Now, to place the lower leg in a certain line, the lower end of the femoral fragment, whatever its length, must be flexed at a given inevitable angle with the upper one: therefore, it is evidently desirable to make the lower fragment as short as possible, since its projection will be in direct ratio with its length. The synovial membrane need not, of course, be taken into account; but the epiphysal line, at the age when it still exists, should not be interfered with, since most of the femoral growth depends on this structure.

We will now pass on to another subject—namely, to excisions of the knee, and to the effect which antisepticism exercises on the duration and prospect of cure—an effect which cannot, I think, be too highly prized.

CASE I.—Chas. B., aged 8, admitted into the Charing Cross Hospital under my care in August 1876, with disease of the knee-joint, evidently arising from osteitis in the femoral epiphysis. Under a treatment which I will not now detail, he made a marked improvement; but in October he became so suddenly worse that the relapse could only be ascribed to some imprudence (the boy is very full of freaks), which he could not be brought to confess. The symptoms became so aggravated, that very soon excision evidently offered the best chance of preserving the limb.

November 10th. I performed the operation, under all antiseptic precautions, in my usual way, making a free straight cut across the front of the joint almost from one hamstring tendon to the other. A good deal of pus escaped. When the bones were sawn through, a large spot of softened bone-tissue had to be cleared with the gouge, both from the inner condyle and from the inner tuberosity of the tibia. A branch of the external articular artery was tied, and three smaller vessels were subjected to torsion. A drainage-tube was placed at the back of the bones, and the limb placed in a splint of thin zinc and plaster of Paris, to be described immediately.

This splint and general arrangements required no change from the time of application until the 30th January, when the wound was healed, except the openings of two little sinuses, and the limb, therefore, was entirely placed in a plaster of Paris bandage, with the necessary windows. There was not during the whole of this time any swelling of soft parts—so frequent an accompaniment of this operation; neither was there any purulent discharge nor rise of temperature. The wound, save where the drainage-tube was kept for some time, and one other spot, healed by the first intention. It was dressed for the first month every other day, then for three weeks every third day, then every fourth day.

The splint—which after several trials I have come to regard as the most efficacious—is a modification of P. H. Watson's. It is constructed of two pieces of thin zinc, cut on paper patterns, carefully fitted to the back of the thigh and leg. The upper one, longer on its outer than on its inner side, extends from about two inches above the place of

Fi. 1.

wound to the great trochanter on the one and to near the fold of the nates on the other side; it must only be broad enough to go one-third round the thigh. The lower splint, commencing about one inch and a half below the wound, supports the calf, then forms a long and narrow part, which terminates in a foot-piece. In application, this

* I must apologise for a mistake in dates, which crept in, I hardly know how, from much copying. Catherine G. was admitted in November, and operated on December 2nd.