shallow; the pulse, however, was good and the inhaler was kept to the face. In a few seconds more, the movements of the chest and abdomen were scarcely perceptible, and the face, which had till now been of good colour, began to be blanched. The inhaler was at once removed; but the pulse now began to fail, and was soon scarcely perceptible at the wrist, though the heart was easily felt beating against the chest-wall. The pillow was pulled from under the head, and the chin thrown forcibly upwards by pressure behind the angles of the lower jaw. The muscles seemed to be fully relaxed, and the surface quite insensitive. The case looked exactly like one of syncope from chloroform, and I expected to have to resort to artificial respiration. On flipping the chest and face, however, with a cold wet towel, slight inspirations were taken, and very soon the head was moved, colour returned to the face, and the pulse was again felt at the wrist. On moving the diseased joint, the child opened her eyes and began to cry, and presently asked if we were putting her leg straight. The inhaler, with two drachms more of ether, was now reapplied, and again the patient began to inspire very freely. Only a few inspirations of ether were allowed, but they were quite enough to produce complete anaesthesia, and, although the pulse did not again flag, respiration soon became extremely feeble, as it had done before. The inhaler was but occasionally applied for short periods, and without being replenished with ether, whilst a dislocation of the hip was reduced, and weights and a long splint were applied to the limb. In about twenty minutes from the commencement of inhalation, when all was finished, the child awoke and talked a little nonsense, but was soon quite rational and comfortable. There was no sickness nor other observed ill effect of the anaesthetic. The ether was of specific gravity .735, and was from the same bottle as had been successfully used the day previous for anaesthetising a boy.

The above case is, I think, of considerable interest, as it points, perhaps more conclusively than the one recorded by Mr. A. S. Morton in the *Lancet* of October 14th, 1876, to failure of the heart's action, as well as of respiration, during the inhalation of ether, without obstruction of the respiratory passages. I have seen difficulty of breathing during ether inhalation, from collection of mucus in the larynx and in the nose; but in this case here recorded there was no evidence of any such condition. Nor do I think that a falling back of the tongue can have produced the symptoms; for it was distinctly in response to the wet towel, and not to the previous pushing upwards of the jaw, that respiration was restored, and, moreover, the pallor and the rapid failure of the pulse point to a different cause; there was at no time any congestion of the face. It is not, I know, uncommon for patients to breathe very lightly when recovering from ether anaesthesia after the completion of an operation; but in such cases they are, I think, always sensitive, and can at once be roused by the infliction of pain; this was not the case in the present instance. Contrary to what has been my experience in cases of chloroform syncope, the pulse gave no warning of the failure of respiration, though it became imperceptible very soon after the breathing had ceased. In conclusion, I may say that the impression of Mr. Denton and myself is that, if we had been using chloroform instead of ether, it would have been less easy to avert a fatal termination to the case.

A CASE OF OPISTHOTONOS CURED BY CHLORAL-HYDRATE.

By HARRY DAVIS, L.R.C.P.Lond., M.R.C.S.Eng., etc.,
Callington, Cornwall.

A SHORT time since, about midnight, I was requested to visit a man immediately, who was thought to be *in extremis*. On my arrival at the house, I found him greatly convulsed, and being supported in an upright posture by two persons, his head drawn back to its extreme point of flexion, and the spinal column arched forwards and very painful, especially near the base of the sacrum and towards the lower extremities. His face presented the usual *risus sardonius*, with retraction of the lower lip, etc. The eyes were fixed and motionless, with slightly contracted pupils. The countenance was very pale and distressed. Speech was nearly obliterated, words being very indistinct, with an ejection of saliva at each attempt to speak. He could neither sit, lie, nor stand; and, on the removal of his support, he would immediately fall backwards. The heart-sounds were weak; pulse regular and small, about 72. I at once ordered him a mixture containing eighty grains of chloral-hydrate, half a drachm of extractum opii liquidum, and camphor mixture to four ounces; a fourth part of the mixture to be taken every four hours until the spasms and pain were relieved. After the third dose was taken, he was much relieved and slept a short time. I then ordered him some powders containing calomel and

jalap; one to be taken every six hours until the bowels were freely acted upon. The back was occasionally rubbed with a liniment composed of turpentine and belladonna liniment. During the succeeding three or four days, he was frequently threatened with the spasmodic attacks; but, on taking a dose of the chloral mixture, he was immediately relieved.

When the spasms had entirely abated, hemiplegia of the left side and paralysis of the right side of the face set in. I then ordered him a mixture containing iodide of potassium with compound tincture of cinchona at a later stage; and occasionally at night, when there was much restlessness, a dose of the sedative mixture. His diet consisted of light farinaceous foods and beef-tea.

After a month's treatment, he became quite convalescent, and soon afterwards resumed his work, and continues to do so up to the present time, without the least sign of inconvenience or remains of the paralysis.

The history of the case, as far as I can gather, is as follows. About eight weeks previously to this attack, the man was employed in repairing a building, when the scaffolding gave way, causing him to fall several feet, when he seemed to have received a blow on the head, as he was picked up insensible. This condition passed off within a few minutes, so that he was able to resume his work without any inconvenience. From the time of the accident till within a few weeks of his illness, he occasionally complained of slight pain in the frontal and temporal regions of the right side of the head, with a certain amount of loss of hearing. A *thick purulent* discharge was then observed to be oozing from the right ear (but never any signs of hæmorrhage), with more pain, and at times very intense, which caused him to give up his work. He then applied to a medical man, who attended him until I was called to see him in his very grave condition. What the remote cause of the convulsions was, I think, is very doubtful. Could it be possible that a man receiving a fracture at the base of the skull, however slight or obscure, continued his work for six weeks with little or no inconvenience? or could it have been an ordinary case of localised inflammation terminating in suppuration? or had he been taking nux vomica or strychnia before I was called to see him?

THERAPEUTIC MEMORANDA.

PITYRIASIS VERSICOLOR TREATED BY MEANS OF GOA POWDER.

At a time when the action of Goa powder is so much under discussion, it may not, perhaps, be out of place to give my experience of its application in a disease for which I believe it has not been previously used. If I relate one "out of several cases", it will illustrate my point sufficiently. A gentleman applied to me for advice with his chest and abdomen covered with pityriasis versicolor, which had existed for several years, and was gradually creeping above his collar; I gave him the following prescription (which I find to be a very convenient form).—R Goa powder gr. 30; acetic acid ℥i 10; glycerine ʒj—and told him to apply it over the affected parts at bedtime for four nights. He saw me again in a fortnight, when the disease had entirely disappeared, and the only marks visible were some livid patches, due to the application, which were gradually disappearing. He then told me that the morning after the first use of the remedy, the parts anointed presented a very red appearance, as if a mustard poultice had been applied, but that the irritation did not increase much, although he persevered with the treatment for the time recommended.

I find the above formula a convenient and efficient application for ringworm, and hope soon to be able to give the history of some obstinate cases of pityriasis rubra which are now being treated by the same preparation.

A. W. MAYO ROBSON, M.R.C.S., L.R.C.P.,
Demonstrator of Anatomy, Leeds School of Medicine.

CHRYSOPHANIC ACID.

I ORDERED chrysophanic acid ointment for an obstinate case of ringworm of the scalp, which extended behind the left ear. After a few applications, the skin of that side of the neck and ear became swollen and inflamed, of a dark or dirty red colour, its surface covered with minute vesicles. The skin around both eyes (about half an inch wide) was stained claret-colour; and conjunctivæ so intensely inflamed that the patient could not open her eyes for days. When the ointment was discontinued, the inflammation gradually subsided. I tried it also in another case of ringworm, when the skin around became much inflamed and of the same dark red colour.

THOMAS E. JONES, Llanrwst.