ous hæmorrhages were to be seen in the retinal segment supplied by the blocked arterial branch. Now, as Knapp says, this might be called a case of hæmorrhagic infarction of the retina. The interest of such a case is that it helps us to complete our knowledge of the effects of embolism in general. It certainly is not widely known that the effects of plugging of arteries is commonly, not anæmia, as one might at first glance suppose, but hyperæmia and extravasation. The case might be called red softening of the retina; the process is just the same in cerebral red softening; there is arterial plugging leading to extravasation of blood and softening of the brain. It is so in the spleen and kidney, etc.; also in what are often called pulmonary apoplexies. In all these cases the blocking leads very quickly to hyperæmia and extravasation. That blocking of arteries can lead to total hyperæmia, to increase of nutrient fluid is, I think, worth some consideration in regard to the pathology of those diseases, such as epilepsy and chorea, in which the increased expenditure of energy implies of necessity increased nutrition by some process. Late in these cases the blood is absorbed and the disc atrophies. A similar change, making allowance for difference in the organs affected, follows in red softening of the brain; there is yellow softening and red splenic and renal blocks becoming washleather-like. Sometimes there is hemiplegia from embolism, and at the same time amaurosis from plugging of the arteria centralis retinæ. I saw a case of this kind in Sir William Jenner's practice with Dr. Gowers. In this case, Sir William's diagnosis was confirmed by the necropsy: a valuable account of the case has been published by Dr. Gowers. This is the only case of recent retinal embolism with hemiplegia that I have seen. I have seen several cases of hemiplegia in patients who had atrophy of one optic disc. It is very important in such cases to note exactly the condition of the disc and its arteries, or we might be misled into the opinion that the atrophy was a consequence of the cerebral disease which caused the hemiplegia. This would be a very serious error; for, so far as I know, no kind of cerebral disease produces atrophy of but one optic nerve except by the process of neuritis, and that is, in physicians' practice, a thing of extreme rarity. As mentioned, Charcot believes that disease of a certain part of one cerebral hemisphere causes blindness of the opposite eye, but he says there are no ophthalmoscopic appearances.

[To be concluded.]

A CONTRIBUTION TO THE NATURAL HISTORY OF CATARRHAL OPHTHALMIA.

By EDWARD NETTLESHIP, F.R.C.S.,
Surgeon to the South London Ophthalmic Hospital.

THE every-day muco-purulent ophthalmia which most surgeons specially designate as "catarrhal" presents some features of interest peculiar to itself, and perhaps of importance in reference to local inflammations of other parts. The view has for some years past been taught, though not, I believe, put upon record, by Mr. Hutchinson, that this disease, in many of its features, resembles the exanthems—a theory based, I believe, on the observations that the disease is almost always symmetrical, that it attacks both eyes almost simultaneously, runs a definite course usually ending in spontaneous recovery, and that it scarcely ever occurs twice in the same person. Some of the facts given below will be found to favour this view, while others appear to point to a different conclusion; but careful records of a very large number of cases can alone decide the important points raised, and I shall for the most part avoid drawing any conclusions from the very limited data given in the present communication.

It may be well to call attention here to the rather vague meaning attached to the term "catarrhal" in connection with the eye. We may apply the term to the slight conjunctivitis which in some persons accompanies nasal catarrh; or to a mild form of disease, sometimes allied to episcleritis, in which the exposed parts of the eyeball become congested and slightly inflamed from exposure to cold or damp, and very liable to relapse; or to the chronic, chiefly palpebral, conjunctivitis sometimes associated with uncorrected ametropia; or to the chronic conjunctivitis so common in old people; or to the muco-purulent ophthalmia which often follows measles, scarlet and other fevers. Yet these forms of disease are very different from one another and from that for which, by common consent, the term is most emphatically used.

The facts given below are collected only from cases of the disease referred to in connection with Mr. Hutchinson's hypothesis, the diagnosis being based in each case on the attack being acute and accompanied by considerable muco-purulent discharge, and, in addition, presenting one or more of the following features:—(a) symmetry, both

eyes being attacked within a short time; (b) a history of contagion; (c) conjunctival ecchymoses.

If the disease be a constitutional one, of which the conjunctivitis is to be looked upon as the eruptive stage, a certain degree and duration of fever must be expected. I have had no opportunity of making continuous daily observations of the temperature, and the few records given below in this connection show no more than that there is a slight rise of temperature in a large proportion of the cases, and that this is greater in severe cases and at an early stage, than when the disease is mild or has lasted a relatively long time. It is, however, very remarkable that any exanthem should be characterised by an eruption limited to so small a part of the surface as that furnished by the conjunctiva, and that it should seldom, if ever, be accompanied by evident symptoms of ill-health. On the other hand, it is equally remarkable that the disease, if it be merely local, should be self-protective; for I do not know that this is true of any contagious local disease, unless favus be considered a case in point, where the scalp may become so scarred, that the hair-follicles are destroyed and no soil remains for the cryptogam to flourish in. It may be urged, however, that the absence or trifling severity of the constitutional symptoms can be scarcely less marked than in some cases of such undoubted specific fevers as chicken-pox and syphilis; whilst the limitation of the eruption to the conjunctiva might be accounted for by appealing to some anatomical and physiological peculiarities in this membrane. That the conjunctiva does possess some properties not found in other mucous membranes is rendered probable by the remarkable chronic changes which it undergoes after prolonged exposure to impure and damp air, changes giving rise to the chronic granular conjunctivitis or trachoma. And the occurrence on the ocular conjunctiva of phlyctenulæ or so-called "pustules" appears to point in the same direction; for they do not seem to be represented by any changes commonly met with on other visible mucous membranes or on the skin. Should further observation prove that a first attack of this "catarrhal" ophthalmia confers immunity from a second, and should a persistent elevation of temperature be found which could not be accounted for by the local inflammation, the exanthem hypothesis would acquire great probability.

In making a short analysis of about two hundred and seventy cases of this "catarrhal" or ecchymotic ophthalmia which have during the last few months* been under my care, the following points will be dwelt upon.

1. *The Temperature.*—This was taken in 91 cases; in 85 of them only once; in the other 6, twice at an interval of several days. The cases were not selected in any way. The observations were all made between about 2.30 and 4 P.M. on the day of the patients' first visit; a six-inch Hawkesley's thermometer was used, and the temperature taken in the axilla. The value of the observations is certainly lessened by their having been made nearly at the time of the daily highest physiological maximum, by many of them having been taken in summer, and by the patients frequently being women or children. Thus, of the 91 patients, 28 were women or girls of 14 and upwards, and 38 children under 14; only 28 were men or boys over 14. Making all possible deductions, however, the figures show that a definite elevation of temperature was present in a considerable proportion of the cases. The increase was never great; in only 13 cases was the temperature over 100 deg. Fahr. In one, a female child aged a year and a half, it reached 103.1 deg. Fahr., but in none of the others did it exceed 101.7 deg. Fahr. The following table gives some further details.

	Total, 86	Males, aged 14 or more.	Females, aged 14 or more.	Children under 14.
Temp. above 100.4 deg. Fahr. (slightly febrile)	5	1 (20 per ct.)		4 (80 per ct.)
Temp. between 99.5 deg. and 100.3 deg. (subfebrile)	36	10 (25 per ct.)	16 (50 per ct.)	10 (25 per ct.)
Temp. between 99.4 deg. and 99.1 deg.	26	8 (30 per ct.)	6 (24 per ct.)	12 (46 per ct.)
Temp. normal (99 deg. or lower)	19	5 (26 per ct.)	4 (21 per ct.)	10 (52 per ct.)

Of the 41 febrile and subfebrile temperatures, 11 (or 25 per cent.) were in men, 16 (or 40 per cent.) in women, and 14 (or 35 per cent.) in children; while, of the 45 remaining which were normal or very nearly so, 13 (or 29 per cent.) were men, 10 (or 22 per cent.) women, and 22 (49 per cent.) children. Thus the women furnished a much larger proportion of high temperatures and a somewhat smaller proportion of normal temperatures than the men. The children gave a proportion of high temperatures intermediate between the men and women, while they much exceeded these in the proportion of normal tempera-

* This paper was written in February.

tures. The latter fact is, no doubt, due to the difficulty in taking the temperatures of young children in an out-patient room, and does not represent the truth.—In six patients, the temperature was taken a second time, several days having elapsed between the two observations; in four of them, the second temperature was normal, the first having been above normal, and in all these both eyes were already affected when the first observation was made. In the other two cases, the second eye was attacked between the date of the first and second thermometric observation, and in each of these the second temperature was the higher.—The majority of the high temperatures were in the more severe cases. In only one of those with febrile or subfebrile temperature was the case mild; viz., a woman aged 42; in nearly all the others, the disease is marked as "severe" or "moderate." On the other hand, in five-sixths of the cases marked "mild", the temperature was practically normal.

Only three patients came under care within twenty-four hours of the onset of the disease in the first eye, and the temperature was above normal in all of them. It was the same with four others who came with both eyes affected, but within twenty-four hours of the onset in the second eye. Again, the cases with high temperatures were, on the average, in a somewhat earlier stage of the disease (counting from the onset in the first eye) than those in which there was no elevation. Further, while none of the cases with the highest temperature were in a later stage than the fourth day (counting from the commencement in the eye last affected), in those with normal temperatures the disease was often at the fifth, sixth, or seventh day, or even later. Lastly, in none of those where the temperature was taken on or after the sixth day was there any material elevation. These are the only facts I have to offer which bear on the relation between the temperature and the stage of the disease.

2. *Evidence of Protection afforded by a First Attack.*—Some kinds of ophthalmia, e.g., phlyctenular cases, are especially liable to relapse; in others, e.g., granular ophthalmia, the patient never really recovers, and, though remaining sometimes for long periods free from active symptoms, he is in constant risk of recurrences. On the other hand, whilst the group of purulent cases furnishes but few instances of second attacks, this is probably due in no degree to the exhaustion of liability on the part of the conjunctiva to purulent inflammation, but simply to the rarity of a second exposure to inoculation with the necessary material.

In the ophthalmia now under consideration, the rarity of second attacks, if it be a fact, can scarcely be referred to the rarity of opportunities for a second inoculation; the disease is so common that numbers of those who have had it once will probably be exposed to the risk of it again. Some other explanation must, therefore, be sought.

Of the two hundred and seventy-one cases of which I have notes, and in nearly all of which the question as to a former attack was asked, twenty-seven gave the history of an attack of some kind of ophthalmia at a previous date. In most of these, no opinion could be formed as to its character; in some, it had certainly not been "catarrhal" ophthalmia; but, in about half a dozen, the account given was such as to leave little doubt that a former attack of the same disease had occurred, and in all of these the attack which I saw, presumably the second, was quite characteristic. The following are short notes of the cases referred to.

CASE 24.—M., 5. Characteristic ecchymotic ophthalmia. Sister and another boy have it. A year and a half ago, the patient, and the same boy who now has it, had bad eyes for a month.

CASE 39.—F., 6. Characteristic ecchymotic case. No contagion. Two years ago, had "blight" in both eyes, and at that time all the other children had it too.

CASE 60.—F., 36. Characteristic case, with history of contagion. Six years ago, had "blight" in both eyes for nearly a month.

CASE 184.—M., 9. Characteristic ecchymotic case. Contagion from a brother. Two years ago, had a similar and equally bad attack, when mother and baby also had it.

CASE 253.—M., 28. Characteristic ecchymotic case. All his three children have it. Three years ago, he had precisely the same thing in both eyes; eldest child (at that time the only one) also had it, and has it again now. Both were soon well in former attack.

The three next cases (Nos. 66, 144, and 225) illustrate not only a second attack (in Nos. 66 and 144), but the escape during the present attack of several who had the disease at an earlier date.

CASE 66.—M., 44. Characteristic case. Three of his eight children

have just had it. A year and a half ago, he had just the same thing; wife and one child had it then, but neither of them has had it this time.

CASE 144.—F., 10. Characteristic ecchymotic case. One child in the same family and several in another family in the same house have just had it. Two years ago, both eyes were bad in just the same way, and were well in a fortnight. Mother and four other children of the family had it then and have escaped now.

CASE 225.—M., 41. Characteristic ecchymotic case. Four of his eight children have just had it. None have had it before. Some years ago, his wife and eldest child had it, and have not now caught it again.

3. *Symmetry.*—In one hundred and fifty-four cases, it is noted that, on admission, the disease was symmetrical, and in one other that it became so after admission. In about seventy-five, there is no note on this point; but it is certain that, in a large number of these, the omission was due to the fact that the cases conformed to the rule, i.e., that they were symmetrical. In forty cases, it is noted that only one eye was affected on admission; but, in three-fourths of these, the disease had begun so recently that there was no probability of the second eye escaping. A small number (nine cases) remain in which from five to eight days had, on admission, already elapsed without the implication of the second eye; but, even in most of these, it is by no means certain (as will be seen from considerations given below) that the disease did not ultimately become symmetrical.

The interval between the beginning of the disease in the two eyes is usually very short. Of eighty-eight cases in which it is recorded, the interval was one day or less in thirty; from one to three or four days in forty; four days in six; five days in five; one week in four; more than one week in three.

It is important to note that, in the cases (twelve in number, and all of them very characteristic) where the interval was longest (five days or more), nearly all the patients were adults, two only being children aged 4 and 10 respectively. It is easy to account for the delay in the affection of the second eye in adults as compared with children, on the ordinary supposition that it is inoculated accidentally by the fingers, etc., from the first eye; this being less likely to happen quickly in proportion as the patient is older and more careful. On the exanthem hypothesis, it would, however, be difficult to explain. Returning to the cases (nine in number) in which, on admission, a long interval (five to eight days) had already passed without the second eye suffering, I find that none of them were children, and that only one was under 20, and it thus seems highly probable, as already mentioned, that in most of them the second eye did suffer afterwards. The disease began in the right eye in ninety-six cases, in the left in seventy-seven, and in both together in six.

4. *Contagiousness.*—The commonly received opinion as to the communicable nature of this ophthalmia is fully borne out by statistical inquiry. In more than one-half of the cases (151), there was a history that the patient had been exposed to the risk of contagion to or from another person suffering from a similar complaint. When it is borne in mind that the patients are usually seen only once, and that inquiries of this kind are often hurriedly made and sometimes omitted, it will be granted that, in all probability, a similar account would have been given by a much larger proportion had time allowed. The total number accounted for by each patient who gave a contagion history was such that, on the average, each of the 150 cases under care represented two others which were not seen; or of 450 cases, 150 came under notice. The disease appears to be far more communicable than purulent ophthalmia derived from gonorrhoea or leucorrhoea; for, although its duration is much shorter, contagion is far more common.

The patients are almost always either children, or married people with families. It is not common to find cases in adults who have had no communication with children. Thus, of the 150 cases with a contagion history, the first patient is stated to be a child in 97 (nine of these being babies in arms), the mother in 6, the father in only 3. The children are more likely to catch the disease than either of the parents; for, of 136 families in which contagion had occurred, the mother had caught the disease in only 37 and the father in 32.—These facts do not favour the belief that exposure to cold, wet, dust, or impure air are causes of this form of ophthalmia.

The disease appears generally to be communicated whilst in an active state, and it is common to find it running through a family of half a dozen persons in as many weeks. But, in a few cases, it is probable that the communication does not take place until quite a late stage, although, in making this inference, I do not forget that there are fallacies which, in hospital clinical work, it is impossible to avoid. It may also be that the malady has an incubation of several days, so that, although the inoculation may occur in an active stage, no symptoms

* It may be noted that the terms "severe", "mild", etc., cannot always be applied with strictness, since the various symptoms of the disease (congestion, ecchymosis, swelling of lids, and discharge) are not always present in corresponding degrees.

show in the recipient until the communicator is nearly well. The following case is perhaps an instance of this.

A lad living in London had "blight" in his eyes; while still suffering somewhat, although said to be nearly well, he went to Gravesend to see his sister, a servant girl aged 16. She gave up her bedroom to him for the night, and he returned to London in a day or two. Eight or nine days after his visit, his sister's left eye inflamed, and the right followed on the day after; five or six days later, she came to me with well marked ecchymotic ophthalmia.

Other instances in which a long interval seems to have elapsed between successive cases are the following.

CASE 136.—M., 17. Characteristic ecchymotic ophthalmia; four days. Three weeks ago, his "mate" had it and is now well.

CASE 71.—F., 18. Characteristic ecchymotic case, with a week's interval between the two eyes (*sic*). Three months ago, her sister had a severe attack in both, lasting about three weeks. No other known contagion.

CASE 107.—M., 35. Characteristic severe ecchymotic case. During the last eight or ten months, all his six children have had it; wife has escaped.

CASE 178.—M., 35. Characteristic case; four days. Six weeks ago, his wife had it, and, within a fortnight of her case, three of his six children suffered. Then a month elapsed without a fresh case, when his own attack began quite suddenly.

CASE 34.—M., 25. Acute ophthalmia of one eye only; two days. Attended only once. Two of his children were suffering from "blight". Between four and five weeks after the onset of disease in the patient, his wife (Case 19) began to suffer, and came under care, with a sufficiently characteristic attack, after the disease had lasted six days.

5. *Natural Course of the Disease.*—In support of the assertion that the cases usually recover spontaneously in a short time, the following may be noted. 1. It is very common to hear of several cases having occurred in the family, and recovered without any treatment, excepting perhaps a little bathing with cold water, before the case under care is brought to the hospital. 2. A large majority of the cases attend the hospital for less than a week; which may be taken as positive proof that they nearly all get well a few days after admission. Thus, of the total number (273), 164 attended only once; 83 twice or three times, their whole attendance extending over from seven to ten days, or occasionally fourteen days; only the remaining 22 were under care three weeks or more. 3. The prolonged cases generally occur either in young children (under four or five), in whom the disease seems slightly prone to run into a chronic condition, or are accounted for by the occurrence of phlyctenulæ as a complication. 4. Severe cases (excepting in some young children) do not, as a rule, last much longer than mild cases.

The disease occurs at all ages. Several of the patients were fifty or more; one was an infant of three weeks, and several were infants of two or three months. About one hundred and forty of the cases were children under fourteen, the remainder being divided almost equally between males and females of fourteen and upwards.

ROUSTCHOUK MILITARY HOSPITAL.

By H. CROOKSHANK, M.R.C.S., etc.,

Surgeon-Major Turkish Military Service; Operating Surgeon to the Hospital.

THIS small but strongly fortified town, which is likely to play an important part during the present Turco-Russian campaign, now boasts of six large temporary hospitals, with an aggregate number of upwards of eight hundred and fifty beds. Four of these hospitals belong to the military authorities, one to the naval, and one to the civil, the two latter having been in existence some years; the naval hospital containing twelve beds; the civil twenty-five, which number, on an emergency, can be increased to one hundred or one hundred and fifty. Two more large buildings have been placed at the disposal of the medical authorities, one of them belonging to the Jewish community, and formerly used as a school for the children of that denomination. These latter will be fitted up with beds and reserved for surgical cases. Two months ago, a portion of the caserne was sufficient to meet the need of the medical service; but, as the number of battalions stationed here increased and each one brought with it a contingent of sick and feeble, it was necessary to obtain more space, and to the caserne was added the École Israélite, then a large building not quite finished, and intended for a school for Ottoman girls; and finally the Haremlik of the Kouak; and over all these buildings gaily floats a large white flag with a red crescent in the centre, which the Turks have chosen as an emblem of

protection and to warn the Russian gunners from pointing their pieces on an assemblage of suffering humanity.

During the last three days, the weather has suddenly changed from beautiful summer to cold winds and rain, which has told very much on the army under canvas and greatly increased the number of admissions into hospital. For two days we have had a continuous down-pour of rain, accompanied by thunder and lightning, which rapidly converted the narrow streets of this town into rivers from eight to twelve inches deep, and has caused the Danube to rise considerably above its usual height even at flood-time. One good effect this deluge has had; namely, to cleanse the roads and to carry into the river all the refuse matter which the inhabitants, in defiance of all sanitary measures, insist upon depositing outside their street doors.

The hospital administration is very good, due in great measure to the admixture of foreign element in the staff, which consists of thirteen doctors, namely, one Armenian, one Greek, five Turks, four German, and two English. The patients are well looked after, there being a good supply of hospital attendants; and the diet is liberal and of very fair quality. The buildings, having been intended for other purposes, are well lighted, properly ventilated, and answer admirably for the purposes to which they are at present applied. They are mostly detached buildings, at some distance from each other; and several of them are surrounded by pleasant well-kept gardens, where the convalescent patients spend most of their time. During the month ending May 12th (April 30th, local time), a total of 1,972 cases were treated at the various hospitals. The largest number being in hospital on one day was 835. On April 13th (April 1st, local time), there were 463 patients in hospital; 1,509 were admitted during the month, making a total of 1,972 cases, of which number 1,106 were dismissed cured or relieved, and 104 died: a mortality of about 5.2 per cent., which is not excessive, considering the fact that most of these men have been eighteen months under arms and passed through a very severe winter in Servia, many of them not having yet recovered from the wounds they received in that campaign. The number of soldiers in this neighbourhood under arms has been estimated at about 25,000, and the average daily admittance into hospital for the month was 65.7, most of them serious cases. The maximum number of cases in hospital at one time was 835, an average of little over 3 per cent., which speaks well for the sanitary condition of this division, the usual percentage being about 1.5.

The following is a list of a few of the most important diseases admitted into hospital, with the result.

DISEASE.	In Hospital.	Dismissed, cured or relieved.	Died.
Anæmia and general debility	30	19	1
Abscesses	49	37	1
Inflammation of glands (various)	36	26	0
Diseases of chest	396	205	20
Phthisis	57	16	25
Diseases of eye	35	18	0
" skin	67	42	0
" ear	19	18	0
Rheumatism (acute and chronic)	114	69	0
Erysipelas	17	11	0
Simulant and hysteria	54	52	0
Gonorrhœa	57	36	0
Veneral disease	59	35	1
Catarrh of intestine	159	113	2
" stomach	80	51	0
Dysentery	63	18	11
Fever (various, but chiefly typhoid)	169	88	18
Influenza cold	113	92	0
Contusions	17	10	0
Ulcers (various)	32	9	0
Scurvy	218	84	10
Various	131	57	15

During the last three days, on account of the great change in the weather, the number of patients admitted into hospital has increased, no less than 354 serious cases having been taken in during the week; so that the numbers stand thus:

In Hospital May 1 (local time).	Admitted during week.	Total.	Dismissed.	Died.	Remaining in Hospital end of week.
762	354	1116	325	22	769

The daily average of beds occupied during the week was 843, and the average daily mortality 3.14. The following were the causes of death: Dysentery, 8; typhoid fever, 5; scurvy, 2; heart-disease, 1; Bright's disease, 1; burns received, 1; pneumonia, 4; pleurisy with effusion, 1; phthisis, 4.