drachms of greenish pus. The interior of the cavity consisted entirely of white brain-substance, and was of a firmer consistence than the rest of the brain. There were also in the cavity two small blood-clots, partially organised. The right lobe of the cerebellum was, comparatively speaking, a mere shell; and the left lobe, as well as the pons Varolii, was to a small extent also implicated. The membranes were perfectly healthy, and there was no apparent communication between the abscess and the suppurating ear.

NOTES OF A CASE OF CARBONIC ACID POISONING, TREATED BY INHALATION OF OXYGEN :

WITH A DESCRIPTION OF A NEW APPARATUS FOR RENDERING
IMPURE AIR RESPIRABLE.*

BY CHARLES B. BALL, M.D.,
Surgeon to the Blaenavon Iron Works.

ON January 16th, 1877, I attended a family who were found insensible in bed that morning. I found a man aged 55, his wife aged 48, and daughter aged 16, in one bed, in a very small room, quite insensible. There were the remains of a large fire in the grate, the chimney of which was imperfect. Upon admitting plenty of air and using the usual restoratives, the woman became sufficiently conscious to say that she remembered being up at 2 A.M., when she experienced considerable difficulty in breathing and a severe headache; after which she remembered nothing further till she was restored to consciousness. The man became semiconscious, but the girl remained perfectly insensible. I had her, therefore, removed to another and well-ventilated room, where artificial respiration was resorted to. She now had violent tetanoid convulsions, at intervals of a few minutes. The muscles of the face were, however, exempt from the general spasm. The pupils of the eyes were of average diameter, but dilated in bright light and contracted when the light was removed. This phenomenon was also noticed in the man's case.

During the day, the man and woman progressed favourably, but the girl became much worse, the convulsions appearing at longer intervals and the intervening coma being more profound. The treatment now adopted in her case was the application of a blister to the back of the neck, stimulating enemata, and the hypodermic injection of pure ether. None of these measures had the desired effect, and at ten o'clock that night she was apparently dying. The convulsions had entirely subsided. The pulse at the wrist could not be felt, and the respirations were few and shallow. Through the kindness of Mr. P. C. Gilchrist, F.C.S., I now obtained a large supply of pure oxygen, which, by a little arrangement of an ether-inhaler, I was enabled to give her to breathe in any degree of dilution required. At first equal parts of air and oxygen were given, but afterwards the proportion of oxygen was increased. It produced a rapid and most marked effect; the pulse soon became perceptible, and quickly increased in force. After the inhalation of about four gallons of gas, it was discontinued, respiration and circulation having been well revived. During the inhalation, the convulsions recurred, having been absent for several hours. Although not sensible during the night, she was decidedly better, being able to swallow fluid nourishment, and the convulsions gradually subsiding. She did not recover consciousness till the morning of the 18th, being forty-eight hours from the time I first saw her.

The different degrees in which the persons suffered, although exposed to the same poison, appears to be worthy of note. The intensity of the girl's symptoms is, I think, accounted for by the fact that she suffers from phthisis, and has therefore a smaller available lung-surface. As the woman got up to walk about the room during the night, she for the time respired a higher, and therefore purer, atmosphere; and this may account for the fact that her symptoms were the lightest of the three. The action of the pupils with regard to light, noticed in two of the cases, was remarkable and very apparent. Although the inhalation of oxygen has been recognised in medicine since the time of Sir H. Davy, yet its use amongst the generality of practitioners is not frequent, and the above case was the first time I have ever used it. Its action impressed me greatly; for I have never witnessed a life so certainly saved by any other therapeutic agent. One great objection to its more general use has been the difficulty of obtaining the gas readily, in a state of purity and in a portable form. This has now been obviated; and I have here a copper flask, into which fifteen gallons of pure oxygen have been compressed. I obtained it from Messrs. Barth

of 26, Duke Street, Bloomsbury, and it appears to be very convenient for medical purposes, being easily carried about, as its weight is only five pounds three ounces when full; the diminution in weight will tell roughly the amount of gas used.

From my experience in this case, I was led to the construction of an apparatus for obviating the bad effects produced by inhaling an atmosphere containing a large percentage of carbonic acid, by the simultaneous inhalation of an increased supply of oxygen; the applicability of some such apparatus being demonstrated by the experiments of Regnault and Reiset, who found that animals could live in an atmosphere containing a large quantity of carbonic acid, provided the percentage of oxygen was proportionally increased. The action of this apparatus is partially illustrated by the annexed diagrams. (Fig. 1.) It consists essen-

Fig. 1.

tially of a tightly fitting face-piece, which communicates with a chamber of a capacity sufficient for one inspiration. The chamber is separated from the outer atmosphere by a sheet of wire gauze, and the air is made to traverse several partitions. Behind the partition next the face-piece, oxygen is allowed to escape from a perforated tube, which is in communication with a reservoir, carried in the manner of a knapsack. The tube is supplied with a tap, which will allow the escape of

Fig. 2.

oxygen up to 30 per cent. of the air respired. During the intervals of respiration, the diaphragms will prevent the escape of oxygen from the chamber; and at each inspiration, the amount of oxygen required will be washed through the valve at the face-piece by the external atmosphere. The expired air is made by a system of valves to pass through an elastic desiccator, and thence supply a lamp. (Fig. 2.) By this means, the lamp can be made a guide as to the amount of oxygen required. By the use of this apparatus, I expect that a collier would be enabled to go into a mine charged with carbonic acid. In the case of an explosion, in which all the oxygen had been converted into carbonic acid and steam, the explorer would be enabled to supply himself with such an amount of oxygen as would neutralise the effects of the carbonic acid. The man could easily carry a supply of oxygen which would last for two hours, even when he was using the maximum quantity required; and, by connecting a second tube and face-piece to the oxygen-reservoir, he could revive any sufferer whom he might find overcome by carbonic acid. I may mention that, in the experiments connected with the construction of this apparatus, I have myself respired, without feeling any discomfort, during a considerable time, a mixture containing 18 per cent. of carbonic acid, but in which the

* Read at the South Wales and Monmouthshire Branch.

oxygen was increased to 30 per cent. This amount of carbonic acid is 6 per cent. more than that found after colliery explosions in which complete combustion has taken place.

THE TREATMENT OF ASCITES BY ABDOMINAL COMPRESSION.

By STEPHEN MACKENZIE, M.D.,

Assistant-Physician to the London Hospital.

THERE are numerous ways of treating the ascites dependent upon contraction of the liver. Alteratives, drastics, diuretics, tonics, compelling the patient to satiate his thirst from his peritoneal cavity, simply keeping the patient at rest in bed, all have their advocates, and are plans under which I, as every other physician, have seen good results in even unpromising cases. Every now and then, however, under any of these plans of treatment, alone or in combination, the peritoneal effusion remains stationary, and its removal becomes tedious. I have found that, in cases where there is any tendency to absorption, this may be greatly accelerated by a simple method—that of abdominal pressure—to be described, and that in cases in which absorption does not appear to be taking place the same plan is often successful in removing the fluid. I have employed this treatment in several cases, with such good results, that it encourages me to hope that others will find it useful. Perhaps I can best illustrate the plan of treatment I have found so successful in some cases of ascites, by narrating the first case in which I put it into operation.

E. S., aged 41, widow, was admitted into the London Hospital under my care on January 16th, 1875. The patient stated that she had not been thoroughly well since her last confinement, nineteen years previously. About one year before admission, in January, she noticed that her waistbands became tight. She attached no importance to the occurrence, however, at first. In February and March, she found that she was passing extremely small quantities of urine. In May, the size of the abdomen was much increased, and its "weight" inconvenienced her when she moved about. The enlargement of the abdomen increased rapidly up to June, when she attended at another hospital as an out-patient. She then measured forty-two inches round the largest part of the abdomen. She attended for three months, and during that time increased in size, and continued to increase in size to the end of September, at about, she thinks, the rate of "an inch a month". Since September, the increase had been less marked, but had slowly continued, until coming under treatment. She had only passed very small quantities of urine. She was the mother of two children, and had miscarried once, owing, she thought, to a strain. There was nothing in her history to support the diagnosis of syphilis; no intemperance could be elicited on close questioning. She had never had jaundice. When admitted, I noted that the patient was much emaciated, but was not decidedly cachectic. The abdomen was greatly distended, of ovoid shape, tense, with extremely distinct fluctuation-wave. The circumference of the abdomen at the umbilicus was forty-six inches, and the same two inches higher. Medium-sized superficial veins were seen coursing over the abdominal walls from below upwards, as could be demonstrated by pressure. There were no veins passing downwards to the thighs; the umbilicus protruded slightly. There was oedema of the lower extremities, from the thighs to the feet, especially on the left side. There was a small patch of discoloration on the inner aspect of the left leg, which, she stated, had been in existence for five years. It came as a small speck of the same colour, and has gradually increased to its present size. There was no history of injury to the part; no nodes. Hepatic dulness in the right mammary line was bounded by the upper border of the fourth rib. Here and in the axilla it was about two inches in extent, and posteriorly on a level with the twelfth dorsal spine, it was four inches deep. The chest organs were pushed up, but presented no noteworthy signs. Vaginal examination showed the uterus and anterior wall of the vagina depressed (cystocele).

On January 22nd, the patient's expression was anxious. She complained of difficulty of breathing, and her respiration was seen to be embarrassed. There was a slight herpetic eruption around the mouth; the patient had not been observed to have a rigor since her admission, but was suffering from catarrh when she came. The bowels had acted daily; the motions were solid and of good colour. She passed about fifty-five ounces of urine *per diem*. It was of an orange colour, free from albumen and bile-pigment. She was ordered to take five-grain doses of iodide of potassium thrice daily.

January 26th. The patient had a less anxious expression, but still looked haggard and as if she were the subject of abdominal disease. The average quantity of urine was about seventy ounces *per diem*.

Her breathing was easier, but entirely thoracic. The measurement of the abdomen at the same point as before was forty-five and a quarter inches. The legs were less swollen than on admission.

February 2nd. Whilst the patient expressed herself as better, and appeared cheerful, she looked more haggard and pinched in the face, and had dark areolæ around the eyes. The sister stated that she began to sleep with her eyes open. The abdomen had considerably diminished in size, was quite lax, and fell over to the side on which she lay. It was forty-two and five-eighths inches in circumference at the level of the umbilicus. The superficial veins were not so distended. She took all the nourishment supplied to her.

February 12th. The patient appeared better. The iodine appeared to be producing toxic effects. She had an acnoid eruption on the face, and had watering and smarting of the conjunctivæ, running at the nose, and soreness of the mouth. She was ordered to discontinue the iodide of potassium, and to take a mixture containing citrate of iron and quinine.

February 23rd. The patient looked and expressed herself as better. She slept and ate well; she passed from fifty to sixty ounces of urine daily, which had a specific gravity of 1,016 to 1,020, and was free from albumen. The largest measurement of the abdomen was now thirty-nine and a half inches. The oedema had wholly disappeared from the legs.

March 30th. The patient's general condition was improved. The abdomen now measured thirty-six inches; its summit was about three-quarters of an inch above the level of the sternum, and it was flat on the surface. The edge of the liver could now be felt three inches below the point of the xiphoid cartilage. No irregularities could be felt on the surface of the liver.

May 11th. The condition of the patient appeared stationary. There was still a considerable amount of fluid in the peritoneal cavity. She passed about forty ounces of urine in the course of the twenty-four hours. *The abdomen was ordered to be tightly bound with a flannel bandage, so as to exercise pressure.*

June 4th. When the bandage was first applied to the abdomen the pressure caused a feeling of sickness, but soon the patient bore it without discomfort; in fact, it appeared to afford her relief. The measurement around the abdomen, at the level of the umbilicus, was now thirty-three inches. She passed about thirty to forty ounces of urine daily, and did not perspire much. I now ordered a properly made abdominal support to be constructed for her by the instrument-maker. It was arranged with straps, so that it could be tightened to a very considerable degree.

June 29th. The patient expressed herself, and appeared to be, in good general health, but she was still much wasted. The abdomen looked much smaller, and measured thirty-one and a half inches at the level of the umbilicus. There was pseudo-fluctuation on percussion, but it did not convey the impression of being due to fluid. The liver could be felt in the same situation as before. Its surface was smooth, its edge firm, sharp, and regular. The spleen was not to be felt in the abdomen. Its dulness reached the seventh rib in the axillary line. She passed, on an average, fifty to sixty ounces of normal urine daily. She ate and slept well. She was now allowed to get up, but was directed to continue wearing the abdominal support. The latter, owing to the diminution of the size of the abdomen, had required to be altered. After being up in the ward for a few days, the patient was allowed to go into the garden; and there then being no reaccumulation of fluid in the abdomen, she was made an out-patient. This was in July 1875. The patient has been continually under my observation from then until now. She has had no recurrence of the ascites. Her general condition of health is feeble, but she is able to do her household work and earn her living. Unusual exertion causes slight oedema of the feet. She has never had albuminuria.

Another case where equally good results followed, is the following. A. T., aged 46, married, was admitted into the London Hospital under my care September 3rd, 1875. She was quite well until two years before admission, when her feet began to swell. The catamenia, hitherto regular, at this time ceased. After the feet had been swollen two or three months, the abdomen began to swell; and it became so large that she came into the hospital eighteen months ago. She remained in the hospital three months, and during this time was twice tapped. On returning home, the swelling increased again, and she was again tapped. After this, the fluid again accumulated; she was readmitted into the hospital and tapped again. She improved, and continued as an out-patient until about eight weeks before her present admission, when her abdomen began to swell again, and it gradually increased, until she came into the hospital. There was nothing in her history to suggest syphilis; no excessive indulgence in alcohol could be elicited. The patient was a stout person, with injected cheeks, puffy face, watery and glistening conjunctivæ. The cardiac dulness could not