exudation of serum or rupture and punctiform hæmorrhages, according to the degree of intensity of blood-pressure? If, then, œdema be admitted as possible, the other symptoms of uræmia—the coma, drowsiness, and typhoid state—are completely accounted for by compression of the brain-tissue.

It may be objected to this, that uræmia occurs in chronic Bright's disease in which there is no dropsy. I have before pointed out a probable reason for the absence of dropsy in this condition; namely, the insidious nature of the disease, the gradual increase of the blood-tension, and the accompanying gradual thickening of the capillaries, which are thus enabled to withstand the increased blood-pressure. But, when this is still further suddenly increased, as by the supervention of an acute attack on the chronic disease, dropsy does occur, and so does uræmia; but, apart from this, the occurrence of a great increase of tension, causing distension of capillaries without exudation, would, by increased pressure on the nervous structures, produce some of the slighter symptoms of uræmia, such as the characteristic headache of Bright's disease.

To return, however, to the convulsions, other grave objections may be raised to my theory of their production.

1. Experimental evidence may be adduced to show that the grey matter of the cortex may be cut, bruised, burnt by caustics, and irritated by various methods, without the production of convulsions. But, on the other hand, besides its irritation by electricity, Nothnagel produced convulsions by the injection of a solution of chromic acid into the grey matter of the cortex. But, more than that—for experimental evidence, unless supported by pathological facts, is worth but little—I think I have already adduced sufficiently strong evidence of the association between punctiform hæmorrhages and convulsions to be found in the deadhouse.

2. It may be said that the convulsions produced the hæmorrhage, and not the hæmorrhage the convulsions; but the entire weight of pathological evidence is the other way. People dying of epilepsy do not usually present these hæmorrhages; in fact, all authorities on this disease emphasise greatly the fact of their great rarity. While in the deadhouse it is common to see ecchymoses below the skin, serous and mucous membranes, the result of impeded venous circulation, as in death by asphyxia from any cause, it is extremely rare to see these hæmorrhages in the brain-tissue under these conditions: though they will occur beneath the arachnoid, I have never seen them, nor can I find any record of their occurrence, in the substance of the brain, from this cause.

3. Epilepsy may be quoted as a fact against my theory, for not only are these hæmorrhages not found in epileptics, but epilepsy is believed by many eminent authorities to be associated with anæmia of the brain rather than hyperæmia; but this view is at least open to question. It would be impossible to discuss this subject here; but however strong the evidence of anæmia of the brain in epilepsy may be, the side of hyperæmia is almost equally strong. If Kussmaul and Tenner think they have proved *experimentally* that anæmia of the brain produces convulsions, so also Schroeder van der Kolk has beyond doubt, I think, proved that dilatation of the capillaries occurs as the result of hyperæmia in epileptics; and this, be it remembered, not generally throughout the brain, which might be the result of general turgescence owing to impeded venous return, but in that centre in the medulla to which he refers the irritation producing the convulsions. He is even able to divide his cases into two classes: tongue-biters, in which the capillaries of the hypoglossal nucleus are especially affected; and those who do not bite the tongue, in which those of the pneumogastric nucleus are alone dilated.

While suggesting capillary hæmorrhage as a cause for convulsions, let it be clearly understood that my remarks are confined to those occurring in conditions of high arterial tension, as in Bright's disease, pregnancy, and other conditions in which so-called uræmia occurs. The convulsions of epilepsy are probably due to some hyperæsthetic or irritable condition of nerve-cells, which may be due to a local hyperæmia, or to some other cause, but certainly not to capillary hæmorrhage.

4. Dr. Hughlings Jackson's theory of spasm of cerebral arteries as a cause for uræmic convulsions remains to be contested: it must stand or fall in great measure with the anæmic theory of epilepsy; but, even admitting that to be true, it appears to me difficult to conceive such severe spasm of cerebral arteries as to so completely cut off all blood-supply to the brain, as Kussmaul and Tenner's experiments seem to indicate as necessary for the production of epilepsy; for, according to them, if but one vessel out of the four large arteries supplying the brain be left uncompressed, the convulsions did not ensue.

5. Dr. George Johnson's theory of "stopcock action" of arterioles, and Dr. Gowers' recent ophthalmoscopic observations in support of this, may also perhaps be adduced as reasons against the hæmorrhagic

theory of convulsions. To take the latter observations first, I think they are open to error. Any one who has used the sphygmograph extensively, will recognise the fact that arterial tension may be high—that is, the tidal wave sustained—even with a relaxed condition of at least the larger vessels, while in other cases the arterial system throughout is contracted; every degree, from great relaxation to tight cord-like contraction of the vessels, may accompany increased arterial tension. In all alike, cerebral hæmorrhages are liable to occur. I think Dr. Gowers' observations can only show this contracted state of the vessels generally or their relaxation, but I do not think that they are any test of arterial tension.

If Dr. Johnson's view of the stopcock action of the arterioles be true, then the tension in the capillaries must surely be low, or not increased, owing to the defending contraction of the arterioles; and it is this fact chiefly that makes me unable to accept his otherwise ingenious theory, for, without admitting that the high tension extends to the capillaries, I cannot see how to explain bloody urine and dropsy in acute Bright's disease, and the hyaline thickening around capillaries and absence of dropsy in chronic Bright's disease.

These, then, are the reasons for and against this theory of the *vascular* cause of what are known as uræmic symptoms, as far as they occur to me. I think they would be better described as the symptoms of high tension in the cerebral capillaries, or, more shortly, as those of *cerebral œdema* or hæmorrhage. They must always be regarded as a precursor of apoplexy, if they occur in a person advanced in life or with degenerate vessels. Should my view prove to be the correct one, the indications it affords for treatment by rapidly diminishing the blood-pressure are of the utmost importance for the safety of the patient.

REPLIES TO DR. C. J. B. WILLIAMS AND TO DR. BROADBENT ON THE MECHANISM OF THE HEART-SOUNDS.

By ARTHUR LEARED, M.D., F.R.C.P.,
Senior Physician to the Great Northern Hospital, etc.

It is a matter of regret to me that the omission of a word in quoting from Dr. Williams' paper afforded him the occasion of making some sweeping and hardly justifiable assertions.* He says that my "misunderstandings and misquotations are something unusual; for example, he (Dr. Leared) quotes me as saying: 'The case is one of striking through a space at a tight barrier,' etc.; whereas my expression was: 'The case is *not* one of striking through a space at a tight barrier,' etc. This reversal of the sense makes nonsense, which he proceeds to deal with accordingly." In another part of the paper, the matter is again taken up. Dr. Williams says: "His omission of the word 'not' in the passage before quoted naturally throws Dr. Leared into a state of confusion, and it may be well to restate the argument in other terms." It is difficult to conceive why Dr. Williams could have allowed himself to make such a statement. But I am at the same time glad that he has put forward certain points of my theory with his own comments. To these I shall have occasion to return. First, let me dispose of the omission of a word and of its alleged effect upon my whole train of reasoning.

It is a remarkable circumstance that the insertion of the missing "not" in its place, instead of throwing me into "a state of confusion," strengthens my argument. I was combating Dr. Williams' assertion "that the whole movement is one of quiet displacement"; and I ask, has he "forgotten the force with which blood issued from the punctured aorta in his vivisection experiments? The force of the adult left ventricle was estimated by Valentin as sufficient to overcome a pressure of four pounds." Now, it is plain that, when I quoted Dr. Williams as saying the case was one of striking, instead of *not* striking, as it should have been, against a tight barrier, the force of my subsequent statement about the power of the ventricle was weakened by the omission of the word. Nevertheless, the mistake was unfortunate, and a cause of unnecessary trouble in this discussion.

Again, Dr. Williams says, in reference to the opinion of Dr. Stone, that "any person, whatever his scientific position, who declares the impossibility of sounds being produced in fluid alone, states a fallacy." "Either Dr. Leared misquoted, or his oracle misunderstood, my words, which were: 'I conclude, then, that the sounds of the heart are not produced by any motions or collisions of the blood, because I find that no such sounds can be produced in fluids alone.'" But, in the paragraph

* Dr. Williams' paper appeared in the BRITISH MEDICAL JOURNAL, February 3rd, 1877.—In Dr. Leared's absence from town, and from pressure on space, the insertion of this paper has been unavoidably delayed.

in my reply to Dr. Williams, in which this last sentence is quoted (BRITISH MEDICAL JOURNAL, December 2nd, 1876), the following sentence (also a quotation from Dr. Williams) goes immediately before: "The whole idea of noisy collision of fluids without air or solids is a mistake." This omission, when supplied, disposes of the qualified expression in the sentence put forward by him, and gives due force to the statement made by Dr. Stone.

It is quite clear, therefore, that, in this instance, the charge of either misquoting or misunderstanding is untenable, and that the context of the sentence quoted by Dr. Williams shows he did say more than he appears to have wished to say.

The points which require comment in the restatement of my theory, which Dr. Williams has been good enough to make, are contained in the following extract: "He (Dr. Leared) made the experiment of forcibly squeezing water from an India-rubber bottle through the side of a reservoir containing water; the result was a bellows-murmur at the orifice. On substituting glycerine for water, the bellows-sound was not heard, and there was difficulty in producing any sound" (I wrote that there was "a comparative difficulty in producing sound"); "but, when the India-rubber bottle was compressed very strongly, a faint sound, resembling the first sound of the heart, was heard near the mouth of the tube; when, however, the moving column was allowed to strike the stethoscope, a louder sound, exactly resembling the heart's first sound, was heard." Upon this, Dr. Williams remarks: "The experiment with glycerine proves my statement of the difficulty of producing sounds in liquids, especially in those that are viscid; but, when the moving column struck against a solid (the stethoscope), the sound was loud enough."*

Now, I wish here to observe that the circumstance of the sound formed in water—current-sound—and that formed in glycerine—shock-sound—having been intensified respectively by the contact of the moving fluid with the stethoscope, does not in the least invalidate my views. What I proved was, that sounds of very different quality are capable of being produced in two liquids, one being possessed of viscosity and the other being free from viscosity. But I have never denied the influence of solids in modifying, intensifying, or conveying sounds so formed. Dr. Williams may argue that the sounds in the cases in question were caused by contact between the liquids and the stethoscope. Even if this be granted, the results remain to be explained; namely, that, in the one case, the sound was a current-sound, in the other, a shock-sound. The bearing of this upon my theory, that shock-sounds and current-sounds merge into each other, is obvious. Is it not a pathological fact that the practised ear finds it difficult at times to decide whether a prolonged first sound or a low-pitched murmur is heard? Does it not

* At this part of Dr. Williams's paper there is a foot-note, which, to say the least, is not characterised by good taste: and it so happens that two other notes are placed immediately under the first mentioned, each being marked, somewhat after the sortie method, by an increased deficiency of the particular in question. In note No. 1, Dr. Williams says, "Dr. Leared seems not to be very successful in his experiments." He says (page 11, note) "I have laboured hard to produce an artificial circulation, which should fulfil all the requisite conditions for the formation of sounds like those of the heart by means of an India-rubber apparatus, but hitherto without success as regards the first sound." The final sentence, "The delicacy and nicety of adjustment necessary in the valves form the great difficulty", is omitted by Dr. Williams. Any one may see from a letter of mine in the *Medical Times and Gazette* of April 7th, 1866, how I was thwarted in my efforts to get such an apparatus manufactured, and how, after expecting it for several months, I was coolly informed that it was not practicable. Soon after this, a caoutchouc instrument, to which the name of the serpent's heart enema-apparatus was given, in which not only was the principle of my intended apparatus followed, but the very shape of the heart-propeller closely copied, was brought out by the firm in question. This instrument, ever since in extensive use, was the result of the act now briefly explained. So much for my unsuccessful experiments.

In note No. 2, speaking of a proposed experiment of mine, Dr. Williams allows himself to write: "I venture to prophesy that this will prove another of his unsuccessful experiments." If the prophet in this instance be inspired, it is to be feared that the inspiration is not of the best quality.

In note No. 3 it is said, "Disregarding my expression 'momentary separation', Dr. Leared erroneously supposes me to mean that, palpable or permanent air is evolved, and proceeds to attack this man of straw." Dr. Williams's words were: "This implies a slight tendency to a vacuum, and momentary separation of the surfaces by gas or vapour." Now, if not "permanent", "palpable" gas or vapour is here implied; and there is, therefore, something more to be attacked than "a man of straw".

"It is unnecessary to follow him through his mistaken argument" (proceeds Dr. Williams), "but it may be remarked that his statement, that the presence of air in the arteries destroys the second sound, is readily explained by its interfering with the action of the semilunar valve, which is adapted for liquids only. It is thus that injection of air into blood-vessels destroys life." I question the statement that, even supposing the second sound to be strictly due to valve-vibration, the presence of air would destroy these vibrations.

The subject of mistakes suggests that this is a good opportunity for bringing one of Dr. Williams's own making under his notice. He says "liquids being more sluggish than air, are less susceptible of the sudden motions which constitute sonorous vibrations". Now, so much is this contrary to fact, that, according to Biot and others, the velocity of sound-propagation in water is more than four times more rapid than in air.

happen that the first observable change preceding the development of a murmur is sometimes prolongation of the first sound?

Dr. Williams admits that sounds of very different qualities are formed in the blood-vessels, not only by obstructions in these vessels, but by alterations in the blood; but he says: "I repeat, the normal sounds of the heart are totally distinct from murmurs." He admits that sound exists potentially in the blood propelled from the ventricles "during the mighty systole", to use his own expression, because not only are murmurs formed by obstruction to the outflow, but also when, without obstruction, the blood is thinner than natural. So far, we agree, and the point of difference between us lies in this. I maintain that not only is thin blood, when forcibly driven against thin blood, capable of being sonorous, but that thick blood, when forcibly driven against thick blood, is not silent, but produces sound of a different quality. In the one case, the facility for friction between the opposed fluids gives rise to a current-sound; in the other case, the interference with friction is the cause of a shock-sound. Nothing, I submit, which Dr. Williams has advanced has shaken this fundamental part of my argument or destroyed the evidence afforded by repeated experiments. I cannot even accept his brief but forcible summing-up of the matter, when he says, "Having negatived the liquid solution of the question", as conclusive.

In reference to the analogy between the noise made by turning a tap through which water is flowing and the second sound of the heart, Dr. Williams writes: "As he (Dr. Leared) has quite misunderstood the too concise explanation which I gave, I think it necessary to compare the two cases more fully." I have carefully read what I wrote, and am quite unable to find any evidence of misconception, unless it be that difference of opinion be taken as synonymous with misconception. However, Dr. Williams explains his views in detail. He endeavours to show that there is no analogy between the two cases, because the noise in the case of the water-tap is only heard when the pipe is several feet in length, and there can be no such pressure in the circulation. But there is considerable pressure and the question of sound becomes one of degree, a comparatively small pressure sufficing to produce a sound only as loud as that of the heart's second sound.

In my former reply to Dr. Williams, I confined myself as much as possible, as I intend to do at present, to his criticisms of, and objections to, my views. I was unwilling to discuss the merits of his theories, both on account of his seniority and the undoubted value of some of his researches. I hoped that he would in the same friendly spirit discuss the subject with the sole aim of arriving at truth. It did not occur to me that he would treat the matter as one finally closed, and his decisions as such against which there could be no appeal. I was myself familiar with the merits of the controversy between himself and Dr. Hope. But I felt that the affair was so long gone by, that this was not the case with the great majority of the profession. Certain unqualified expressions used by Dr. Broadbent make it certain that he was in this position. But a Nemesis has arisen in an unexpected quarter. In an article founded upon the present discussion in the pages of a contemporary (*The Doctor*, March 1st), in which my theory is not supported, several of Dr. Williams's statements are questioned, and his claims to originality in the matter of the heart-sounds in general are treated with an unsparing hand. Dr. Williams lays much stress upon his invention of terms, etc., and, as he found the terms "shock-sound" and "current-sound" employed by me useful in his explanations, he takes care to inform the reader that I borrowed these terms from French writers. Shock-sound was adopted from Gendrin; but I beg to assure Dr. Williams that current-sound, which conveys the idea derived from my theory that the sound is formed in and by the blood itself, is altogether my own. Dr. Williams thus begins what is intended to be crushing.

"But the most conclusive proof that the walls of the ventricle, and not the blood, are the seat of the systolic sound is found in those observations in which the sound was heard when the heart contained no blood." Here it will be convenient to meet the statements of Dr. Broadbent on the same subject; for the latter is quoted by Dr. Williams in support of his position. To both observers, I have to say that we are still totally at variance on the question of fact. Notwithstanding Dr. Broadbent's clever badinage about the thing being "only absolutely impossible to me", I still maintain the impossibility of deciding whether the feeble sound heard on the application of a stethoscope to the empty but still contracting heart is not due to friction between the heart and the instrument. "But", says Dr. Broadbent, "Dr. Sibson found...that there is a point on the anterior aspect of the heart...which is itself almost motionless. It was here that the stethoscope, which was flexible and had a small cap, was applied." There is much virtue in an *if*, and so, in the above sentence, is there much

* Vide Dr. Broadbent's reply, BRITISH MEDICAL JOURNAL, February 10th, 1877.

virtue in *almost*. Let the reader experiment with a flexible or a rigid stethoscope; let him moisten the bell-end as suggested; let him contract the muscles of his abdomen, of his arm, or of his hand; and let him do this so carefully, that the surface to which the instrument is applied is *almost motionless*, and I will answer for it that a sound or an impulse will be conveyed to his ear. And I will further answer for it that a similar sound would be produced by the contracting empty heart, and that, under the influence of a foregone conclusion, this would be regarded as the expiring remnant of the true first sound of the heart.

Dr. Broadbent writes in reference to this matter: "The statement which he quotes from Dr. Halford, 'that neither first nor second sound was heard', is beside the question. Dr. Leared must have heard sound under these conditions, as I have, or his assertion about it was simply unmeaning. Had he come to the same conclusion as Dr. Halford, I should probably never have written my remarks at all." There is some confusion in this statement. I quoted Dr. Halford's assertion that, in the contracting empty heart, "neither first nor second sound was heard" with the conviction that this involuntary testimony of a man who was supporting a theory differing both from Dr. Williams's and my own was very much to the question. If, under the circumstances mentioned, sound was heard by Dr. Halford, he presumably attributed it to friction, because he distinctly says no sound of the heart was heard. What is intended to be conveyed in the statement about this experiment of Dr. Halford, that I must have heard sound under these conditions, or my assertion about it was unmeaning, I really cannot unravel. Then follows the strange assertion that, had I come to the same conclusion as Dr. Halford, his (Dr. Broadbent's) remarks would probably never have been written. Now, if "the same conclusion" refer to the origin of the sound in the empty heart, that has just been disposed of. If it refer to the valvular theory as advocated by Dr. Halford, it is "beside the question". It can hardly be supposed that Dr. Broadbent means that he would be *equally* satisfied with the muscular or the valvular theory, provided only my theory be excluded. A special question raised by Dr. Broadbent is the following:—"If we deny to the valves any share in the production of the sounds, we must positively explain their silence. I tried in vain to obtain Dr. Leared's attention to this." It will give a fair idea of the unnecessarily strong language in which my friend has constantly indulged in the discussion to quote the only statement of his to which this allegation can be referred. "I do not understand Dr. Leared to deny that sudden tension of membranous valves and tendinous cords must be attended with the production of sound." I passed this over only because there was so much to be discussed, and because it seemed of little importance, far less, for instance, than Cruveilhier's case and other things about which I failed to elicit any opinion from Dr. Broadbent. I hasten, however, to make good the omission in as few words as possible, although much could be said on the subject.

The reason, then, why the valves do not yield the essential portions of the heart-sounds is this. When in action, they are equally supported at both sides, not being adapted for the rough work implied by the theory of comparatively loud sound-producing vibrations. Such valves are well known in the arts by the name of equilibrium valves. It is observable that, in their zeal for the muscular theory of the first sound, no stress is laid by Dr. Williams on the valvular theory and very little by Dr. Broadbent. It is now and then spoken of as an accessory and in what may be termed a "permissive" manner. And yet this valvular theory so ably advocated by Dr. Billing, compared with the muscular theory, has the great recommendation of making the cause of both sounds identical in kind.

Here I take the opportunity of saying that I have never denied the possibility of *some* sound being produced by muscle and valve. In this matter, I follow the rule *de minimis lex non curat*. What I maintain is, that the body or essential part of the heart's sounds is not caused by one or both of these factors.

Some of the objections raised by Dr. Broadbent in the form of questions (BRITISH MEDICAL JOURNAL, December 23rd, 1876) have been commented upon by Dr. Williams. "If", he says, "the large arteries are the principal seat of sound, the sound would be increased by an hypertrophied heart and diminished by a dilated heart; whereas the contrary is the fact. Also that, when arterial pressure is increased, the first as well as the second sound ought to be louder, according to my views." These points have been considered in my former reply to Dr. Broadbent, which Dr. Williams had evidently not seen when his last paper was written.

In reference to my remarks on the difficulty connected with deductions resting solely upon position of sounds, Dr. Broadbent writes: "I see no end to the discussion, if we must cease to accept the recognised methods of localising the source of a sound whenever this is re-

quired by the exigencies of Dr. Leared's theory. Errors arising from the conduction of sound do not prevent us from forming a confident diagnosis in disease." The statement in the first of these sentences is uncalled for by anything I have written as to that in the second. I have only to observe that Dr. Broadbent is lucky if he have not, in common with the best auscultators, been now and then betrayed, as determined after death, into a too confident diagnosis when founded on the localisation of murmurs alone. The new theory of anæmic murmurs of Dr. G. Balfour, founded on conduction alone, was mentioned by me, which Dr. Broadbent would probably not endorse; at all events, he passed it over without remark.

The last but most important thing to be considered is what is said about my question, Why is it that, in anæmia, *the first sound of the heart only*, and the second never, becomes replaced by a murmur? Dr. Broadbent reiterates his previous argument, in which he is followed by Dr. Williams almost *totidem verbis*. That argument is, that the normal first sound is not replaced by, but is accompanied by a basic murmur. It was Dr. Broadbent who made the *supposed* discovery. "To this my objection admitted by Dr. Leared was fatal." "When (Dr. Broadbent writes just previously) he has restated the proposition put by him as a *crux* which I challenged, with the modifications required by his admission that the murmur accompanies instead of replacing the sound, I will again consider it." There is more smartness than logic in much that Dr. Broadbent writes. There is no necessity for any restatement of my proposition. Dazzled by the imaginary brilliancy of the discovery, he was unable to see the real bearings of the case, and my frankness only tended to increase his difficulty. My question is put in terms sufficiently definite for all practical purposes. It is affirmed of the second sound alone that it is never replaced by a murmur, and of the first that it only is liable to be changed into a murmur. I do not know that it is stated by any standard authority that, in some instances, this murmur is accompanied by a shock-sound. Now, with regard to this shock-sound, I agree with Dr. Broadbent that the relative loudness of sounds is not a simple and definite matter for discussion. If it were, I should join issue with him on this subject. Perhaps some of his "exceptional cases would be an interesting clinical study". This was said by him in reference to my statement (to which I still adhere), that instances occur in which the first sound of the heart is louder at the base than at the apex. But *de sonibus* (like *de gustibus*) *nil disputandum*; at all events, about such sounds and on paper.

If I had written, "Why is it that, in *some* cases of anæmia, the first sound of the heart alone undergoes the change in question?" Dr. Broadbent's objection would have had no more point than as the case stands; for I deny the universality of his statement that the anæmic systolic murmur is always accompanied by a shock-sound. If, then, it be a fact, as I maintain it to be, that the normal basic first sound is sometimes replaced by a low-pitched anæmic murmur, not drowned by a loud rasping murmur, such as is common in organic affections, my position is safe.

The coexistence of a shock-sound with an anæmic murmur is capable of explanation in various ways. It is possible that the sound produced by the blood issuing from the ventricles may sometimes be split into normal and abnormal sounds. More probable is the occurrence of the persistence of the normal sound at the outlet of one great vessel, while it has degenerated into a murmur in the other; and that the normal and abnormal sounds are heard together. Finally, there is the possible agency of other sources of sound, which I have never repudiated.

Dr. Williams says "that, in some cases of anæmia, I have found a short murmur preceding the second sound". As a matter of fact, I do not question the accuracy of the observation, although I believe it to be peculiarly his own. The sound described by Dr. Williams is pre-diastolic, precedes the second sound, and does not either replace or accompany that sound, and, therefore, does not interfere with my statement that the second sound is never replaced by a murmur in anæmia.* Nothing, I maintain, has shaken the significance of the solid fact stated in my original proposition. It would, indeed, be a different matter if the converse of the proposition could be proved; namely, that, while the first sound always remained a shock-sound in anæmia, the second sound was sometimes changed to a murmur; for my argument is, that, if a thin fluid like water circulated in the body, there would be no first sound such as is formed by the circulation of healthy blood, but, instead of it, there would invariably be a murmur; but that, under the same circumstances, the second sound would still be a normal shock-sound. A shock-sound is sometimes, not always, heard along with the anæmic cardiac murmur. As there is sometimes no sound heard but the murmur, this *must be* a substituted sound and virtually proves my

* On this point Dr. Walshe says: "I have never encountered a positive example of aortic blood-murmur diastolic in time" (*A Practical Treatise on the Diseases of the Heart and Great Vessels*, London, 1873, p. 378).

case. If more or less shock-sound be heard with the murmur, this does not disprove my case. The questions, then, arise, What is the source of this shock-sound? Is it the true heart-sound, or is it only an appendage to the murmur which has replaced the normal sound? The present theory does not rest for support on the advocacy of its author. It has many adherents, and amongst them such eminent men as Dr. Walshe. His published works and his well earned reputation as one of the ablest and most accurate observers with regard to everything that concerns auscultation and physical diagnosis that this country has produced, makes his testimony of special value. I may be pardoned, therefore, for giving the following extract from his last work. After having considered the merits of the muscular theory of the first sound as well as several others, including my own, he says: "In fine, the theory advocated by Dr. Leared appears more in accordance with the principles of physics and better reconcilable with numerous facts of clinical experience (many of which will appear in the sequel) than any of its rivals."*

Considering the turn which this discussion has taken, I have no wish to prolong it; but that matter does not rest with myself alone, and, if necessary, I shall be again ready to maintain what I believe to be the true theory of the sounds of the heart. The rule observed throughout has been to meet fairly every objection brought forward.

SURGICAL MEMORANDA.

DISLOCATION OF THE TIBIA FORWARD.

HAVING noticed in the JOURNAL for May 5th the record of some cases of rare dislocation by Mr. Bradley of Manchester, I am induced to communicate the details of the following case, in the belief that they will be of interest to him, and perhaps to the profession generally.

On March 31st, 1875, while in practice in Cape Colony, I was requested to proceed to a farm distant about three hours' ride from the village in which I resided, to see a man who, four and a half days previously, had been kicked by an ox. I found the patient suffering intense pain, lying in a hut near which he had been working when the accident occurred. In the region of the left knee-joint, there was extreme deformity, evidently due to displacement of the bones of the leg forward; the tibia was riding on the front of the lower end of the femur, while the latter appeared to be rotated inwards, as the internal condyle was pressing strongly on the skin in the popliteal space. A circular spot of skin, black, hard, and dry, was adherent to this condyle, and followed it when reduction was accomplished. The leg was shortened to the extent of three inches. The foot and leg were cedematous, but much less so than might have been expected. There were several large bullæ, full of bloody serum, on the outer and posterior aspects of the joint; the largest of these being on the spot at which the blow had taken effect—immediately behind the head of the fibula. By extension and manipulation, the dislocation was readily reduced without the aid of chloroform. The parts behind the joint were dressed with a watery solution of carbolic acid, and the limb put up in an improvised long splint, without extension.

I left the colony immediately afterwards, and saw this man no more; but he ultimately did well, as a friend, writing to me three months later, stated that — had repeatedly sent for more lotion, and was last seen hobbling about leaning on a stick.

ALEXANDER NAPIER, M.D., Crosshill, Glasgow.

THERAPEUTIC MEMORANDA.

THE EMPLOYMENT OF LIQUOR AMMONIÆ ACETATIS IN DYSMENORRHEA.

THE following case may prove interesting, as affording additional evidence of the usefulness of liquor ammoniæ acetatis in painful menstruation.

On June 22nd, at 6 A.M., I was called to see M. C., aged 19, who was stated to be in great pain. On arrival, I found the patient writhing in agony, her face pale, and the body covered with profuse perspiration. She is a healthy-looking and well nourished girl, but not of a marked plethoric habit; and I was informed that, with the exception of the dysmenorrhœa, her health had always been good. Since menstruation commenced, four years ago, she has always suffered more or less at her periods, but never so much as on the present

occasion. I was told that menstruation had come on some little time before I was called, and that a slight "show" had appeared. I prescribed a drachm of liquor ammoniæ acetatis with a little spirit of chloroform every hour until relieved. On visiting the patient again in the course of a few hours, I was informed that she had taken two or three doses of the medicine with the most satisfactory results, and she now expressed herself as quite free from pain.

In the afternoon of the same day, the pain returned slightly, but was again relieved by the medicine, and since then there has been no recurrence.

I find that the same satisfactory results have been obtained by my brother, Mr. C. W. Drew, previously in the same patient, and also in other similar cases where everything else has failed. How the liquor ammoniæ acetatis effects this result is not quite clear, but certainly, in some cases, it seems to act as a specific. It may be that the same influence is exerted on the uterine surfaces as on the cutaneous, increasing secretion, making the flow more free, and thus lessening the tension of the engorged vessels.

CLIFFORD L. DREW, M.B.

REPORTS

OF

MEDICAL AND SURGICAL PRACTICE IN THE HOSPITALS AND ASYLUMS OF GREAT BRITAIN.

HOSPITAL NOTES.

ST. THOMAS'S HOSPITAL: MR. MACCORMAC'S WARD.

Excision of Hip-Joint.—A boy, delicate-looking but playful and free from pain, had undergone excision of the hip-joint a few weeks before. Previously to the operation, he suffered great pain, and hypodermic injections of morphia were needed daily. General constitutional disturbance had been marked, and signs of commencing phthisical changes were found in both lungs. Immediately after the operation, pain ceased and his general health improved. Another little boy had just been admitted for hip-joint disease, with suppuration. There was considerable shortening of the limb; the head of the femur appeared to have been absorbed and sinuses existed. On examining the pelvis through the abdominal walls, much inflammatory thickening could be felt on its inner surface opposite the acetabulum; evidently the inflammation, and probably the suppuration, extended through into the pelvis. Mr. MacCormac remarked that, in such cases, it is usually the junction of the three portions of the innominate bone at the acetabulum that gives way; the pus then enters the pelvis, and, as it accumulates there, tends to strip up the periosteum from the bone. Hence this swelling in the pelvis is an indication for excision rather than against it, as is often taught. After excision, there will be free exit for the pus; removal of the dead bone will, it is hoped, lessen the discharge, and the sections of healthy bone may produce fibrous union. Directions were given for a careful examination of the lungs and urine to be made previously to a final decision on operative procedure.

Hæmophilia.—A man aged 42 had received a blow on the cheek, and dangerous hæmorrhage resulted from the socket of the last molar tooth. On admission, the blood welled up continuously, but not in large quantities. Various styptics were used without success during the first week. The loss of blood amounted to about half a pint a day, and threatened life. Another plan was then tried. A piece of India-rubber was rendered soft in hot water, placed over the bleeding jaw, and the mouth firmly closed upon it; it was then moulded so as to form a case over the bleeding point; pressure was thus effectually made, and the bleeding has not recurred during the last week. Ergot has been given internally, and ergotine by hypodermic injection. The patient has five brothers, of whom three are "bleeders"; his two sisters are not so. Neither his father nor mother are thus afflicted, but his mother's father, uncles, and aunts were "bleeders". Such conditions seem to be transmitted to children by the mother rather than by the father, and a mother may transmit the tendency without appearing unhealthy herself.

Cicatrix after a Burn.—A boy was admitted with a very extensive granulating surface on the right side of his chest, passing up to the axilla, as the result of a burn. There was as yet no cicatrix, but it was certain that such would form and tie down the arm. A month ago, Mr. MacCormac dissected a long flap of skin from the dorsal region, and brought it horizontally forward under the arm on to the granulating surface, leaving its upper extremity at the spine. A part of this flap

* *Op. cit.*, page 57.