would just remark upon the great danger of stamping everything as diphtheria where there is found fibrinous exudation. We know that it is found as a result of scald from hot water, and I have seen a throat after the removal of tonsils that I could not have distinguished from diphtheria.

Three years ago, my coachman lost his voice, and I found a thick tough exudation over the whole of the posterior wall of the larynx. There had been bad measles in the house in which he lodged, and bad drainage. There was no glandular enlargement or puffing of the cellular tissue. I believed it to be diphtheria, and treated it as such; but it was bad for many weeks, and only yielded to iodide of potassium or warm weather. There was *no history whatever of syphilis*. The next winter, it was just in the same condition; and this year again it was like the two preceding winters, although each time he was living in different houses and under different circumstances, and no other members of his family had suffered in a similar manner. My coachman attributed it, and I now think with reason, to the exposure to cold, as he is never in any way ill excepting in the bad weather of winter.

In bad scarlatina, in bad measles, and at the end of many exhausting diseases, membranous exudations are to be found; and I think it very probable that of late years many cases have been attributed to the specific contagion of diphtheria, which really owned no such origin.

MALARIAL FEVER AT ROME.

By LAUHLAN AITKEN, M.D., Rome.

THE existence of a malarial element in the climate of Rome is so generally admitted, that it is quite unnecessary to enter into the history of the subject, to point out the foci of infection in the Campagna, to discuss the different effects of the poison, or even to mention the many varieties of malarial fever which authors describe. In this paper, I shall only deal with the effects of the malaria on unacclimatised strangers, English or American, who have been under my care. So dormant is the poison during the winter and spring months, that acclimatised members of the nationalities mentioned, who can afford to live well and to reside in a healthy quarter, are, so far as my six years' experience goes, never affected; but visitors, who are careless of consequences and heedless of the ordinary warnings, often overwork themselves in sight-seeing, become heated in the sun, and stand for hours in museums, galleries, and churches, exhausting their nervous systems, or expose themselves more directly on the Campagna, or even within the walls of the city, to a chill which results in an attack of so-called Roman fever. What this fever is the residents are often very puzzled to know, as strangers include under the generic term all kinds of ailments from a simple cold to a severe enteric fever; and it is sufficient to fall ill in Rome to be credited everywhere with being a victim of that vague disease. The result is naturally a wide divergence of views on the subject, and an alarm proportionate to the amount of confusion created. *Omne ignotum pro terribili* is the idea of all who speak of Roman fever.

That the malarial poison does produce, more particularly in the unacclimatised, a type of fever which can scarcely be classed under any such heading as intermittent or remittent, is certain. To this form the name "subcontinuous typhoid" has been given by Professor Baccelli of Rome. The term, fortunately, has not as yet become widely known; for, though the type of fever may be regarded as subcontinuous, yet, in whatever sense we take the word typhoid, the epithet is misapplied. If we regard it as used to express the group of symptoms called typhoid, or the typhoid state, my experience contradicts such an use, as none of the forty-five cases I have treated throughout have presented those symptoms which British physicians at least class under this heading. On the other hand, it is evident that Baccelli does not look on this fever as the product of the union of the malarial poison with that of decomposing excrementitious matter, as he carefully distinguishes his subcontinued typhoid from true ileo-typhus (enteric fever). Indeed, Baccelli seems to contradict himself in his lectures on the subject, as in one part he says that "the two morbid entities—the typhoid and malarial poisons—each losing its autonomy, become so fused together, but with the malarial predominating, that they give rise to the special type—subcontinuous typhoid"; while in other pages of the same pamphlet he asserts that "the subcontinuous typhoid is a fever resulting from malarial infection, pernicious as to type"; and again, "that, when a subcontinuous typhoid breaks out, every other morbid element, rheumatic, bilious, or typhoid, becomes subordinate to the malarial". It is, therefore, difficult to determine in what exact sense Baccelli uses the term typhoid; and I prefer to regard those

fevers as truly malarial, though presenting a remarkably continued course.* I am far from denying the occurrence of the combined type of fever which may appropriately be called typho-malarial. Indeed, I have treated such cases; but, as might be expected, the more severe poison very soon predominates, and the bowel-symptoms and eruption make the diagnosis of an enteric fever clear.

Etiology.—The fever to be described results, then, from malarial poisoning; the predisposing causes being, in the cases I have seen, unacclimatisation; lowered vitality from anxiety of mind, from chronic diseases, or from previous acute complaints, particularly the malarial; pregnancy and the puerperal state; or a combination of some of these; while the exciting cause in every instance I have known has been a direct chill when the body was overheated, or its power of resisting cold diminished by any such causes as have just been mentioned, or by exposure during the night, when the heat of the body is naturally lower. A few lines from my case-books will illustrate this.

E. P., a young lady aged 22, goes to an evening party after a fatiguing day's sight-seeing; dances until well on in the morning; stands, when perspiring, at the door awaiting her carriage; gets a chill, and the next day is laid up.

J. G., a gentleman aged 28, has a hard day's work after the hours on the Campagna, arrives much overheated at one of the gates after sunset, dismounts there, and drives to his hotel in an open cab without putting on a greatcoat. Two days afterwards, he shivers, and he has a somewhat prolonged fever.

S. S., a young lady aged 17, just convalescing from bronchitis, sits at an open window one cold afternoon in December, gets chilled, and has the most severe attack yet treated by me.

The Rev. Mr. D., aged 30, reading with some Oxford students, has one of the headaches to which he is liable; and, after a long heating walk to get rid of it, he sits down about sunset on the grass outside one of the gates, and gets what proves to be a mild attack.

D. W., aged 19, a young gentleman, rising one morning in December with a stitch in his side—dry pleurisy it turns out to be—takes his cold bath as usual, does not react at all, and has also a mild attack, which masks the pleuritic symptoms entirely for more than ten days.

An exactly similar account of the origin of the complaint might be given in nearly all the forty to fifty cases mentioned.

It is certain that the young are much oftener the victims than those advanced in life, which partly arises from their greater imprudence, and partly from their greater predisposition to contract fevers. So much is this the case that, though quite as many, if not more, elderly people come to Rome, yet among my cases four only have been over forty years of age. So far as sex is concerned, the numbers have been pretty equal; and the severity of the attack has apparently depended more on the previous state of the patient than on the fact of the chill having occurred within or without the walls of the city.

Incubation.—This could not be determined in every case; but, in nearly all in which the time of getting the chill was definitely fixed, it did not extend over two or three days. In several, the first symptom—more or less decided shivering—began within a few hours. In the young lady mentioned, in whom the attack followed a bronchitis, the temperature was 102.2 deg. within seven hours from the time of the chill. In another young lady, the shivering and headache began within a few hours from the time at which she had felt cold when sitting in a draught at an afternoon tea. In one gentleman (over forty), the incubation-stage seemed much longer. He was ill for about eight days with indefinite symptoms; loss of appetite; furred tongue; urine laden with lithates; evening headache and disturbed sleep, but no rise of temperature—pointing, as I supposed, to functional liver-disorder, but quite unrelieved by the treatment adopted; when he was sent to Albano, where he became worse instead of better, and from which he returned after a few days with a well-developed attack. At what exact period of his stay at Albano, the fever first showed itself, is uncertain. It may be objected, that the symptoms previous to the appearance of the fever did not necessarily constitute the incubation-stage; but I incline to think they did, from the non-specific treatment adopted doing no good at all. In the other three cases of elderly people being affected, I was unable to fix the date of the chill. Indeed, two of the patients had been under the care of other physicians before I was sent for; and in the fourth I could not definitely determine the length of the incubation, which probably varies with the age and power of resistance of the patient, as well as with the dose of the poison. General languor, an inclination to sigh and yawn, insomnia, aching pains in the muscles, and loss of appetite, are symptoms common to the incubation-period of all forms of fever.

* Although venturing to differ from Professor Baccelli as to the name of the fever, I would here express my appreciation of the great value of his communications on the subject.

Symptoms.—The fever begins usually with more or less decided shivering, but never, in my experience, to the extent of the chattering of the teeth, blueness of the lips and face, and shaking of the whole body, seen in an ague-fit. Sometimes there is only the feeling of cold water running down the back, and occasionally nothing more than a little shuddering or the sensation of goose-skin. In one young lady, aged 24, a prolonged syncopal attack was the first symptom, recurring more or less every day for about ten days at the same hour. This daily recurring faintness I saw in another young lady, in whom the fever began in the regular way with shivering. Neither of the patients had ever been subject to fainting fits, which obviously took the place of the cold stage of an ague. It may begin as a genuine intermittent, and take the subcontinued form. Baccelli asserts that it often does so, but I have never seen it amongst the strangers, though I have known it terminate both as a quotidian and a tertian intermittent. Headache is invariably present in the beginning of the attack, always frontal and quite distinguishable from the neuralgiae of the ophthalmic divisions of the fifth nerve, supraorbital and ocular, which pressure on the supra-orbital notch and on the eyeball, and the attempt to rotate the eyes, often prove to exist. Occasionally, the neuralgia is occipital; and in one old lady, aged 65, violent unilateral neuralgiae of the branches of the fifth and of the cervico-occipital nerves, of an intermittent character, constantly reappeared during a somewhat prolonged attack of the fever. The eyes are generally congested at the beginning, while, during the fever, the conjunctivæ become yellowish, as Baccelli also observes.

Few of the nerve-symptoms were prominent, either in the beginning or throughout the complaint. Delirium was rare and transitory, except in my two youngest patients, aged 14 and 17 respectively. Slight confusion of ideas, marked listlessness, and sleeplessness, were common to all, but no deafness apart from that caused by drugs.

Temperature.—The thermometer shows a sudden and rapid rise of temperature, after the initial symptoms; from 102 deg. to 105 deg. being very commonly found quite early in the complaint, within a few hours from the shivering, for instance; and until means are taken to diminish it, either by cold applications or by the use of quinine, this increase of heat virtually remains unchanged. During the progress of the fever, however, there seem to be rapid fluctuations of temperature several times in the twenty-four hours, and the remissions are not invariably in the morning. Indeed, the temperature in several of my patients has been lower in the evening than in the morning, and that for many days consecutively; but, as all my cases have been treated with antipyretics—quinine, usually in large doses—the charts only indicate the course of the fever thus modified.

Case of E. P., an English lady, aged 23. First seen on the evening of the second day of the fever. The case was of average severity, and there were no complications. She was treated throughout with quinine, but not quite in such large doses as usual.

Days.	M.	E.	Days.	M.	E.
2	..	102.6	12	..	100.0
3	..	101.6	13	..	100.1
4	..	102.0	14	..	99.2
5	..	101.6	15	..	99.0
6	..	102.0	16	..	98.8
7	..	100.6	17	..	98.9
8	..	101.5	18	..	98.9
9	..	100.5	19	..	99.0
10	..	100.5	20	..	98.8
11	..	101.4	21	..	98.8

S. S., aged 17, English. Case severe; complicated with constant vomiting at the beginning, and bronchitis from the tenth day of the fever. Subcutaneous injection of quinine from the fourth to the eighth days of the fever. Reported at length in the BRITISH MEDICAL JOURNAL for April 1st, 1876.

Days.	M.	E.	Days.	M.	E.
1	..	102.5	10	..	100.9
2	..	100.5	11	..	101.0
3	..	100.5	12	..	101.1
4	..	102.0	13	..	99.5
5	..	102.1	14	..	98.8
6	..	103.1	15	..	98.7
7	..	99.5	16	..	98.7
8	..	101.0	17	..	98.5
9	..	101.0			

A typical case ought to have the temperature indicated every few hours, but in private practice that is nearly impossible.

Circulatory System.—The circulation never seems so rapid as the rise of temperature would lead us to expect. Even quite at the beginning of the attack, the pulse is seldom more than 100 per minute, and, as soon as antipyretics are absorbed, it falls, so that it is quite usual to find the heart beating only from 70 to 90, with a temperature at from 101 deg. to 104 deg. It is nearly always quicker in the evening than in the morning, but not more so proportionately than in health. Feebleness, irregularity, and intermittency have often been observed after the

fever had gone on some time. Occasionally, there is congestion of the cutaneous vesicles of the face, neck, chest, and abdomen; and I have seen limited erythematous patches, in two or three cases, over the same parts; but there is no eruption characteristic of the complaint.

Sweating is a nearly constant symptom. As a rule, it has been moderate in amount and has come on in the early morning hours, but very often it has appeared at other times without any apparent cause. In some patients, it has been a symptom throughout; while in others, it has first shown itself towards the end of the complaint. In only a very few has it been so excessive as to require the use of waterproof sheeting to protect the bedding.

Digestive System.—The tongue from the beginning is covered with a thick white fur. I have notes of the occasional appearance of red prominent papillæ at the tip, but never of its being red all over, nor of cracks or fissures on it at any time, nor of any hæmorrhage from the tongue or gums. In one or two of the younger patients, there has been epistaxis, and the lips have bled, but not to any extent. When the patient improves, the white coating begins to disappear from the tip and edges first. The mouth is rarely dry, and patients do not complain much of thirst. The appetite is lost, but nausea and vomiting are unfrequent; though, in one severe case, vomiting was the leading symptom in the beginning of the attack. The liver and spleen are both enlarged. The former organ is congested, and Baccelli believes there is probably catarrh of the bile-ducts in many cases. I have, however, never seen jaundice; and, while the conjunctiva often becomes yellow during the progress of the complaint, the presence of bile in the urine has not been detected. The spleen is invariably somewhat larger than usual, but it seems to yield readily to the action of quinine, as only in one case did it assume great dimensions and project into the epigastrium. It was then very tender to the touch, and pressure on it excited severe retching. But though there may be exceptional cases, the rule seems to be that the spleen does not enlarge to the extent described by writers on intermittent and remittent fevers. In microscopical examinations of the blood, no excess of leucocytes has been found, or any of the pigment-particles which other observers have noticed. In the bowels there is little noteworthy. Constipation seems the rule, necessitating purgatives, and sometimes very strong purgatives. When there has been diarrhoea, it has been of a bilious character; more bile escaping from the congested liver, or the reabsorption in the bowels being more sluggish. No pea-soup stools occur, nor general or localised tenderness on pressure over any part of the abdomen except the epigastrium, or over the liver or spleen if enlarged. Occasionally there is slight meteorismus, but, as Baccelli notices too, it is never persistent. At the beginning, the urine is acid, scanty, of high specific gravity, and filled with urates. As soon as treatment takes effect, it becomes neutral or alkaline, clear, and with only slight deposit of epithelium and of phosphates, but the chief microscopic feature is the nearly constant presence of pigment-particles. Analyses of urea have been made, but are not yet worth reporting. Albumen has scarcely ever been found; a fact in marked contrast to my observations in enteric fever here.

Course.—This fever seems to have no definite duration. The shortest of my cases—a gentleman aged 39—was apyretic on the seventh day, and continued so; but I suspect he had fever two days before I was sent for. The longest case lasted thirty-six days; but this was due to a relapse, as the patient, a young lady, was without fever from the seventeenth to the twenty-second day. The termination of the case is often indicated by the gradual cleaning of the tongue, by the fever assuming a true remittent or intermittent type, and by an increase of the perspiration in the morning hours; and if this be excessive, the temperature has been frequently found subnormal at all hours of the day. Very rarely do anything like critical deposits occur in the urine.

Complications.—Two patients had unilateral pleurisy with considerable effusion, which, however, disappeared quickly. One—a lady aged 51—had pleurisy and endocarditis, and only slowly recovered. One patient had bronchitis, and one thrombosis of the left leg. None of my own patients had any other complication; but one lady, aged 28, whom I attended as consulting physician ten days after the fever began, miscarried at the eighth month, and died from *post partum* hæmorrhage. Baccelli describes complicated cases as special forms of the fever—the bilious, the rheumatic, the pneumonic, etc.—but I know nothing of these.

Prognosis.—None of the forty-five patients treated throughout by myself died; and only three—the two youngest treated and the lady who had the complication with pleurisy and endocarditis—were very dangerously ill. One young gentleman, aged 18, whose temperature was 102.4 deg., insisted on getting up on the tenth day, and went out for a walk. He repeated this on two subsequent days, was none the worse, and was apyretic on the fourteenth day. Some of the patients

had been delicate, and were travelling by the advice of physicians, but none had any well defined chronic malady. In uncomplicated cases, therefore, in average health, the prognosis is very favourable.

Diagnosis.—The only other fever with which this could easily be confounded is a mild enteric fever, as they certainly have some features in common; but the short incubation, and the rapid rise of temperature even to 104 deg. soon after a chill, seem almost absolutely to exclude the idea of sewage-poisoning. The response of the fever to treatment, its indefinite duration, and the total absence of any eruption like that of enteric fever, with the negative character of the bowel-symptoms, seem quite as certainly to disprove the existence of any specific intestinal glandular lesions, even if the *post mortem* examinations made by others were not quite satisfactory on the point, hyperæmia of the Peyerian patches being the utmost that is found; while the frequent termination of the cases as remittents, or even as genuine quotidian and tertian intermittents, seems quite as eloquently to prove its origin from malaria alone. Enteric fever, indeed, is as marked in its features in Rome as in London. I have lost five out of fifteen cases, two from bowel-hæmorrhage; and the abdominal symptoms have been prominent in all. It is obviously essential to distinguish the two fevers, as an early and energetic specific treatment undoubtedly lessens the severity and shortens the duration of the malarial form; whereas, if neglected in the beginning, it becomes rebellious to remedies and very tedious in its course. The prognosis, too, is very different, and it is probably from the two fevers being constantly confounded together that Rome has acquired its unenviable reputation for typhoid; whereas, as I have shown in the BRITISH MEDICAL JOURNAL for September 16th, 1876, fewer cases of enteric fever originate in Rome in proportion to population than in any other large continental city.

The subcontinued type of the fever described, and the rapid variations of the temperature, distinguish it clearly from the malarial remittents of British authors; but it frequently passes into a remittent fever. For other fevers it could not well be mistaken.

Treatment.—When the ordinary antipyretic mixtures have been given in the beginning of those fevers, they have had no appreciable effect, except perhaps in making the pulse a little slower. A good purge is usually given, the much abused calomel very often, and then quinine in some form. At first, it was given in small and divided doses: two grains every two hours, or three to four grains every four hours. But most of the patients have had the drug in one or two large doses in the forenoon, or at least before 2 P.M., if possible, although the thermometer too often fails to show a remission at that time. Unless the patient's objections are insurmountable, the quinine is given in solution. Formerly, it was combined with bromide of potassium, or alkaline mixtures were given at the same time. Now, I almost always use the hydrobromic acid, as recommended by Dr. Milner Fothergill. Both the bromide and hydrobromic acid counteract to a certain extent the unpleasant effects of large doses of quinine; but the hydrobromic acid is much the more effectual of the two. The headache, neuralgia, malaise, and general aching pains either disappear altogether or are much alleviated, as soon as the patient comes under the influence of the quinine. It notably slackens the pulse and lowers the temperature. Fifteen grains will bring the pulse down from five to ten beats, and the temperature from one to three degrees, within two hours of its administration in solution. The average amount given in the beginning is from fifteen to thirty grains daily. I have found no patient intolerant of the remedy, even when their repugnance to it has been based on the opinions of the leading physicians of Great Britain and America. Most of them, after the first doses, are too sensible of the benefit derived to doubt the fact that the disease indicates the means of cure. If the patient will not take it in solution, it is given as a powder in a rice-wafer, followed by some drops of nitric or hydrochloric acid. Quinine enemata have rarely been used, and subcutaneous injections of the drug were necessitated in only one case by constant sickness, and by the young patient not tolerating the enema-syringe. Cold applications, too, are always used. Frequent cold sponging of the body, the ice-bag to the forehead and back of the head, iced compresses over the spleen, are all useful. Tepid baths rapidly reduced in temperature were required in a boy aged 14, who was very delirious. They were given eight or ten times with excellent results. The diet is restricted to fluids; and stimulants are seldom needed, except during convalescence. Salicylate of soda has been given in doses varying from thirty to ninety grains at short intervals; and, in the forenoon hours, if practicable, sixty to ninety grains have nearly the same power in reducing temperature as fifteen or twenty grains of quinine, but the effect seems to disappear more rapidly. Salicin, tincture of eucalyptus, and Warburg's tincture, have also been used; but little reliance is placed on anything but quinine.

Recurrence.—It is uncertain whether this type of malarial fever tends

to recur. In those patients whose history, subsequent to the fever, is known, it has not done so. If it did, it would probably be as some variety of intermittent fever. No injurious sequelæ have come under my notice.

Frequency of Occurrence among Visitors.—It is obviously impossible for any one physician to give an exact idea on this point. The fever goes by so many different names—Roman, bilious, nervous, gastric, typhoid—that it is very difficult to know when it is the type described that is meant. Besides, there are no proper data for calculating the number of visitors in a season.

CLINICAL MEMORANDA.

REMARKS ON SCARLATINOID ERUPTION IN THE AGUISH DISORDERS OF CHILDREN.

IN Dr. Cheadle's interesting paper on the occurrence of ague in young children in London (BRITISH MEDICAL JOURNAL, April 13th, 1878, p. 523), he remarks that he had never previously observed a general scarlatinoïd eruption in young children, nor can discover any instance of its occurrence on record. I have never witnessed so general an eruption as existed in his first case; but I have long been aware that cutaneous hyperæmias occur in aguish disorders, such as I have termed malarïoid remittent, having more or less resemblance to scarlatinal rash; and I referred to them in my work as roseolous. In the case mentioned at page 765, the child, aged 3½, became red all over, legs, face, and all, two or three times a day. She had several pyrexial attacks in the twenty-four hours, and I found her temperature in one of them to be 102.2. In the following case, which occurred to me in 1869, the cutaneous hyperæmia was usually a less striking feature than the anæmia; but both phenomena were well marked, and are both attributable to the same cause, just as the cold and hot stages of a common ague are.

E. B., aged 3, admitted October 14th with herpes of the face, was described as having suffered for one or two years with seizures occurring two or three times a day, in which she turned "deadly white"—"as white as a child laid out for dead". She was also very restless at night, very feverish, and "burned" very much. Usually the burning affected the hands; but on November 24th her chest and face were quite red, as in scarlatina, for three hours; and about the middle of December she was one day quite prostrate and scarlet all over, slept all day until the evening, then woke up, lost her redness, and appeared quite well. Beside the vasa spasm and paresis, she had fits of panting and difficulty of breathing, lasting several hours or even a whole day. These were probably analogous attacks of bronchial spasm. In general condition, she was languid and drooping, and her flesh was flabby. With citrate of iron and quinine, five grains three times a day, administered for three months, she improved steadily, and completely recovered. In such a case as this, the diagnosis of malarïoid remittent was sufficiently plain; and usually the rash is, I think, more suggestive of roseola than scarlet fever. Yet, in some cases, for a time doubt may prevail, especially should there happen to be some sore-throat.

I have no note as to whether this child had ever resided in a malarious district; but that London is capable of giving rise to aguish disorder is, I hold, certain. In addition to the evidence given in my work at page 761, I may mention the case of a student who had never had ague, or been in the way of it, before he came to London, where he had several attacks, and could not get rid of it until he left. Another case was that of a young lady who always lived in a high dry locality, except that she spent one night at Stratford near London. She had well-marked ague for three or four weeks, and felt its effects long afterwards. Jacoud's experience of Paris is quite similar.

C. HANDFIELD JONES, M.B.

STRAPPING THE CHEST IN THE TREATMENT OF CHRONIC PULMONARY CONSUMPTION.

IN private practice, I have lately had a case of pulmonary consumption in which great benefit was produced by strapping the upper part of the chest. The patient was a female aged 33, the wife of a tradesman; she had had one child. The disease apparently began, as chronic pneumonia of the apex, in February 1877; and when I was first consulted, exactly a year afterwards, there was sufficient evidence of a cavity at the top of each lung, and the constitutional symptoms were very grave. The case was clearly hopeless; but at first there was a marked improvement from the internal use of the hypophosphite of lime, and from the external application of strips of soap-plaster, ex-