pneumonia surrounding small and large racemose masses of aggregated tubercle. These masses are generally more or less yellow in colour from the process of caseation, and present in some parts excavations of evidently recent date. Scattered miliary tubercle is generally present, though the aggregated form is more commonly met with. In this variety, we find occasionally ulceration of the intestines and the presence of tubercle in the solitary glands; but the serous membranes are free.

The clinical symptoms may be described as those of chronic tubercular phthisis in an exaggerated form. The cough is very troublesome, and is early accompanied by large expectoration; anorexia is generally present; the tongue varies, but is sometimes red and irritable in appearance, and, as may be seen in the following case, diarrhoea often occurs. The pulse is rapid, varying from 100 to 130, and the respirations are hurried and count between thirty and forty.

The temperature is distinctly pyrexial, with occasional but rare periods of collapse, the high figures being maintained more uniformly towards the end than is the case in scrofulous pneumonia, the reason probably being that excavation is accompanied by fresh tuberculation even during the last weeks of the patient's life.

Let us take an instance of this form.

Maurice A., aged 22, married, was admitted into the South Branch of the Brompton Hospital on April 7th, 1874.

History.—He had lost several maternal aunts, of phthisis. He had suffered from general weakness and prostration for eight months, and for the last four had had cough, with expectoration, accompanied by loss of flesh and night-sweats. He complained of pains in the back and indigestion after meals. His aspect was very anæmic; cough troublesome; expectoration mucopurulent; tongue furred; appetite moderate; weight 9 st. 7 lbs.

Physical Signs.—Dulness, flattening, and cavernous sounds were detected on the right side as low as the fourth rib.

April 30th. He had gradually become worse; the temperature and pulse had risen and the appetite had fallen off. He had lost three pounds in weight. In addition to the above signs, crepitation was audible under the left clavicle, and tubular sounds in the second intercostal space. He was transferred to the main hospital and confined to bed.

May 8th. The cough was worse; the temperature, which was taken five times a day, was 100 deg. Fahr., except at 8 A.M., when it fell to 98.4 deg. Fahr.; the respirations were 30; pulse 104. The crepitation-sounds had increased in extent on the left side, and cavernous sounds were audible below the fourth rib, on the right side.

May 24th. He had wasted considerably. Slight diarrhoea came on yesterday, there being two or three loose motions in the twenty-four hours. The temperature was somewhat higher, being 101.2 deg. Fahr. at 8 P.M. Respirations 102; pulse 120.

June 4th. The bowels were still relaxed, motions passing twice a day. There was oedema of the left foot; no albumen in the urine.

June 17th. Weight 8 st. 4½ lbs., showing a loss of 1 st. 4½ lbs. since April 13th. He complained of much pain under the right nipple; the skin was perspiring. *Physical Signs:* On the right side, dulness was present over the upper half; dry cavernous sounds to the fourth rib; croaking sounds below. On the left side, tubular sounds were audible over the upper fourth in front; marked dulness over the upper two thirds.

June 22nd. The breath had become very difficult; aspect pale and anxious; tongue red, furred in the centre. Pulse 120; respirations 48. Crepitation was audible over the whole right side. The left leg was swollen and oedematous. There was a lump in the popliteal space, tender on pressure, due probably to plugging of the vein.

This patient died on the 26th. Below are the averages of the day temperatures from 8 A.M. to 12 P.M., taken for a period of about three weeks.

SUMMARY.	8 A.M.	11 A.M.	2 P.M.	5 P.M.	8 P.M.	10 P.M.	12 P.M.
Averages.....	99.1	99.7	100.4	100.3	100.9	101.1	101.4
No. of observations.	9	22	22	23	24	15	9

Observations were hourly taken during the night on two occasions, and the result was, that the temperature was maintained at 100 deg. Fahr. until about 7 A.M., when a distinct fall commenced, reaching as low on one occasion as 96 deg. Fahr. The greater part of the observations were, however, high.

Autopsy, twenty-four hours after Death.—Both lungs were somewhat collapsed. The right lung weighed 2 lbs. 11 oz., and was firmly adherent; the upper lobe was completely excavated; the lower and middle lobes were consolidated, with aggregations of miliary tubercle in which centres of softening were visible; the middle lobe contained a cavity of recent formation. The left lung weighed 2 lbs. 5 oz.; the pleura was adherent; the upper third was consolidated, with aggrega-

tions of miliary tubercle caseating in parts, with some small cavities, one the size of a walnut. The lower lobe contained a few grey aggregations, the rest being in a state of red hepatisation. The liver, nutmeggy, weighed 4 lbs. 7 oz. Intestines: The solitary glands of the ileum were caseous; some had broken down and formed ulcers; the cæcum was a mass of ulceration.

This case only lasted altogether six months and a half, and it may be safely affirmed that the principal changes in the lungs and intestines took place during the two months and a half the patient was under observation in the hospital. The tuberculosis of the left lung, the pneumonia of the lower lobe, the fresh miliary tubercle in the right, and the rapid extension of the cavity, were all noted during the last few weeks of life, while the regular thermometric record enabled us to compare this phenomenon with the lung-changes. The high temperature so continuously maintained was probably due to the various and successive fresh crops of tubercle being formed, and the diarrhoea to the ulceration of the cæcum, though, considering the extent of surface involved, it is remarkable that the flux was not more profuse. The prognosis in most of these patients is gloomy, as the lung-disorganisation proceeds generally to utter destruction; but, in some instances, the fever diminishes and the tubercular masses remain quiescent, during which time the patient regains strength and flesh, and may live on in a precarious state.

TUBERCULAR MENINGITIS.

By REGINALD SOUTHEY, M.D. Oxon., F.R.C.P.,
Physician to St. Bartholomew's Hospital.

THE subject of tubercular meningitis has become invested with some special interest at the present moment by the recent Penge case, and, without presuming to offer any opinion upon the facts, far too few in themselves, observed by the medical witnesses at that trial, it is possible that a further knowledge of the clinical symptoms attached to some undoubted cases of tubercular meningeal disease in adults might engage the attention of medical practitioners.

In a previous clinical lecture upon the same subject published in the JOURNAL of January 22nd, 1876, page 97, I directed notice to two hundred and sixty-seven well authenticated cases of this affection placed in a table, from which it became easy to infer the relative frequency of the disease at different periods of life. Of those cases, 25.8 per cent. occurred in children under five years of age, 17.2 per cent. befell children between six years and ten years of age; between the ages of eleven and fifteen years, the disease became exceedingly rare, 7 cases only out of 267, or 2.6 per cent., only presenting themselves; whereas, between sixteen and twenty years of age 31 cases were recorded, and, again, a similar number between the ages of twenty-one and twenty-five, or at both periods of life respectively, 11.6 per cent. Furthermore, as life advances, not only are the symptoms attached to the disease far less distinct than when it occurs in childhood, but the malady begins and pursues its fatal course with singular insidiousness, and is frequently misunderstood even by well experienced medical men until the autopsy reveals its true nature.

The following cases which I have collected together establish my statement.

CASE I.—In June, 1867, a young woman between nineteen and twenty, comely, well nourished, fair-skinned, but having rather coarse black hair, was admitted by one of my colleagues into Faith Ward. She complained of various aches and pains. Headache was her chief trouble, and she appeared generally nervous and excitable. She had walked to the hospital from some little distance; was fatigued, answered questions precisely and rationally, but was brusque to rudeness in her manner. There was no account of vomiting; no constipation, no irregularity of pulse; but her temperature was 102 deg. Fahr., and this important fact determined her admission. She walked upstairs without assistance; became drowsy directly after getting into bed; but was easily roused up by the nurse to take her nourishment, which she swallowed without difficulty. Six hours later, she was found in a state of profound coma, out of which she could not be stirred, and, thirty hours after her admission, she died.

POST-MORTEM EXAMINATION.—The body was more than usually plump. There was a large amount of fat about the heart. Extensive pericarditis was found on both surfaces, presenting layers of thick butter-like lymph. Several whitish fibrinous bodies, like warts or papillary growths, stood out from the muscular substance of the heart, growing apparently from interfibrillar hyperplastic connective tissue roots that were deeply interpolated between the muscular bundles of the ventricular walls. Microscopical examination of them showed

identity of structure with ordinary tubercle-growths; they consisted of small cells and fine granules closely packed together. The convolutions of the brain were slightly flattened; some excess of clear serum existed in the ventricles. Granulations were found upon several vessels of the choroid plexus, with some lymph attached here and there to their surfaces. There was lymph also in the subarachnoid spaces upon the upper surface of the brain. The base of the brain was the seat of extensive fibrinous lymph-exudation, especially in the cleft of both Sylvian fissures.

CASE II.—William Fuller, aged 45, a fire engine-driver, was admitted under my care in Radcliffe Ward in 1867. He was a spare, sallow-complexioned, black-haired man, with flushed face and injected ferret conjunctivæ. He had been ill for a few days only, complaining of pain in his head and being feverish, wandering in his talk at night, and being excessively prostrate. Three nights previously to his admission, he had driven his engine to a fire, and had been out all night and drenched to the skin. At my examination of him, he was clear and precise in his account of his illness, which he attributed to a feverish cold; but his manner was nervous and excited; his answers to all my questions curt and ill-tempered. His skin was hot (102 deg. F.); tongue furred and dry; pulse quick, 108, but regular. There were some mucous râles in both lungs, but no evidence of pneumonia. The heart-sounds were clear; the abdomen was not full or tender; the urine was scanty, containing no albumen. He had no appetite, and was not very thirsty; he vomited what he drank. Typhus was at that time epidemic in London, and his case was suspected to be fever; but no rash was observed, and no diagnosis was made.

On the same night, September 27th, the patient became violently delirious, and was with great difficulty restrained in bed. On the 28th, he was seen by Mr. Wood, our resident apothecary at that date, who ordered a large blister to the nape of his neck. He was believed to have complained of pain at the back of the head and of stiff neck. He became comatose, and died the following morning. The *post mortem* examination discovered general tubercular meningitis, with tubercle in the lungs and other viscera. The only notes I possess of the case are very imperfect, but not wholly without value.

CASE III.—James Reed, 29, cook on board a steamboat, was admitted under my care in Luke Ward on November 12th, 1868. He was a thin, small-made, sallow-complexioned man, with lank brown hair and a razor-ridged nose. *History.*—The patient had been complaining for eight weeks, with loss of appetite, constant inclination to vomit, frequent sickness, and supraorbital headache. He had emaciated much, but still had been able to work a little up to three weeks before his admission.

Condition on Admission.—He was thin and feverish; the tongue was furred in patches and inclined to be dry; he had a foul taste in the mouth; the bowels were constipated, the abdomen retracted, the skin dry; urine high-coloured, scanty, specific gravity 1032, non-albuminous. The right testicle was swollen, hard as a stone, but not tender, and entirely painless. He had not noticed when it first became enlarged, but thought it had been so for some weeks. The heart-sounds were feeble, but normal; the respirations hurried; no moist sounds could be heard over the lungs; there was increased resonance over the left thorax. The nights were sleepless; intelligence was perfect; temperature 100.5 deg. (?) The real character of his nights was ascertained after he had been in a few days: he was restless the early part of the night, then slept heavily for an hour or two, and woke up complaining of his head; he frequently remarked upon his headache being aggravated by sleeping and by lying on his left side. *Treatment.*—He was ordered milk, lime-water, and ice, and a simple enema to open his bowels.

The testicle was carefully examined by me, and formed the subject of some clinical remarks; it felt hot and hard; the epididymis chiefly was the seat of enlargement; examined by candle-light, it was seen to be no hydrocele; it was not simple inflammatory orchitis. He was aware of no blow or injury that had provoked it; there had existed no previous urethral discharge. In size it was scarcely more than half as large again as the left or sound testicle, and the enlargement was not general, or the induration of that stony hardness which usually betokens scirrhus. Further, the inguinal glands were enlarged solely upon the affected side, and were neither so hard as syphilitic, nor so large as cancerous glands usually are. Against the hypothesis of cancer was the absence of pain and tenderness. Against syphilitic gummata was the feel of the testicle and the situation of the enlargement. Syphilitic growths commence usually in the interstitial substance in the body of the testis, between the seminal tubules; they spread to the periphery and protrude the tunica albuginea at some one prominent spot. Tubercular disease begins usually as an epididymitis that spreads from the wall of the vas deferens inwards; at an early stage, the epididymis

feels hard, swollen, and clay-like, but it is some time before the body of the testis is implicated. Exceptionally, I am aware that tuberculous disease may originate in the interlobular structures of the body of the testis; it would then be difficult to distinguish from syphilitic disease. The history, the slow progress, the absence of pain, the local heat, the small tenderness, the situation of the principal enlargement, all favoured the diagnosis of tubercular disease of the testis; and the general symptoms—the fever, the headache, the vomiting—indicated some secondary deposit of tubercle in the substance of the brain or its membranes.

Course and Progress.—Three days after admission, a thick, yellow, slightly blood-stained discharge came away from his right ear; there was no pain or tenderness, or sign of inflammation in the external meatus; the headache appeared temporarily relieved by this discharge. The discharge, which amounted to from half a drachm to two drachms in the twenty-four hours, continued; and it was observed that he was freer from pain and vomiting when it was more abundant. A week after his admission, the testicle began to give him pain; indeed, as the pain in the head subsided, the pain in the testicle began and increased. The intellect remained clear; the appetite continued indifferent; the temperature usually between 100.5 deg. and 102.5 deg.; pulse regular, 96, small; bowels costive. He felt giddy upon attempting to rise up suddenly; but this passed off, and he was able to get out of bed without help. He was usually sick, once daily, early after his breakfast. Subcutaneous injections of morphia were administered on most nights.

The following notes were made on December 14th:—His appetite improved; he had eaten a mutton chop; the tongue was cleaner; he slept better; he had not vomited for two days; the testicle remained in the same state; there was a continual slight purulent discharge from the left ear; pulse 108.

December 17th. He was feebler and more drowsy all day; but could be roused, and then answered questions rationally; he recognised and spoke to his wife, and got out of bed by himself.

December 22nd. He was completely insensible since early in the morning; pulse 84, irregular; the mouth was slightly drawn to the left side. He was believed by the nurse to have had a fit, though there was no distinct evidence of his having been convulsed. The sphincters were relaxed; no reflex movements were excitable.

December 23rd. He remained in an unconscious state; but swallowed mouthfuls of fluid well and easily. Reflex movements of both arms and legs were excitable. The abdomen became tympanitic, and he gradually sank and died, without having recovered consciousness, on December 25th, the morning of the third day from that on which he had lost consciousness.

The *post mortem* examination showed general tubercular meningitis, an excess of turbid serum in the ventricles of the brain, which escaped upon its being opened; adherence of the Sylvian fissures; tubercular lymph at the base of the brain; the brain-substance very soft, in parts almost diffident. The cerebellum was firm, but containing three distinct globular tubercular masses of greenish yellow colour, each about a quarter of an inch in diameter. The left lung was the seat of extensive miliary tubercle scattered generally throughout its substance; the pleura was universally adherent; the epididymis of the right testicle was much enlarged and breaking down throughout into caseous pus; some portions of the body of the testis were becoming involved in the degeneration.

REMARKS.—Although very imperfectly recorded in my own notebooks, I attach much value to this case, illustrating as it does the insidiousness of the cerebral symptoms in a very remarkable manner. Looked at pathologically, the cerebellar masses and the tubercular disease of the epididymis might have been of about the same date. The lung-mischief and the meningeal must both, I think, have begun after his admission into the hospital. But here was a man able to stand and help himself, although feeble, rational, with no paralysis of motion or sensation, possessed of faculties of sight and hearing (the latter having been only partially affected upon the left side) to within three days of his death.

CASE IV.—Maria Mason, a general servant, who had been hard worked, was admitted on September 18th, 1875. She was a careworn woman, with an antero-posterior spinal curvature; there was some thickening in the lumbar region, prominence of spines of the lowest dorsal and first and second lumbar vertebrae, and tenderness upon deep pressure over them. Her face was flushed and her manner excited. She had been supported, but not carried, upstairs. Her temperature was high. There were abundant flea-bites over her trunk, and some few raised spots, that were supposed by some who had seen them to be pathognomonic of typhoid fever. Her tongue was dry and furred, and there were sores both upon her lips and teeth. It is only fair to state that she was admitted as a case of typhoid fever. The account given

by her friends was that she was strange in her head, had been slightly delirious at night for some two or three nights, and had talked nonsense by day as well; that she took very little food and vomited frequently. Her own account of herself was given me with much precision: she remembered dates and facts accurately; and, as I found afterwards, her history of her past and present illness corresponded with the accounts given by those with whom she had lived.

She had been laid up in her situation for more than a week, but had had no diarrhoea: her chief complaints had been pain in her head and back, and almost constant sickness after every kind of food; her catamenia had been irregular for some months. She was aware that her back was rather bent, and that she stooped in consequence of it, but was sure that this had only taken place in the course of the last eighteen months. She had been housemaid to a family where a good deal of water had to be carried upstairs for baths; and a year and a half previously she had fallen upstairs while carrying two heavy water-cans, and then strained her back severely. She was able to continue at her work that same day, but was shortly afterwards laid up with pain and tenderness in her back; and then left her place, returning home to her mother's house, where she was laid up in bed for three months. During those three months, she was paraplegic; and her account of this paraplegia was as precise as could be; thus, first she had hyperæsthesia in the right leg, then loss of power of motion in the same leg; and, within a few weeks, the same sequence of events occurred in the opposite limb, although the ultimate loss of motor power in this left leg never was so complete as it had been in the right. For three months, she was quite unable to stand or lift herself, and at night-time her legs used to wake her up by their involuntary startings. Her sphincter ani never was affected, but she was occasionally unable to retain her urine.

After complete rest in bed, without doctor or physic, but with the most careful nursing upon the part of her mother, she gradually recovered power, first to move herself, and then to stand, and finally to walk. She obtained another situation, and had held this for twelve months, working very hard up to the commencement of her present illness. Curiously enough, as showing family predisposition, she had a sister suffering with chronic spinal disease.

Examination of the Patient.—She lay chiefly upon her back, I was told; but was rather curled up upon her right side when I first saw her, and nearly always afterwards assumed this posture by preference. She moved very little, but had complete capacity of motion, and perfect sensation in both arms and legs. There was slight ptosis of the upper eyelid of the right eye: her eyesight, she said, was bad; but she saw and correctly described objects upon the opposite wall of the ward. Both pupils were of equal size, and reacted naturally to light; the conjunctivæ were red and injected. There were occasional twitchings of the facial muscles. There was no fullness or tenderness of the abdomen, which was rather empty; no increased dulness over either liver or spleen. She had slight occasional cough, but no abnormal chest-sounds. The heart's action was regular, its sounds clear. She complained to me of a feeling of constriction across her chest, and of dull heavy pain in her occiput and down the back of her neck. Pulse 104, regular; temperature 101.6; respirations 23. Vomiting, headache, nape pain; previous (strumous?) vertebral disease; the nature of the fever; a morning temperature of 100.8, an evening of 101.6; no symptom of typhoid fever, except continued fever with night delirium; no evidence of pneumonia—these things, taken all together, guided me to the opinion I gave that the case was probably one of tubercular meningitis, the tuberculosis being secondary to some caseous pus collection in connection with her former spinal disease. The treatment consisted of milk, essence of beef, and one drachm of bromide of potassium every six hours.

Course and Progress.—September 19th. She had passed a quiet night; had had no vomiting. Fluid nourishment had been taken at frequent intervals of time in small quantities, and was well supported. Urine occasionally passed in her bed; but she appeared at once aware of this, and called the nurse's attention to it. There had been one action of the bowels, loose, dark coloured. The lips were dry and peeling; tongue red and dry. Morning, pulse 96, regular; temperature 101.4; respirations 20. Evening, pulse 105, regular; temperature 102.4; respirations 36.

September 20th. She had had a restless night; her mental condition was unaltered. She answered questions correctly. The skin was dry and harsh-feeling. The bowels acted once daily; the motion was dark and loose. Morning, pulse 110; temperature 101; respirations 40. Evening, pulse 96; temperature 101.4; respirations 32.

September 21st. Her posture was unchanged; she was quite constantly lying upon her right side, and curled up, resting her head upon her right hand, and complaining of pain at the back of her neck when her head was moved. She answered questions deliberately, though

not always correctly, forgetting recent things. When asked to show her tongue, she did so. There was more marked ptosis of the right eyelid, with increased secretion from the eyes. The pupils were equal, and equally sensitive. She swallowed what food was given her easily and well, but would not attempt to feed herself. The tongue was dry; the bowels acted once *per diem*; urine passed chiefly unconsciously under her. Morning, pulse 100; temperature 101.2; respirations 24. Evening, pulse 99; temperature 102; respirations 40.

September 22nd. Simple semiautomatic acts, like putting out the tongue and taking nourishment, were fairly well performed, but no more complex intellectual acts. The patient lay in a drowsy, half-dreaming state, in which she occasionally muttered. She groaned when disturbed. The skin appeared hypersensitive to impressions of cold or touch: the slightest touch of the soles of the feet was responded to by the drawing up of her legs. The eyelids were much gummed-together, but she was still able to open the left; she did not appear to see with either eye. There was no reflex movement when her eyes were rapidly approached with a finger. She was restless, tossing about the limbs, throwing off the bedclothes, and clutching and picking at the coverlet. No action of the bowels had taken place. The urine passed involuntarily. Morning, pulse 100; temperature 101.6; respirations 60. Evening, pulse 105; temperature 100.6; respirations 30.

September 23rd. She was not quite so drowsy nor so restless. The tongue was cleaner and less dry. Nourishment was taken more readily. She had no sickness, and opened her eyes. The ptosis of the right lid was scarcely noticeable; vision doubtful; abdomen more sunken; any movement of the head was followed by evident signs of pain. Morning, pulse 104; temperature 101.4; respirations 30. Evening, pulse 100; temperature 99; respirations 36.

September 24th. She remained in the same semicomatose state. The pupils were dilated, not reacting to light. The eyelids were half open. Occasional movements were made by all her limbs. She was setting aside imaginary objects with her hands; but these motions were tremulously performed and badly coordinated. The muscles and tendons of the hands were twitching much when I held her hand to count her pulse. She mumbled indistinctly to herself when roused, and called out sharply when her pillow was moved. Pulse 120; temperature 98.8; respirations 30.

September 25th. The coma became more profound; swallowing impossible; the breathing quicker, and at times catching and irregular; the pulse irregular, and scarcely perceptible at the wrist. A slight convulsion shortly preceded death.

At the *post mortem* examination, twenty hours after death, old caries of the second lumbar vertebra was discovered, with a collection of caseous pus upon its anterior aspect. The spinal cord had evidently been compressed in some measure by this old abscess, and the sheath of the cord still bore distinct marks of pressure-thickening in this situation. The cord itself was here very soft, and easily broken down; but there was no evidence of recent disease visible either in the cord or theca. Scattered miliary tubercle was found in both lungs, and in the sheath of the bronchi, but no extensive deposits; also in the kidneys. There was abundant lymph at the base of the brain, with matting together of the nerves; some excess of fluid in the ventricles, but no flattening of the cerebral convolutions, and therefore little compression.

There remain two more cases to narrate and some remarks to append, which must stand over for a second communication.

A CASE OF STARVATION.

By THOMAS ELLIOTT, M.D. Dub.

THE difference and diversity of medical opinion exhibited in the Penge case induce me to record the following case, which I trust may interest such of the readers of the BRITISH MEDICAL JOURNAL as notice medico-legal questions. It was not alleged, in this case, that the deceased had not received food; but it was clearly proved at the coroner's inquest that it had been insufficient. For five or six months, though she complained at times to her husband of not having enough to eat, she remained in this sad plight; and when at length relief from the parish official came, it proved to be of little avail. It not being a case of acute starvation, the *post mortem* examination did not reveal anything particular beyond the extreme emaciation.

Ann G., aged 57, a married woman, was admitted from South Noranton into the Mansfield Union about 6 P.M. on Thursday, September 27th, 1877. Being extremely deaf, it was impossible to elicit any history of the case from her. As gleaned from the coroner's inquest, it appeared that, her husband being short of work, they had lived in a destitute condition for five or six months, and she had been without