grains; and, as no dangerous effects have attended the somewhat free experiments hitherto made with it in the healthy subject, it may be given with confidence. I would propose that an infusion of one drachm of the powder, in an ounce of water, be injected at a blood-heat into the rectum, and repeated every two hours until the full influence of the drug is established. If there be intolerance of the simple infusion, it may be medicated by the addition of liquor opii and starch, so as to facilitate its retention. If the active principle of the drug could be obtained, it would, of course, be much more easily managed in the form of a concentrated solution for hypodermic injection. The liquid jaborandi of Messrs. Ferris and Co. I have found to be a reliable test. The dose is from five to ten drops. It is therefore presented in a handy form for injection either *per rectum* or hypodermically.

ON THE TREATMENT OF HIP-JOINT DISEASE BY EXTENSION WITH MOTION,

AS PRACTISED BY THE AMERICAN SURGEONS, INSTEAD OF LONG-CONTINUED REST AND IMMOBILITY.*

By WILLIAM ADAMS, F.R.C.S.,

Surgeon to the Great Northern Hospital, and to the National Hospital for the Paralysed and Epileptic; Consulting Surgeon to the National Orthopædic Hospital; etc.

THE object of the present paper is to direct attention to the recent advances which have been made in the treatment of hip-joint disease by the American surgeons, to whom we are indebted for the discovery of two very important principles, and also for their practical application by means of most ingeniously contrived instruments.

The first principle is that of extension, as a means of relieving the most acute pain in joint-diseases, especially applicable to the knee and hip-joints.

The second principle is that of extension combined with motion during the progress of disease, the patient being allowed to walk about, so as to promote recovery with free motion in the joint, instead of the ordinary result of ankylosis obtained by long-continued rest and immobility.

There can be no doubt that the discovery and practical application of these two principles have completely revolutionised the treatment of joint-diseases, and changed our opinion with regard to the pathological conditions existing, especially as to the production of acute pain which, formerly, was believed to depend upon acute inflammation, requiring active local, as well as general antiphlogistic treatment, such as leeches, blisters, calomel and opium, etc. It has now been proved to depend upon undue articular pressure and contact of inflamed surfaces, produced by reflex muscular contraction, and capable of relief by mechanical means alone, producing extension, whether this be applied by means of the weight and pulley, or by the screw and cog-wheel.

The object of extension is not, as generally supposed, to separate articular surfaces, but to overcome reflex muscular contraction, and, by relaxing the muscular rigidity, to prevent undue pressure of inflamed articular surfaces, or their margins, when the joint is held in a flexed position by muscular contraction.

Extension, for the relief of pain, is applied by means of the weight-and-pulley apparatus whilst the patient is in bed, the extension always being made in the same direction as the long axis of the limb. It will vary, therefore, when first applied, as the leg may be flexed, adducted, or abducted; but as soon as the reflex muscular action has been overcome, the weight will be made to pull in the direct horizontal line. After the pain has subsided, the object with which the weight-extension is continued is to secure sufficient rest to the joint, and at the same time permit a limited amount of motion, as the patient varies his position in bed. It is not found, by experience, that the limited amount of motion thus permitted in any way interferes with, or retards, the favourable progress of the case; but, on the contrary, this may be looked forward to with confidence in the great majority of cases after the pain has been relieved by the weight extension. The health of the patient also improves rapidly after the relief of pain.

The successful application of weight-extension as a remedy for pain, however acute it may be, in joint-disease, is the greatest discovery of modern times in the treatment of these affections. The originality of this discovery is claimed by Dr. Henry G. Davis of New York. In his work on *Conservative Surgery*, published in the year 1867, at

page 206, Dr. Davis observes: "We consider it fully established that, when disease about a joint renders the movement of that joint painful, the joint is always liable to be destroyed by uninterrupted pressure effected through the contraction of the muscles passing over it." And again at page 212, "In diseases of joints, we were the first to point out, as an always present factor in their destruction, the existence of unremitting pressure, as effected by contraction of muscles passing over the joints, causing constant forcible apposition of the surfaces within the joint. This is a general principle appertaining not only to the hip, but to all joints similarly affected. The application of this universal principle should guide us in any joint-affection, whether the disease be internal or external, as soon as the disease renders the movements of the joint painful. When this fact was fully established in our mind, we were led to seek the best way of counteracting the contraction of the muscles, and soon came to the conclusion that a constantly acting force, however moderate, must eventually weary muscles by giving them no respite. We adopted the weight as applied by means of the cord, pulley, and adhesive strips for this purpose, when the patient was confined to his bed or couch. This answered the indication perfectly, relieving all pain and constitutional disturbance, enabling the patient to enjoy his days free from pain, and to rest quietly at night, to relish his food, and to be nourished by it; in short, it robbed the disease of all its terrors".

Dr. Davis does not precisely state the period at which he arrived at the above conclusion as to the cause of the acute pain and its method of treatment; but it must have been anterior to the year 1855, which is the date affixed to his extension splint by the instrument-makers Otto and Reynders of New York. At page 308 he states: "In 1856, I had fully established the treatment of morbus coxarius upon the plan of overcoming the pressure effected by the muscles, through the means of *continued elastic extension*, and the mode with its results was very flatteringly commented upon by one of the editors of the *American Medical Monthly* of this city, in March 1857."

There can be no doubt that interarticular pressure, or the pressure of inflamed articular surfaces—one surface being generally pressed against the margin of the opposite articular surface—must take place when the joint is drawn into a flexed position by the rigid contraction of the muscles, more or less of a spasmodic character, which always coexists with the period of acute pain. Mr. Hilton has particularly pointed out this fact in his *Lectures on Rest and Pain*, and has explained it to be the result of reflex nervous irritation, depending upon the anatomical fact of the various muscles surrounding the joint being supplied by branches of the same nerves, which are also distributed to the interior of the joint. The explanation of this period of acute pain in joint-disease, and its dependence upon interarticular pressure, produced by the abnormal contraction of the muscles surrounding the articulation, has been brought before the profession in a very able paper by Mr. Howard Marsh, in the *St. Bartholomew's Hospital Reports*.*

The English idea has always been rest and immobility to the joint. The American idea, during the last ten years, has been extension with motion, *i.e.*, preserving motion in the joint whilst the pain is relieved by extension.

In the treatment according to the English system, immobility of the joint is obtained by various instruments and splints; from that piece of surgical antiquity, the *long straight splint*, reaching from the axilla to the foot, necessitating the confinement of the patient in the horizontal position for many months, and many other contrivances, such as metal and leather splints to the joint, which permit the patient to move about on crutches, to the now somewhat fashionable Thomas's splint, invented by Mr. Thomas of Liverpool, and described in his recently published work.

All these means succeed, more or less, in relieving pain and promoting recovery, although ankylosis is frequently produced, and this has generally been regarded as the most desirable termination; but in many cases they all fail in relieving pain, for want of the American extension principle, and also they do not prevent the occurrence of dislocation, or partial dislocation, the effect of which is to produce shortening of the limb with permanent lameness.

An example of this occurred recently in my own practice, dislocation, or partial dislocation, taking place whilst the patient was wearing a Thomas's splint. A young gentleman, aged 6, was brought to me in March of the present year, suffering from slight symptoms of hip-joint disease; no pain, but limping, with muscular rigidity, and and slight flexion of the hip-joint. I ordered Thomas's splint, which

* Read in the Section of Surgery at the Annual Meeting of the British Medical Association in Manchester, August 1877.

* On the Prejudicial Effect of Interarticular Pressure in Joint-Disease; and the Application of Continuous Extension by means of a Weight, as a Remedy for this Condition. By F. H. Marsh.—*St. Bartholomew's Hospital Reports*, vol. ii. London: Longmans, Green, and Co.

was made by Mr. Krohne, and applied by myself, with his assistance. I also ordered an extension-apparatus to be used at the same time with Thomas's splint, as the absence of extension appeared to me to be its chief defect. The splint, however, appeared to answer its purpose so well, and the child could be so readily moved about and taken into the open air, and at the same time apparently improving, that the extension-apparatus was not used by the parents, and I saw no more of the patient until sent for in consequence of the occurrence of pain, chiefly at night. This took place nine weeks after the application of Thomas's splint, and it continued two or three weeks before I was sent for; and I then discovered that a dislocation, or partial dislocation, of the head of the femur had taken place; shortening of the limb was apparent, the top of the great trochanter being nearly on a level with the anterior superior spinous process of the ilium. After this the pain ceased, and the disease will probably be arrested; but there must remain permanent shortening, with lameness, and probably complete loss of motion. I am, however, now trying persistent extension night and day, in the hope of diminishing these evils, which could not have occurred if the American system of extension had been relied upon, or extension combined with Thomas's splint, which I ordered. On July 17th, Professor Sayre went with me to see this patient, as he was desirous of seeing cases in which Thomas's splint had been applied.

In my own practice, during the last ten years, since the publication of Dr. Davis's book, I have, in the treatment of hip-joint disease, relied chiefly upon weight-extension, at the commencement, continued day and night with the patient in bed, especially if pain exist; but, as improvement takes place, I allow the patient to walk about with the assistance of crutches, wearing at the same time a firm leather splint moulded to the hip, reaching from the waist to the knee. This secures rest and immobility to the joint, whilst general exercise of the body is permitted; but during the daytime, whilst the patient is not walking about, the leather splint is removed, and the weight-extension applied whilst the patient is on the sofa. Immobility, therefore, is not long continued; and in all cases where the disease has been arrested in the early stage, after treatment lasting from one to two years, motion at the hip-joint has always been preserved, and no lameness whatever remains. Patients treated in this way some years ago are now able to walk and take horse exercise without any fear of the recurrence of pain or any inconvenience.

When disease has not been arrested in the first stage, but has advanced to abscess, with destruction of the joint, motion is generally lost, and ankylosis results; but in some cases, in which I have relied upon extension throughout the disease, an useful though limited amount of motion has resulted.

Before concluding this paper, I will allude to the second principle in the treatment of hip-joint disease, for which we are indebted to the American surgeons—viz., that of extension combined with motion, allowing the patient to walk about during the progress of disease, so as to promote recovery with free motion in the joint, instead of the ordinary result of ankylosis obtained by long continued rest and immobility.

The most recent American advance has been to combine extension with the preservation of motion, and at the same time to allow the patient to walk about without crutches, the weight of the body being transmitted to the ground by a steel instrument applied to the limb from the waist to the foot, or rather a little below the foot. Attached to the steel band round the waist, are two firm inelastic perineal bands, which, passing from before backwards below the tuberosity of the ischium on each side, receive the weight of the body. A raised boot of one inch in thickness is worn on the foot of the sound limb. The limb is connected with the foot-piece of the instrument by means of two long strips of plaster, extending from a little above the ankle to the middle of the thigh, and retained in position by a circular bandage. Attached to the lower extremity of each strip of plaster is a buckle, to which is fastened a strap coming upwards from the foot-piece of the instrument. When thus connected, extension is made by means of a rack and pinion in the longitudinal bar of the instrument at about the middle of the calf. Inversion and eversion of the foot can also be controlled by a circular cog-wheel placed a little below the hip-joint.

The instruments now generally in use in America for the application of extension with motion, whilst the patient is allowed to walk without crutches, are those invented by Dr. Taylor and Professor Sayre of New York, also one by Dr. J. C. Hutchinson of Brooklyn, all of which are modifications of the instrument first invented for this purpose in the year 1855 by Dr. Henry Davis of New York.

My own experience in the use of these instruments is very limited; but, during my visit to America last year, I had the opportunity of seeing them applied in a large number of cases, and as it appeared to me

with great advantage. In one case, that of a young lady, who was residing in Dr. Taylor's private establishment in New York, where patients are received for the treatment of various deformities, the hip-joint disease appeared to be in a more active stage, judging from the pain she suffered, than I should have thought the walking instrument could have been applicable; still, when the extending force was applied pretty nearly to its full extent by Dr. Taylor, she was enabled to walk without pain, and therefore it seemed to be a test-case of the value of extension. Children with hip-joint disease in a more chronic form are frequently seen walking about the streets of New York wearing these supports, and are enabled to get in and out the tramway-cars without difficulty.

Dr. Sayre applied his walking instruments (represented in fig. 2), made by Ernst of London, a fortnight ago, to one of my cases, a young gentleman six years of age, from Ireland, and he was enabled to walk about with ease and comfort; but, as he was previously free from pain, the case was less a test of the value of extension than that previously alluded to.

These instruments and their mode of application are described by Dr. Sayre in his *Lectures on Orthopaedic Surgery*, published in 1876; and, as we have now the pleasure of welcoming Dr. Sayre amongst the distinguished guests present at this meeting of our Association, I am glad to state that he will exhibit the hip-joint instrument of his own construction, with the latest improvements, and make some observations on its application.

The accompanying woodcuts, which were kindly forwarded to me by Dr. L. Sayre of New York, for the purpose of illustrating this

Fig. 1.—Dr. Sayre's short Hip-Joint Splint.

Fig. 2.—Dr. Sayre's long Hip-Joint Splint.

paper, represent the two hip-joint instruments used by him for combining extension with motion in the treatment of hip-joint disease. Both these instruments are figured and their application described in Dr. Sayre's recently published *Lectures on Orthopaedic Surgery*, pp. 262 and 269.

ALCESTER.—A considerable portion of the report for the quarter ending September 30th was taken up with the details of the action taken for the prevention of the spread of scarlatina and typhoid. As regards the former, he regrets not having a disinfecting chamber, and relates a successful prosecution for the exposure of some children while in an infected state. As regards the latter—typhoid—Mr. Fosbroke relates the history of two outbreaks in some isolated cottages, which occurred at first from polluted water-supply arising from defective drains, and afterwards from the excreta of the infected having been thrown into the common privy without disinfection. The annual death-rate was only 14.9 per 1,000 living.